

North Providence (Providence)

| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE ³ | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|--|---|------------------------------------|---------------------------------|-------------------|---------------------------------|
| POPULATION CHARACTERISTICS | | | | | |
| % of 60+ LGBT | | 2.1% | (1.3% - 2.9%) | 2.0% | (1.6% - 2.4%) |
| Age-sex adjusted 1-year mortality rate | | 4.8% | (4.2% - 5.3%) | 4.8% | (4.6% - 4.9%) |
| % 60+ lived at same address 25 years or more | | 52.2% | (46.4% - 58.0%) | 48.1% | (46.8% - 49.5%) |
| WELLNESS and PREVENTION | | | | | |
| % any physical activity within last month | w | 63.6% | (59.1% - 68.0%) | 70.0% | (69.0% - 71.0%) |
| % injured by a fall within last year | | 13.4% | (7.2% - 19.6%) | 10.0% | (8.7% - 11.3%) |
| % ever had a hip fracture | | 3.6% | (2.9% - 4.3%) | 3.9% | (3.7% - 4.0%) |
| % with self-reported fair or poor health status | | 24.6% | (20.2% - 29.1%) | 20.4% | (19.4% - 21.5%) |
| % with 15+ physically unhealthy days last month | | 15.8% | (12.0% - 19.6%) | 13.9% | (13.0% - 14.8%) |
| % with physical exam/check-up in past year | | 92.4% | (89.7% - 95.0%) | 91.9% | (91.2% - 92.6%) |
| % met CDC preventive health screening goals | | 37.4% | (31.1% - 43.6%) | 39.5% | (38.0% - 41.1%) |
| % flu shot past year | | 58.2% | (52.7% - 63.6%) | 59.1% | (57.8% - 60.4%) |
| % pneumonia vaccine | | 73.3% | (67.5% - 79.0%) | 73.8% | (72.4% - 75.2%) |
| % shingles vaccine | | 24.1% | (17.9% - 30.4%) | 30.3% | (28.7% - 31.8%) |
| % cholesterol screening | | 90.2% | (87.3% - 93.1%) | 88.4% | (87.6% - 89.3%) |
| % mammogram within last 2 years (women) | | 80.8% | (75.1% - 86.4%) | 81.8% | (80.4% - 83.1%) |
| % colorectal cancer screening | | 71.4% | (66.4% - 76.4%) | 76.1% | (75.0% - 77.3%) |
| Oral Health | | | | | |
| % with complete tooth loss | w | 37.9% | (33.6% - 42.1%) | 32.4% | (31.4% - 33.4%) |
| % with annual dental exam | W | 65.4% | (60.4% - 70.4%) | 74.7% | (73.5% - 75.8%) |
| NUTRITION / DIET | | | | | |
| % with 5 or more servings of fruit or vegetables per day | | 21.2% | (17.1% - 25.2%) | 23.0% | (21.9% - 24.1%) |
| % obese | | 28.0% | (23.2% - 32.9%) | 25.4% | (24.2% - 26.5%) |
| % high cholesterol | W | 80.2% | (78.7% - 81.7%) | 78.0% | (77.7% - 78.3%) |
| % current smokers | | 9.7% | (6.9% - 12.5%) | 8.9% | (8.2% - 9.7%) |
| % excessive drinking | B | 5.4% | (3.3% - 7.5%) | 8.9% | (8.1% - 9.6%) |
| MENTAL HEALTH | | | | | |
| % with 15+ days poor mental health last month | | 7.8% | (5.4% - 10.3%) | 7.5% | (6.8% - 8.1%) |
| % 60+ talked with family or friends almost daily | | 75.7% | (70.2% - 81.3%) | 75.3% | (74.1% - 76.5%) |
| % ever diagnosed with depression | W | 33.4% | (31.7% - 35.2%) | 30.0% | (29.7% - 30.3%) |

| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE ³ | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|--|---|------------------------------------|---------------------------------|-------------------|---------------------------------|
| CHRONIC DISEASE | | | | | |
| % with Alzheimer's disease or related dementias | | 13.5% | (12.2% - 14.8%) | 14.4% | (14.2% - 14.7%) |
| % with diabetes | W | 42.4% | (40.5% - 44.3%) | 35.7% | (35.4% - 36.1%) |
| % with stroke | | 11.7% | (10.5% - 12.8%) | 12.5% | (12.2% - 12.7%) |
| % with chronic obstructive pulmonary disease | W | 27.7% | (26.0% - 29.3%) | 24.1% | (23.8% - 24.4%) |
| % with asthma | | 14.8% | (13.5% - 16.1%) | 14.0% | (13.7% - 14.2%) |
| % with hypertension | W | 82.1% | (80.6% - 83.5%) | 79.0% | (78.8% - 79.3%) |
| % ever had a heart attack | | 6.2% | (5.3% - 7.1%) | 5.4% | (5.3% - 5.6%) |
| % with ischemic heart disease | W | 48.5% | (46.5% - 50.4%) | 45.9% | (45.5% - 46.2%) |
| % with congestive heart failure | W | 27.0% | (25.3% - 28.7%) | 24.8% | (24.5% - 25.1%) |
| % with atrial fibrillation | | 14.8% | (13.5% - 16.1%) | 15.2% | (15.0% - 15.5%) |
| % with osteoarthritis/rheumatoid arthritis | W | 54.5% | (52.5% - 56.5%) | 52.0% | (51.6% - 52.4%) |
| % with osteoporosis | W | 23.0% | (21.5% - 24.6%) | 21.0% | (20.7% - 21.3%) |
| % with glaucoma | | 26.9% | (25.3% - 28.5%) | 26.6% | (26.2% - 26.9%) |
| % with cataract | | 68.6% | (66.9% - 70.3%) | 67.9% | (67.6% - 68.2%) |
| % women with breast cancer | | 12.0% | (10.4% - 13.5%) | 10.7% | (10.4% - 11.0%) |
| % with colon cancer | W | 4.1% | (3.4% - 4.8%) | 3.2% | (3.1% - 3.3%) |
| % men with prostate cancer | | 14.4% | (12.4% - 16.5%) | 13.8% | (13.4% - 14.1%) |
| % with lung cancer | | 1.7% | (1.2% - 2.2%) | 2.1% | (2.0% - 2.2%) |
| % with hypothyroidism | W | 23.1% | (21.5% - 24.6%) | 21.1% | (20.8% - 21.4%) |
| % with anemia | W | 58.5% | (56.7% - 60.3%) | 52.2% | (51.9% - 52.6%) |
| % with benign prostatic hyperplasia | W | 43.8% | (40.9% - 46.7%) | 40.3% | (39.8% - 40.8%) |
| % with chronic kidney disease | | 24.3% | (22.6% - 25.9%) | 23.3% | (23.0% - 23.6%) |
| Summary chronic disease measures | | | | | |
| % with 4+ chronic conditions | W | 69.0% | (67.3% - 70.8%) | 63.9% | (63.6% - 64.3%) |
| % with 0 chronic conditions | | 7.2% | (6.3% - 8.2%) | 8.4% | (8.2% - 8.6%) |
| ACCESS TO CARE | | | | | |
| % with a regular doctor | b | 97.9% | (97.0% - 98.8%) | 96.5% | (96.2% - 96.9%) |
| % did not see a doctor when needed due to cost | | 9.5% | (5.5% - 13.4%) | 6.3% | (5.6% - 6.9%) |
| SERVICE UTILIZATION | | | | | |
| Physician visits per year | * | 8.8 | (8.5 - 9.1) | 8.0 | (7.9 - 8.0) |
| Emergency room visits/1000 persons 65+ years per year | * | 708 | (648 - 767) | 628 | (618 - 638) |
| Part D monthly prescription fills per person per year | | 54.8 | (53.7 - 56.0) | 54.2 | (53.9 - 54.4) |
| Home health visits per year | | 4.3 | (3.6 - 4.9) | 3.7 | (3.6 - 3.9) |
| Durable medical equipment claims per year | * | 2.2 | (2.0 - 2.4) | 2.0 | (1.9 - 2.0) |
| Inpatient hospital stays/1000 persons 65+ years per year | | 303 | (271 - 334) | 284 | (279 - 289) |
| Inpatient hospital readmissions (as % of admissions) | * | 24.3% | (20.5% - 28.1%) | 16.9% | (16.3% - 17.5%) |
| Skilled nursing facility stays/1000 persons 65+ years per year | | 112 | (96 - 129) | 100 | (97 - 103) |
| COMMUNITY VARIABLE & CIVIC ENGAGEMENT | | | | | |
| % 60+ who are satisfied with neighborhood | | 75.7% | (71.1% - 80.3%) | 80.0% | (78.9% - 81.1%) |
| % 60+ who believe local service orgs understand needs | W | 32.2% | (26.6% - 37.8%) | 44.1% | (42.7% - 45.6%) |
| % 60+ who believe he/she can make a difference | W | 42.0% | (36.2% - 47.8%) | 51.6% | (50.2% - 53.0%) |
| % 60+ who believe working together can make a difference | | 81.0% | (76.9% - 85.2%) | 81.0% | (79.9% - 82.1%) |
| % 60+ who volunteer at least once per month | | 18.3% | (14.0% - 22.6%) | 22.8% | (21.6% - 23.9%) |
| % 60+ who attend community events at least once per month | w | 37.3% | (32.5% - 42.1%) | 44.2% | (43.1% - 45.4%) |

Notes. ¹ Community-state differences that the margins of error do not overlap each other are noted B, W, w, or *.

² All community and state margins of error are 95% confidence intervals, except when a lower case b or w is noted in which case 90% confidence intervals are used.

³ C indicates that the community rate is censored due to inadequate sample size.