

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02860 (Pawtucket)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02860 ESTIMATE	MARGIN OF ERROR ²	PAWTUCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			77.7%	(76.2% - 79.1%)	77.0%	(75.9% - 78.1%)	78.0%	(77.7% - 78.2%)
% with diabetes	w		42.5%	(40.7% - 44.4%)	39.7%	(38.4% - 41.1%)	35.7%	(35.4% - 36.0%)
% with stroke			12.5%	(11.4% - 13.7%)	12.8%	(11.9% - 13.6%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			23.9%	(22.4% - 25.4%)	23.4%	(22.3% - 24.5%)	24.1%	(23.8% - 24.3%)
% with asthma	w		16.8%	(15.5% - 18.2%)	16.3%	(15.3% - 17.2%)	14.0%	(13.8% - 14.2%)
% with hypertension	w		81.3%	(80.0% - 82.7%)	80.7%	(79.7% - 81.7%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.8%	(5.0% - 6.6%)	6.5%	(5.9% - 7.2%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			44.2%	(42.4% - 46.0%)	45.0%	(43.6% - 46.4%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			26.5%	(24.9% - 28.1%)	26.9%	(25.7% - 28.1%)	24.8%	(24.5% - 25.1%)
% with anemia			52.4%	(50.6% - 54.1%)	52.8%	(51.5% - 54.1%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		27.0%	(25.4% - 28.6%)	26.4%	(25.2% - 27.6%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		66.0%	(64.4% - 67.7%)	65.8%	(64.6% - 67.1%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			8.8%	(7.8% - 9.8%)	10.0%	(9.2% - 10.8%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w		711	(658 - 764)	662	(623 - 701)	628	(620 - 637)
Part D monthly prescription fills per person per year	w	w	59.3	(58.2 - 60.3)	57.2	(56.4 - 58.0)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.5	(2.3 - 2.7)	2.2	(2.0 - 2.3)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			295	(267 - 322)	291	(271 - 311)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			96	(82 - 109)	98	(88 - 108)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			13.9%	(12.7% - 15.0%)	14.9%	(14.0% - 15.9%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	b		46.5%	(44.7% - 48.3%)	47.5%	(46.1% - 48.9%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			20.7%	(19.3% - 22.1%)	21.3%	(20.2% - 22.3%)	21.0%	(20.8% - 21.2%)
% with glaucoma	b		24.7%	(23.2% - 26.2%)	24.4%	(23.3% - 25.5%)	26.6%	(26.3% - 26.8%)
% with cataract			66.2%	(64.6% - 67.9%)	65.0%	(63.7% - 66.2%)	67.9%	(67.6% - 68.2%)
% women with breast cancer	b		8.9%	(7.7% - 10.2%)	9.5%	(8.5% - 10.5%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer	b		10.8%	(9.0% - 12.5%)	11.4%	(10.1% - 12.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			20.7%	(19.3% - 22.1%)	21.1%	(20.0% - 22.2%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	b		36.3%	(33.6% - 38.9%)	37.3%	(35.3% - 39.2%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b		6.5	(6.3 - 6.7)	6.9	(6.7 - 7.1)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02860 ESTIMATE	MARGIN OF ERROR ²	PAWTUCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.3%	(3.8% - 4.8%)	4.7%	(4.3% - 5.1%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture	b		3.0%	(2.4% - 3.5%)	3.4%	(2.9% - 3.8%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		35.3%	(33.6% - 36.9%)	33.3%	(32.1% - 34.5%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			15.4%	(14.1% - 16.7%)	15.2%	(14.2% - 16.2%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.2%	(2.6% - 3.8%)	3.3%	(2.8% - 3.7%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.1%	(1.6% - 2.6%)	2.3%	(1.9% - 2.7%)	2.1%	(2.0% - 2.2%)
Home health visits per year			4.0	(3.5 - 4.4)	3.8	(3.5 - 4.2)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w	w	10.9%	(10.2% - 11.7%)	8.6%	(8.2% - 9.1%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			16.2%	(12.6% - 19.7%)	16.9%	(14.5% - 19.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			7.6%	(5.2% - 10.0%)	7.5%	(5.8% - 9.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		26.8%	(22.6% - 30.9%)	26.0%	(23.1% - 28.9%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty	w		11.8%	(8.5% - 15.2%)	10.9%	(8.6% - 13.3%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			7.7%	(5.5% - 9.9%)	7.9%	(6.2% - 9.6%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			17.5%	(14.1% - 21.0%)	17.2%	(14.7% - 19.8%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.