

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02861 (Pawtucket)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02861 ESTIMATE	MARGIN OF ERROR ²	PAWTUCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			76.1%	(74.4% - 77.8%)	77.0%	(75.9% - 78.1%)	78.0%	(77.7% - 78.2%)
% with diabetes		b	35.9%	(33.8% - 38.0%)	39.7%	(38.4% - 41.1%)	35.7%	(35.4% - 36.0%)
% with stroke			12.9%	(11.6% - 14.3%)	12.8%	(11.9% - 13.6%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			22.6%	(20.9% - 24.3%)	23.4%	(22.3% - 24.5%)	24.1%	(23.8% - 24.3%)
% with asthma			15.5%	(14.1% - 17.0%)	16.3%	(15.3% - 17.2%)	14.0%	(13.8% - 14.2%)
% with hypertension			79.6%	(78.0% - 81.3%)	80.7%	(79.7% - 81.7%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack	w		7.4%	(6.3% - 8.4%)	6.5%	(5.9% - 7.2%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			45.6%	(43.5% - 47.7%)	45.0%	(43.6% - 46.4%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		27.0%	(25.2% - 28.9%)	26.9%	(25.7% - 28.1%)	24.8%	(24.5% - 25.1%)
% with anemia			53.0%	(51.0% - 55.0%)	52.8%	(51.5% - 54.1%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			24.9%	(23.1% - 26.8%)	26.4%	(25.2% - 27.6%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions			65.2%	(63.3% - 67.1%)	65.8%	(64.6% - 67.1%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions	b		11.8%	(10.5% - 13.1%)	10.0%	(9.2% - 10.8%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			590	(533 - 647)	662	(623 - 701)	628	(620 - 637)
Part D monthly prescription fills per person per year			55.3	(54.1 - 56.5)	57.2	(56.4 - 58.0)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year		b	1.8	(1.6 - 2.0)	2.2	(2.0 - 2.3)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			282	(254 - 311)	291	(271 - 311)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			98	(83 - 113)	98	(88 - 108)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			16.1%	(14.7% - 17.6%)	14.9%	(14.0% - 15.9%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	b		48.5%	(46.3% - 50.7%)	47.5%	(46.1% - 48.9%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			22.1%	(20.5% - 23.7%)	21.3%	(20.2% - 22.3%)	21.0%	(20.8% - 21.2%)
% with glaucoma	b		23.8%	(22.1% - 25.5%)	24.4%	(23.3% - 25.5%)	26.6%	(26.3% - 26.8%)
% with cataract	b		62.9%	(60.9% - 64.8%)	65.0%	(63.7% - 66.2%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			10.4%	(8.8% - 12.0%)	9.5%	(8.5% - 10.5%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			11.9%	(9.9% - 13.9%)	11.4%	(10.1% - 12.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			21.7%	(20.1% - 23.3%)	21.1%	(20.0% - 22.2%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			37.5%	(34.5% - 40.5%)	37.3%	(35.3% - 39.2%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b	w	7.4	(7.1 - 7.7)	6.9	(6.7 - 7.1)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02861 ESTIMATE	MARGIN OF ERROR ²	PAWTUCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.8%	(4.3% - 5.4%)	4.7%	(4.3% - 5.1%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			3.8%	(3.1% - 4.6%)	3.4%	(2.9% - 3.8%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression			30.8%	(28.9% - 32.6%)	33.3%	(32.1% - 34.5%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			14.6%	(13.1% - 16.0%)	15.2%	(14.2% - 16.2%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.3%	(2.6% - 4.0%)	3.3%	(2.8% - 3.7%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.4%	(1.8% - 3.0%)	2.3%	(1.9% - 2.7%)	2.1%	(2.0% - 2.2%)
Home health visits per year			3.5	(3.0 - 4.1)	3.8	(3.5 - 4.2)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports		b	5.7%	(5.1% - 6.3%)	8.6%	(8.2% - 9.1%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty	w		18.0%	(14.4% - 21.6%)	16.9%	(14.5% - 19.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			7.4%	(4.6% - 10.2%)	7.5%	(5.8% - 9.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		25.0%	(20.7% - 29.3%)	26.0%	(23.1% - 28.9%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			9.7%	(6.6% - 12.8%)	10.9%	(8.6% - 13.3%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			8.3%	(5.6% - 11.0%)	7.9%	(6.2% - 9.6%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			16.8%	(13.3% - 20.4%)	17.2%	(14.7% - 19.8%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.