

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02889 (Warwick)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02889 ESTIMATE	MARGIN OF ERROR ²	WARWICK ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			78.4%	(76.8% - 79.9%)	80.0%	(79.2% - 80.8%)	78.0%	(77.7% - 78.2%)
% with diabetes	w		38.2%	(36.3% - 40.1%)	37.9%	(36.8% - 38.9%)	35.7%	(35.4% - 36.0%)
% with stroke	w		14.0%	(12.8% - 15.3%)	13.4%	(12.8% - 14.1%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			25.0%	(23.4% - 26.5%)	25.9%	(25.0% - 26.8%)	24.1%	(23.8% - 24.3%)
% with asthma			15.4%	(14.1% - 16.7%)	15.1%	(14.4% - 15.9%)	14.0%	(13.8% - 14.2%)
% with hypertension			79.4%	(77.9% - 80.9%)	80.6%	(79.8% - 81.5%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack	w		6.9%	(6.0% - 7.8%)	6.4%	(5.9% - 6.9%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease	w		50.4%	(48.4% - 52.4%)	50.5%	(49.4% - 51.6%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			26.5%	(24.8% - 28.2%)	26.4%	(25.5% - 27.4%)	24.8%	(24.5% - 25.1%)
% with anemia	w		55.4%	(53.6% - 57.2%)	57.3%	(56.2% - 58.3%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			23.7%	(22.0% - 25.4%)	25.1%	(24.1% - 26.0%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		66.2%	(64.5% - 68.0%)	67.7%	(66.8% - 68.7%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			9.0%	(8.0% - 10.1%)	7.9%	(7.3% - 8.4%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w		688	(637 - 738)	683	(647 - 718)	628	(620 - 637)
Part D monthly prescription fills per person per year			55.3	(54.2 - 56.4)	54.5	(53.8 - 55.1)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year			2.0	(1.8 - 2.2)	2.1	(2.0 - 2.2)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			306	(278 - 334)	306	(290 - 322)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			97	(82 - 111)	97	(89 - 105)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			14.9%	(13.6% - 16.2%)	15.3%	(14.5% - 16.0%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	w		55.1%	(53.1% - 57.1%)	55.6%	(54.5% - 56.7%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			20.5%	(19.1% - 22.0%)	22.8%	(22.0% - 23.7%)	21.0%	(20.8% - 21.2%)
% with glaucoma	b		23.7%	(22.1% - 25.2%)	25.9%	(25.0% - 26.8%)	26.6%	(26.3% - 26.8%)
% with cataract	b		65.3%	(63.5% - 67.0%)	67.4%	(66.4% - 68.4%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			10.9%	(9.4% - 12.4%)	10.8%	(10.0% - 11.6%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			13.4%	(11.5% - 15.3%)	13.8%	(12.7% - 14.8%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			22.3%	(20.7% - 23.8%)	23.7%	(22.9% - 24.6%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	b		35.7%	(33.0% - 38.3%)	38.5%	(37.0% - 40.0%)	40.3%	(39.9% - 40.7%)
Physician visits per year			8.0	(7.7 - 8.2)	8.4	(8.2 - 8.5)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02889 ESTIMATE	MARGIN OF ERROR ²	WARWICK ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			5.4%	(4.8% - 5.9%)	5.1%	(4.8% - 5.4%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			4.0%	(3.3% - 4.8%)	4.3%	(3.9% - 4.7%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		32.7%	(31.0% - 34.4%)	32.3%	(31.3% - 33.2%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			13.6%	(12.4% - 14.9%)	13.7%	(13.0% - 14.5%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.5%	(2.9% - 4.2%)	3.5%	(3.1% - 3.8%)	3.2%	(3.1% - 3.3%)
% with lung cancer	w		3.1%	(2.4% - 3.7%)	2.8%	(2.4% - 3.1%)	2.1%	(2.0% - 2.2%)
Home health visits per year			3.9	(3.4 - 4.5)	3.7	(3.4 - 4.0)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	b		5.5%	(4.9% - 6.0%)	5.1%	(4.8% - 5.4%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			17.0%	(13.4% - 20.5%)	14.1%	(12.3% - 16.0%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			5.6%	(3.4% - 7.9%)	6.0%	(4.6% - 7.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			17.0%	(13.8% - 20.3%)	18.9%	(16.7% - 21.1%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			10.3%	(7.1% - 13.5%)	8.3%	(6.7% - 9.8%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			5.4%	(3.1% - 7.8%)	6.0%	(4.7% - 7.3%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			13.6%	(10.7% - 16.4%)	13.3%	(11.3% - 15.2%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.