

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02895 (Woonsocket)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02895 ESTIMATE	MARGIN OF ERROR ²	WOONSOCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	w		80.8%	(79.4% - 82.2%)	80.8%	(79.4% - 82.2%)	78.0%	(77.7% - 78.2%)
% with diabetes	w		40.2%	(38.4% - 42.0%)	40.2%	(38.4% - 42.0%)	35.7%	(35.4% - 36.0%)
% with stroke			13.5%	(12.4% - 14.7%)	13.5%	(12.4% - 14.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease	w		34.5%	(32.9% - 36.1%)	34.5%	(32.9% - 36.1%)	24.1%	(23.8% - 24.3%)
% with asthma	w		17.2%	(15.9% - 18.4%)	17.2%	(15.9% - 18.4%)	14.0%	(13.8% - 14.2%)
% with hypertension	w		83.8%	(82.6% - 85.1%)	83.8%	(82.6% - 85.1%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			6.4%	(5.5% - 7.2%)	6.4%	(5.5% - 7.2%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease	w		58.8%	(57.0% - 60.6%)	58.8%	(57.0% - 60.6%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		34.4%	(32.7% - 36.1%)	34.4%	(32.7% - 36.1%)	24.8%	(24.5% - 25.1%)
% with anemia	w		54.5%	(52.8% - 56.2%)	54.5%	(52.8% - 56.2%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			23.6%	(22.1% - 25.2%)	23.6%	(22.1% - 25.2%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		70.8%	(69.3% - 72.4%)	70.8%	(69.3% - 72.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			7.7%	(6.8% - 8.7%)	7.7%	(6.8% - 8.7%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w		889	(830 - 949)	889	(830 - 949)	628	(620 - 637)
Part D monthly prescription fills per person per year	w		61.8	(60.7 - 63.0)	61.8	(60.7 - 63.0)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.6	(2.4 - 2.8)	2.6	(2.4 - 2.8)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year	w		428	(394 - 461)	428	(394 - 461)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year	w		135	(119 - 152)	135	(119 - 152)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation	w		17.6%	(16.3% - 18.8%)	17.6%	(16.3% - 18.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis			53.3%	(51.4% - 55.1%)	53.3%	(51.4% - 55.1%)	52.0%	(51.7% - 52.3%)
% with osteoporosis	w		23.5%	(22.0% - 24.9%)	23.5%	(22.0% - 24.9%)	21.0%	(20.8% - 21.2%)
% with glaucoma	b		24.2%	(22.7% - 25.7%)	24.2%	(22.7% - 25.7%)	26.6%	(26.3% - 26.8%)
% with cataract			68.1%	(66.5% - 69.7%)	68.1%	(66.5% - 69.7%)	67.9%	(67.6% - 68.2%)
% women with breast cancer	b		9.0%	(7.7% - 10.3%)	9.0%	(7.7% - 10.3%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer	b		11.0%	(9.4% - 12.7%)	11.0%	(9.4% - 12.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism	w		24.0%	(22.6% - 25.5%)	24.0%	(22.6% - 25.5%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			39.7%	(37.1% - 42.3%)	39.7%	(37.1% - 42.3%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b		7.6	(7.4 - 7.9)	7.6	(7.4 - 7.9)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02895 ESTIMATE	MARGIN OF ERROR ²	WOONSOCKET ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate	w		5.6%	(5.1% - 6.2%)	5.6%	(5.1% - 6.2%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			4.1%	(3.4% - 4.8%)	4.1%	(3.4% - 4.8%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		32.0%	(30.4% - 33.6%)	32.0%	(30.4% - 33.6%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			15.3%	(14.0% - 16.5%)	15.3%	(14.0% - 16.5%)	14.4%	(14.2% - 14.6%)
% with colon cancer	w		4.5%	(3.8% - 5.2%)	4.5%	(3.8% - 5.2%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.2%	(1.7% - 2.7%)	2.2%	(1.7% - 2.7%)	2.1%	(2.0% - 2.2%)
Home health visits per year			3.8	(3.3 - 4.4)	3.8	(3.3 - 4.4)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w		11.5%	(10.8% - 12.2%)	11.5%	(10.8% - 12.2%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			16.6%	(13.2% - 20.0%)	16.6%	(13.2% - 20.0%)	13.8%	(13.2% - 14.2%)
% 65+ with vision difficulty			4.4%	(2.9% - 5.8%)	4.4%	(2.9% - 5.8%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		25.1%	(21.6% - 28.6%)	25.1%	(21.6% - 28.6%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			8.5%	(6.0% - 10.9%)	8.5%	(6.0% - 10.9%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			5.6%	(3.2% - 8.0%)	5.6%	(3.2% - 8.0%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			17.2%	(14.1% - 20.3%)	17.2%	(14.1% - 20.3%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.