

## RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

## 02903 (Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	BETTER / WORSE CITY RATE <sup>1</sup>	02903 ESTIMATE	MARGIN OF ERROR <sup>2</sup>	PROVIDENCE ESTIMATE	MARGIN OF ERROR <sup>2</sup>	STATE ESTIMATE	MARGIN OF ERROR <sup>2</sup>
<b>SERIOUS CHRONIC DISEASE</b>								
% high cholesterol			75.2%	( 72.5% - 77.9% )	75.1%	( 74.2% - 75.9% )	78.0%	( 77.7% - 78.2% )
% with diabetes	w		41.8%	( 38.5% - 45.0% )	40.4%	( 39.4% - 41.4% )	35.7%	( 35.4% - 36.0% )
% with stroke			11.3%	( 9.3% - 13.3% )	12.3%	( 11.7% - 13.0% )	12.5%	( 12.3% - 12.6% )
% with chronic obstructive pulmonary disease			24.9%	( 22.2% - 27.6% )	22.7%	( 21.9% - 23.5% )	24.1%	( 23.8% - 24.3% )
% with asthma	w		17.5%	( 15.1% - 19.8% )	15.3%	( 14.6% - 16.0% )	14.0%	( 13.8% - 14.2% )
% with hypertension			79.1%	( 76.5% - 81.6% )	78.6%	( 77.8% - 79.4% )	79.0%	( 78.8% - 79.3% )
% ever had a heart attack			4.5%	( 3.2% - 5.8% )	4.9%	( 4.5% - 5.3% )	5.4%	( 5.3% - 5.6% )
% with ischemic heart disease		w	49.3%	( 46.0% - 52.6% )	44.9%	( 43.9% - 45.9% )	45.9%	( 45.5% - 46.2% )
% with congestive heart failure	w		29.6%	( 26.6% - 32.6% )	25.8%	( 24.9% - 26.6% )	24.8%	( 24.5% - 25.1% )
% with anemia			54.6%	( 51.5% - 57.7% )	52.5%	( 51.5% - 53.4% )	52.2%	( 51.9% - 52.5% )
% with chronic kidney disease	w		29.4%	( 26.4% - 32.4% )	26.6%	( 25.7% - 27.5% )	23.3%	( 23.1% - 23.6% )
% with 4+ chronic conditions	w		68.3%	( 65.4% - 71.3% )	64.5%	( 63.6% - 65.4% )	63.9%	( 63.6% - 64.2% )
% with 0 chronic conditions			7.9%	( 6.2% - 9.6% )	9.5%	( 9.0% - 10.1% )	8.4%	( 8.2% - 8.6% )
Emergency room visits/1000 persons 65+ years per year	w		916	( 756 - 1077 )	744	( 712 - 776 )	628	( 620 - 637 )
Part D monthly prescription fills per person per year	w	w	62.5	( 60.2 - 64.8 )	56.9	( 56.2 - 57.6 )	54.2	( 54.0 - 54.4 )
Durable medical equipment claims per year	w		2.5	( 2.1 - 2.9 )	2.4	( 2.3 - 2.5 )	2.0	( 1.9 - 2.0 )
Inpatient hospital stays/1000 persons 65+ years per year			297	( 246 - 348 )	285	( 270 - 300 )	284	( 280 - 289 )
Skilled nursing facility stays/1000 persons 65+ years per year			98	( 73 - 122 )	99	( 91 - 107 )	100	( 98 - 103 )
<b>MILD CHRONIC DISEASE</b>								
% with atrial fibrillation			14.0%	( 11.8% - 16.1% )	12.7%	( 12.1% - 13.4% )	15.2%	( 15.0% - 15.4% )
% with osteoarthritis/rheumatoid arthritis			51.0%	( 47.6% - 54.3% )	49.3%	( 48.2% - 50.3% )	52.0%	( 51.7% - 52.3% )
% with osteoporosis			22.1%	( 19.5% - 24.7% )	22.4%	( 21.6% - 23.2% )	21.0%	( 20.8% - 21.2% )
% with glaucoma			25.4%	( 22.7% - 28.1% )	26.5%	( 25.7% - 27.3% )	26.6%	( 26.3% - 26.8% )
% with cataract			64.8%	( 61.8% - 67.8% )	64.7%	( 63.8% - 65.6% )	67.9%	( 67.6% - 68.2% )
% women with breast cancer			10.7%	( 8.1% - 13.3% )	9.5%	( 8.8% - 10.3% )	10.7%	( 10.4% - 10.9% )
% men with prostate cancer			12.5%	( 9.4% - 15.6% )	13.0%	( 12.1% - 14.0% )	13.8%	( 13.5% - 14.1% )
% with hypothyroidism	b		18.4%	( 16.0% - 20.8% )	18.9%	( 18.1% - 19.6% )	21.1%	( 20.9% - 21.4% )
% with benign prostatic hyperplasia			36.3%	( 31.7% - 40.8% )	38.9%	( 37.5% - 40.3% )	40.3%	( 39.9% - 40.7% )
Physician visits per year	b		7.1	( 6.6 - 7.5 )	6.8	( 6.7 - 6.9 )	8.0	( 7.9 - 8.0 )

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	BETTER / WORSE CITY RATE <sup>1</sup>	02903 ESTIMATE	MARGIN OF ERROR <sup>2</sup>	PROVIDENCE ESTIMATE	MARGIN OF ERROR <sup>2</sup>	STATE ESTIMATE	MARGIN OF ERROR <sup>2</sup>
<b>DISABILITY and LATER LIFE CHRONIC DISEASE</b>								
Age-sex adjusted 1-year mortality rate			4.8%	( 3.8% - 5.9% )	5.0%	( 4.7% - 5.3% )	4.8%	( 4.7% - 4.8% )
% ever had a hip fracture			3.0%	( 2.0% - 4.0% )	3.8%	( 3.5% - 4.2% )	3.9%	( 3.8% - 4.0% )
% ever diagnosed with depression	w		36.6%	( 33.6% - 39.6% )	34.0%	( 33.1% - 34.9% )	30.0%	( 29.8% - 30.3% )
% with Alzheimer's disease or related dementias			16.7%	( 14.3% - 19.1% )	16.6%	( 15.9% - 17.3% )	14.4%	( 14.2% - 14.6% )
% with colon cancer			3.3%	( 2.2% - 4.3% )	2.7%	( 2.4% - 3.0% )	3.2%	( 3.1% - 3.3% )
% with lung cancer			2.3%	( 1.4% - 3.3% )	2.1%	( 1.8% - 2.3% )	2.1%	( 2.0% - 2.2% )
Home health visits per year			3.7	( 3.0 - 4.4 )	4.4	( 4.1 - 4.8 )	3.7	( 3.7 - 3.8 )
% of 65+ getting Medicaid long term services and supports	w	w	15.2%	( 13.6% - 16.8% )	11.3%	( 10.9% - 11.7% )	6.2%	( 6.1% - 6.3% )
% 65+ with hearing difficulty			13.8%	( 7.6% - 20.0% )	13.5%	( 11.7% - 15.4% )	13.8%	( 13.2% - 14.3% )
% 65+ with vision difficulty			6.1%	( 1.9% - 10.3% )	7.3%	( 5.8% - 8.8% )	5.2%	( 4.9% - 5.5% )
% 65+ with ambulatory difficulty			25.2%	( 18.3% - 32.1% )	23.6%	( 21.3% - 25.8% )	19.9%	( 19.3% - 20.5% )
% 65+ with cognition difficulty			11.7%	( 6.6% - 16.9% )	9.3%	( 7.8% - 10.8% )	7.8%	( 7.3% - 8.2% )
% 65+ with self-care difficulty			11.6%	( 6.3% - 16.8% )	8.7%	( 7.2% - 10.2% )	6.6%	( 6.2% - 6.9% )
% 65+ with independent living difficulty			20.1%	( 12.7% - 27.4% )	17.9%	( 15.7% - 20.1% )	13.7%	( 13.2% - 14.2% )

Notes.<sup>1</sup> Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

<sup>2</sup> All zip code, city, and state margins of error are 90% confidence intervals.