

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02904 (Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02904 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol		w	79.4%	(78.0% - 80.7%)	75.1%	(74.2% - 75.9%)	78.0%	(77.7% - 78.2%)
% with diabetes	w	w	43.3%	(41.5% - 45.1%)	40.4%	(39.4% - 41.4%)	35.7%	(35.4% - 36.0%)
% with stroke			11.7%	(10.7% - 12.8%)	12.3%	(11.7% - 13.0%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease	w	w	29.2%	(27.7% - 30.8%)	22.7%	(21.9% - 23.5%)	24.1%	(23.8% - 24.3%)
% with asthma	w		15.7%	(14.5% - 16.9%)	15.3%	(14.6% - 16.0%)	14.0%	(13.8% - 14.2%)
% with hypertension	w	w	82.8%	(81.5% - 84.0%)	78.6%	(77.8% - 79.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.8%	(5.0% - 6.5%)	4.9%	(4.5% - 5.3%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease	w	w	49.4%	(47.6% - 51.2%)	44.9%	(43.9% - 45.9%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		27.7%	(26.1% - 29.3%)	25.8%	(24.9% - 26.6%)	24.8%	(24.5% - 25.1%)
% with anemia	w	w	59.2%	(57.5% - 60.8%)	52.5%	(51.5% - 53.4%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		26.1%	(24.6% - 27.7%)	26.6%	(25.7% - 27.5%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w	w	70.4%	(68.8% - 71.9%)	64.5%	(63.6% - 65.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions	w	w	6.9%	(6.1% - 7.8%)	9.5%	(9.0% - 10.1%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w		764	(702 - 825)	744	(712 - 776)	628	(620 - 637)
Part D monthly prescription fills per person per year	w		57.1	(56.0 - 58.1)	56.9	(56.2 - 57.6)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.5	(2.3 - 2.7)	2.4	(2.3 - 2.5)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			315	(285 - 345)	285	(270 - 300)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year	w		119	(103 - 134)	99	(91 - 107)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			14.0%	(12.9% - 15.2%)	12.7%	(12.1% - 13.4%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	w	w	55.0%	(53.2% - 56.8%)	49.3%	(48.2% - 50.3%)	52.0%	(51.7% - 52.3%)
% with osteoporosis	w		24.2%	(22.7% - 25.6%)	22.4%	(21.6% - 23.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			26.3%	(24.9% - 27.8%)	26.5%	(25.7% - 27.3%)	26.6%	(26.3% - 26.8%)
% with cataract		w	67.9%	(66.3% - 69.5%)	64.7%	(63.8% - 65.6%)	67.9%	(67.6% - 68.2%)
% women with breast cancer	w	w	12.8%	(11.4% - 14.3%)	9.5%	(8.8% - 10.3%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			15.0%	(13.1% - 16.9%)	13.0%	(12.1% - 14.0%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism		w	21.9%	(20.5% - 23.3%)	18.9%	(18.1% - 19.6%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	w	w	45.8%	(43.2% - 48.5%)	38.9%	(37.5% - 40.3%)	40.3%	(39.9% - 40.7%)
Physician visits per year	w	w	8.3	(8.1 - 8.6)	6.8	(6.7 - 6.9)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02904 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.7%	(4.2% - 5.2%)	5.0%	(4.7% - 5.3%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			3.6%	(3.0% - 4.3%)	3.8%	(3.5% - 4.2%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		35.3%	(33.6% - 36.9%)	34.0%	(33.1% - 34.9%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias		b	14.6%	(13.3% - 15.8%)	16.6%	(15.9% - 17.3%)	14.4%	(14.2% - 14.6%)
% with colon cancer	w	w	4.0%	(3.4% - 4.7%)	2.7%	(2.4% - 3.0%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.0%	(1.5% - 2.4%)	2.1%	(1.8% - 2.3%)	2.1%	(2.0% - 2.2%)
Home health visits per year			4.1	(3.5 - 4.6)	4.4	(4.1 - 4.8)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w		10.3%	(9.6% - 11.0%)	11.3%	(10.9% - 11.7%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			12.9%	(9.3% - 16.4%)	13.5%	(11.7% - 15.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			6.7%	(3.7% - 9.6%)	7.3%	(5.8% - 8.8%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		25.4%	(20.8% - 30.0%)	23.6%	(21.3% - 25.8%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			9.1%	(5.7% - 12.4%)	9.3%	(7.8% - 10.8%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			10.1%	(6.4% - 13.7%)	8.7%	(7.2% - 10.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			16.0%	(12.0% - 20.0%)	17.9%	(15.7% - 20.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.