

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02906 (Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02906 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	b		73.7% (71.8% - 75.5%)		75.1% (74.2% - 75.9%)		78.0% (77.7% - 78.2%)	
% with diabetes	b	b	25.2% (23.4% - 27.1%)		40.4% (39.4% - 41.4%)		35.7% (35.4% - 36.0%)	
% with stroke			12.1% (10.9% - 13.4%)		12.3% (11.7% - 13.0%)		12.5% (12.3% - 12.6%)	
% with chronic obstructive pulmonary disease	b	b	16.5% (15.1% - 17.9%)		22.7% (21.9% - 23.5%)		24.1% (23.8% - 24.3%)	
% with asthma			13.4% (12.1% - 14.8%)		15.3% (14.6% - 16.0%)		14.0% (13.8% - 14.2%)	
% with hypertension	b	b	70.3% (68.4% - 72.2%)		78.6% (77.8% - 79.4%)		79.0% (78.8% - 79.3%)	
% ever had a heart attack	b	b	3.5% (2.8% - 4.1%)		4.9% (4.5% - 5.3%)		5.4% (5.3% - 5.6%)	
% with ischemic heart disease	b	b	39.0% (36.9% - 41.1%)		44.9% (43.9% - 45.9%)		45.9% (45.5% - 46.2%)	
% with congestive heart failure	b	b	18.1% (16.6% - 19.6%)		25.8% (24.9% - 26.6%)		24.8% (24.5% - 25.1%)	
% with anemia	b	b	48.0% (46.0% - 50.0%)		52.5% (51.5% - 53.4%)		52.2% (51.9% - 52.5%)	
% with chronic kidney disease		b	21.7% (20.0% - 23.4%)		26.6% (25.7% - 27.5%)		23.3% (23.1% - 23.6%)	
% with 4+ chronic conditions	b	b	56.7% (54.7% - 58.7%)		64.5% (63.6% - 65.4%)		63.9% (63.6% - 64.2%)	
% with 0 chronic conditions			9.2% (7.9% - 10.5%)		9.5% (9.0% - 10.1%)		8.4% (8.2% - 8.6%)	
Emergency room visits/1000 persons 65+ years per year	b	b	516 (473 - 559)		744 (712 - 776)		628 (620 - 637)	
Part D monthly prescription fills per person per year	b	b	48.5 (47.2 - 49.8)		56.9 (56.2 - 57.6)		54.2 (54.0 - 54.4)	
Durable medical equipment claims per year	b	b	1.6 (1.4 - 1.8)		2.4 (2.3 - 2.5)		2.0 (1.9 - 2.0)	
Inpatient hospital stays/1000 persons 65+ years per year	b	b	202 (179 - 225)		285 (270 - 300)		284 (280 - 289)	
Skilled nursing facility stays/1000 persons 65+ years per year	b	b	75 (61 - 90)		99 (91 - 107)		100 (98 - 103)	
MILD CHRONIC DISEASE								
% with atrial fibrillation			14.1% (12.7% - 15.4%)		12.7% (12.1% - 13.4%)		15.2% (15.0% - 15.4%)	
% with osteoarthritis/rheumatoid arthritis	b		48.6% (46.4% - 50.8%)		49.3% (48.2% - 50.3%)		52.0% (51.7% - 52.3%)	
% with osteoporosis	w		24.4% (22.7% - 26.0%)		22.4% (21.6% - 23.2%)		21.0% (20.8% - 21.2%)	
% with glaucoma	w		28.7% (27.0% - 30.5%)		26.5% (25.7% - 27.3%)		26.6% (26.3% - 26.8%)	
% with cataract		w	69.8% (67.9% - 71.8%)		64.7% (63.8% - 65.6%)		67.9% (67.6% - 68.2%)	
% women with breast cancer	w	w	12.7% (11.1% - 14.4%)		9.5% (8.8% - 10.3%)		10.7% (10.4% - 10.9%)	
% men with prostate cancer			15.1% (13.0% - 17.3%)		13.0% (12.1% - 14.0%)		13.8% (13.5% - 14.1%)	
% with hypothyroidism			19.9% (18.3% - 21.4%)		18.9% (18.1% - 19.6%)		21.1% (20.9% - 21.4%)	
% with benign prostatic hyperplasia			42.3% (39.2% - 45.4%)		38.9% (37.5% - 40.3%)		40.3% (39.9% - 40.7%)	
Physician visits per year		w	8.1 (7.8 - 8.4)		6.8 (6.7 - 6.9)		8.0 (7.9 - 8.0)	

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02906 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate	b	b	4.0%	(3.4% - 4.6%)	5.0%	(4.7% - 5.3%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			4.1%	(3.4% - 4.8%)	3.8%	(3.5% - 4.2%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		33.2%	(31.4% - 35.1%)	34.0%	(33.1% - 34.9%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			14.8%	(13.4% - 16.2%)	16.6%	(15.9% - 17.3%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.1%	(2.5% - 3.7%)	2.7%	(2.4% - 3.0%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.0%	(1.5% - 2.5%)	2.1%	(1.8% - 2.3%)	2.1%	(2.0% - 2.2%)
Home health visits per year		b	3.3	(2.8 - 3.8)	4.4	(4.1 - 4.8)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	b	b	3.8%	(3.2% - 4.3%)	11.3%	(10.9% - 11.7%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			13.8%	(9.2% - 18.5%)	13.5%	(11.7% - 15.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			4.0%	(2.2% - 5.9%)	7.3%	(5.8% - 8.8%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	b	b	14.2%	(10.6% - 17.8%)	23.6%	(21.3% - 25.8%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty	b	b	3.5%	(1.7% - 5.3%)	9.3%	(7.8% - 10.8%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			6.5%	(3.4% - 9.7%)	8.7%	(7.2% - 10.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			13.6%	(9.6% - 17.5%)	17.9%	(15.7% - 20.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.