

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02908 (Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02908 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	b		75.3%	(73.5% - 77.2%)	75.1%	(74.2% - 75.9%)	78.0%	(77.7% - 78.2%)
% with diabetes	w		41.6%	(39.3% - 43.9%)	40.4%	(39.4% - 41.4%)	35.7%	(35.4% - 36.0%)
% with stroke			12.7%	(11.3% - 14.1%)	12.3%	(11.7% - 13.0%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			25.1%	(23.2% - 26.9%)	22.7%	(21.9% - 23.5%)	24.1%	(23.8% - 24.3%)
% with asthma			14.7%	(13.3% - 16.2%)	15.3%	(14.6% - 16.0%)	14.0%	(13.8% - 14.2%)
% with hypertension			79.4%	(77.6% - 81.1%)	78.6%	(77.8% - 79.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.6%	(4.7% - 6.6%)	4.9%	(4.5% - 5.3%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			47.1%	(44.8% - 49.4%)	44.9%	(43.9% - 45.9%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		28.5%	(26.5% - 30.5%)	25.8%	(24.9% - 26.6%)	24.8%	(24.5% - 25.1%)
% with anemia			54.6%	(52.5% - 56.7%)	52.5%	(51.5% - 53.4%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		26.9%	(24.9% - 28.9%)	26.6%	(25.7% - 27.5%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		66.6%	(64.6% - 68.7%)	64.5%	(63.6% - 65.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			9.8%	(8.5% - 11.1%)	9.5%	(9.0% - 10.1%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w		720	(660 - 779)	744	(712 - 776)	628	(620 - 637)
Part D monthly prescription fills per person per year	w		56.7	(55.2 - 58.2)	56.9	(56.2 - 57.6)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.4	(2.1 - 2.6)	2.4	(2.3 - 2.5)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			284	(253 - 315)	285	(270 - 300)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			118	(99 - 137)	99	(91 - 107)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation		w	15.0%	(13.5% - 16.4%)	12.7%	(12.1% - 13.4%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis			51.9%	(49.6% - 54.2%)	49.3%	(48.2% - 50.3%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			21.9%	(20.2% - 23.6%)	22.4%	(21.6% - 23.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			27.3%	(25.4% - 29.1%)	26.5%	(25.7% - 27.3%)	26.6%	(26.3% - 26.8%)
% with cataract	b		65.5%	(63.5% - 67.6%)	64.7%	(63.8% - 65.6%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			9.3%	(7.7% - 10.8%)	9.5%	(8.8% - 10.3%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			13.7%	(11.4% - 16.0%)	13.0%	(12.1% - 14.0%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			21.1%	(19.4% - 22.8%)	18.9%	(18.1% - 19.6%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			40.4%	(37.1% - 43.7%)	38.9%	(37.5% - 40.3%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b		6.9	(6.6 - 7.2)	6.8	(6.7 - 6.9)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02908 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate	w		5.8%	(5.2% - 6.5%)	5.0%	(4.7% - 5.3%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture	w	w	5.5%	(4.6% - 6.5%)	3.8%	(3.5% - 4.2%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		33.8%	(31.8% - 35.8%)	34.0%	(33.1% - 34.9%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias	w		18.5%	(16.8% - 20.2%)	16.6%	(15.9% - 17.3%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.3%	(2.6% - 4.1%)	2.7%	(2.4% - 3.0%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.0%	(1.5% - 2.6%)	2.1%	(1.8% - 2.3%)	2.1%	(2.0% - 2.2%)
Home health visits per year	w		5.2	(4.3 - 6.1)	4.4	(4.1 - 4.8)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w		11.2%	(10.3% - 12.1%)	11.3%	(10.9% - 11.7%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			16.1%	(11.8% - 20.4%)	13.5%	(11.7% - 15.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			8.7%	(5.2% - 12.3%)	7.3%	(5.8% - 8.8%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		26.0%	(20.9% - 31.1%)	23.6%	(21.3% - 25.8%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			8.7%	(5.5% - 12.0%)	9.3%	(7.8% - 10.8%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			6.7%	(3.8% - 9.6%)	8.7%	(7.2% - 10.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty	w		21.4%	(16.6% - 26.1%)	17.9%	(15.7% - 20.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.