

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02909 (Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02909 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			76.3%	(74.3% - 78.3%)	75.1%	(74.2% - 75.9%)	78.0%	(77.7% - 78.2%)
% with diabetes	w	w	46.5%	(44.0% - 49.0%)	40.4%	(39.4% - 41.4%)	35.7%	(35.4% - 36.0%)
% with stroke			12.2%	(10.7% - 13.8%)	12.3%	(11.7% - 13.0%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			24.8%	(22.8% - 26.8%)	22.7%	(21.9% - 23.5%)	24.1%	(23.8% - 24.3%)
% with asthma	w		16.3%	(14.6% - 18.0%)	15.3%	(14.6% - 16.0%)	14.0%	(13.8% - 14.2%)
% with hypertension			80.5%	(78.7% - 82.4%)	78.6%	(77.8% - 79.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.6%	(4.5% - 6.7%)	4.9%	(4.5% - 5.3%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			46.3%	(43.8% - 48.8%)	44.9%	(43.9% - 45.9%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		28.7%	(26.5% - 31.0%)	25.8%	(24.9% - 26.6%)	24.8%	(24.5% - 25.1%)
% with anemia			53.9%	(51.6% - 56.2%)	52.5%	(51.5% - 53.4%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		28.9%	(26.6% - 31.1%)	26.6%	(25.7% - 27.5%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions			65.4%	(63.2% - 67.6%)	64.5%	(63.6% - 65.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions	b		10.1%	(8.8% - 11.5%)	9.5%	(9.0% - 10.1%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	w	w	943	(855 - 1032)	744	(712 - 776)	628	(620 - 637)
Part D monthly prescription fills per person per year	w		58.7	(57.2 - 60.3)	56.9	(56.2 - 57.6)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.7	(2.4 - 3.0)	2.4	(2.3 - 2.5)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year	w	w	375	(332 - 419)	285	(270 - 300)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			114	(91 - 136)	99	(91 - 107)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation	b		11.1%	(9.6% - 12.5%)	12.7%	(12.1% - 13.4%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	b		47.2%	(44.7% - 49.7%)	49.3%	(48.2% - 50.3%)	52.0%	(51.7% - 52.3%)
% with osteoporosis		b	19.8%	(17.9% - 21.6%)	22.4%	(21.6% - 23.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			25.7%	(23.6% - 27.7%)	26.5%	(25.7% - 27.3%)	26.6%	(26.3% - 26.8%)
% with cataract	b	b	60.7%	(58.4% - 62.9%)	64.7%	(63.8% - 65.6%)	67.9%	(67.6% - 68.2%)
% women with breast cancer	b		7.4%	(5.8% - 9.1%)	9.5%	(8.8% - 10.3%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer	b	b	9.5%	(7.5% - 11.5%)	13.0%	(12.1% - 14.0%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism	b		17.1%	(15.3% - 18.8%)	18.9%	(18.1% - 19.6%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	b		35.5%	(32.3% - 38.8%)	38.9%	(37.5% - 40.3%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b	b	6.2	(5.9 - 6.5)	6.8	(6.7 - 6.9)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02909 ESTIMATE	MARGIN OF ERROR ²	PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.8%	(4.1% - 5.6%)	5.0%	(4.7% - 5.3%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture	b	b	2.6%	(1.9% - 3.4%)	3.8%	(3.5% - 4.2%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		32.9%	(30.7% - 35.1%)	34.0%	(33.1% - 34.9%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			15.5%	(13.7% - 17.3%)	16.6%	(15.9% - 17.3%)	14.4%	(14.2% - 14.6%)
% with colon cancer	b		2.3%	(1.6% - 3.0%)	2.7%	(2.4% - 3.0%)	3.2%	(3.1% - 3.3%)
% with lung cancer			1.7%	(1.1% - 2.3%)	2.1%	(1.8% - 2.3%)	2.1%	(2.0% - 2.2%)
Home health visits per year	w		5.6	(4.6 - 6.6)	4.4	(4.1 - 4.8)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w		11.1%	(10.1% - 12.1%)	11.3%	(10.9% - 11.7%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty	b		9.6%	(6.4% - 12.9%)	13.5%	(11.7% - 15.4%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			5.5%	(2.6% - 8.5%)	7.3%	(5.8% - 8.8%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty	w		27.6%	(21.8% - 33.4%)	23.6%	(21.3% - 25.8%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			11.5%	(7.6% - 15.4%)	9.3%	(7.8% - 10.8%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			9.4%	(6.4% - 12.4%)	8.7%	(7.2% - 10.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			18.2%	(13.5% - 22.8%)	17.9%	(15.7% - 20.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.