

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02910 (Cranston)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02910 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			79.0%	(76.5% - 81.4%)	78.9%	(78.0% - 79.8%)	78.0%	(77.7% - 78.2%)
% with diabetes			35.1%	(32.1% - 38.1%)	36.9%	(35.8% - 38.0%)	35.7%	(35.4% - 36.0%)
% with stroke	b	b	9.6%	(7.9% - 11.3%)	12.1%	(11.4% - 12.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			23.7%	(21.2% - 26.2%)	25.0%	(24.1% - 26.0%)	24.1%	(23.8% - 24.3%)
% with asthma			14.8%	(12.7% - 16.9%)	14.1%	(13.3% - 14.8%)	14.0%	(13.8% - 14.2%)
% with hypertension	w		81.9%	(79.6% - 84.2%)	80.4%	(79.5% - 81.3%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			6.8%	(5.3% - 8.2%)	5.0%	(4.6% - 5.5%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			45.3%	(42.2% - 48.5%)	46.4%	(45.2% - 47.5%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			24.1%	(21.5% - 26.8%)	26.5%	(25.5% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia			53.2%	(50.2% - 56.2%)	54.7%	(53.6% - 55.7%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			25.7%	(23.0% - 28.4%)	25.9%	(24.9% - 26.9%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions			65.4%	(62.5% - 68.2%)	66.3%	(65.3% - 67.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			8.4%	(6.8% - 10.1%)	8.1%	(7.5% - 8.7%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			579	(507 - 651)	627	(596 - 658)	628	(620 - 637)
Part D monthly prescription fills per person per year			53.2	(51.5 - 54.9)	52.6	(51.9 - 53.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year			2.1	(1.8 - 2.4)	2.1	(2.0 - 2.2)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			298	(245 - 350)	307	(289 - 325)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			130	(99 - 161)	110	(100 - 120)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			15.1%	(13.0% - 17.2%)	14.1%	(13.4% - 14.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis			51.8%	(48.6% - 54.9%)	53.2%	(52.0% - 54.4%)	52.0%	(51.7% - 52.3%)
% with osteoporosis		b	19.2%	(16.8% - 21.5%)	22.4%	(21.5% - 23.3%)	21.0%	(20.8% - 21.2%)
% with glaucoma			27.8%	(25.2% - 30.5%)	26.9%	(25.9% - 27.8%)	26.6%	(26.3% - 26.8%)
% with cataract			67.6%	(64.8% - 70.4%)	67.0%	(66.0% - 68.1%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			12.6%	(10.0% - 15.2%)	11.3%	(10.4% - 12.2%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			16.5%	(13.1% - 20.0%)	13.5%	(12.4% - 14.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			20.2%	(17.8% - 22.6%)	21.0%	(20.2% - 21.9%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			43.2%	(38.6% - 47.8%)	43.0%	(41.3% - 44.7%)	40.3%	(39.9% - 40.7%)
Physician visits per year			8.3	(7.8 - 8.7)	8.6	(8.5 - 8.8)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02910 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			3.9%	(3.2% - 4.7%)	4.5%	(4.2% - 4.8%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			3.4%	(2.4% - 4.5%)	4.2%	(3.8% - 4.7%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression			29.2%	(26.5% - 31.9%)	30.9%	(29.9% - 31.8%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			13.4%	(11.3% - 15.4%)	14.0%	(13.3% - 14.8%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.8%	(2.6% - 4.9%)	3.5%	(3.1% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.1%	(1.2% - 2.9%)	2.4%	(2.1% - 2.8%)	2.1%	(2.0% - 2.2%)
Home health visits per year			3.6	(2.8 - 4.4)	4.6	(4.2 - 5.0)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	b	b	4.3%	(3.5% - 5.1%)	5.5%	(5.2% - 5.8%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			18.2%	(11.9% - 24.5%)	13.6%	(11.6% - 15.7%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			7.5%	(4.2% - 10.7%)	5.5%	(4.3% - 6.6%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			22.5%	(16.9% - 28.2%)	20.2%	(17.8% - 22.6%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			5.1%	(2.6% - 7.5%)	6.0%	(4.7% - 7.3%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			5.2%	(1.9% - 8.5%)	6.7%	(5.1% - 8.3%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			12.6%	(8.8% - 16.5%)	12.2%	(10.3% - 14.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.