

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02914 (East Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02914 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol		b	75.8%	(73.7% - 77.8%)	79.3%	(78.2% - 80.4%)	78.0%	(77.7% - 78.2%)
% with diabetes			36.3%	(33.9% - 38.7%)	38.5%	(37.1% - 39.9%)	35.7%	(35.4% - 36.0%)
% with stroke			10.9%	(9.4% - 12.4%)	12.8%	(11.9% - 13.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			24.9%	(22.9% - 26.9%)	26.2%	(25.0% - 27.4%)	24.1%	(23.8% - 24.3%)
% with asthma			14.4%	(12.7% - 16.0%)	15.9%	(14.9% - 16.9%)	14.0%	(13.8% - 14.2%)
% with hypertension		b	79.3%	(77.3% - 81.2%)	82.3%	(81.3% - 83.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			4.6%	(3.6% - 5.6%)	6.2%	(5.6% - 6.8%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease	b		43.0%	(40.6% - 45.5%)	46.2%	(44.8% - 47.6%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			26.3%	(24.1% - 28.4%)	26.3%	(25.0% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia		b	49.8%	(47.4% - 52.2%)	55.4%	(54.1% - 56.8%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			24.9%	(22.8% - 27.0%)	27.3%	(26.1% - 28.6%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions		b	63.3%	(61.0% - 65.6%)	68.3%	(67.0% - 69.6%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions	b		10.3%	(8.8% - 11.7%)	8.1%	(7.4% - 8.9%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			574	(515 - 632)	616	(580 - 651)	628	(620 - 637)
Part D monthly prescription fills per person per year	b	b	52.4	(51.1 - 53.8)	56.3	(55.5 - 57.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year			2.2	(1.9 - 2.4)	2.2	(2.1 - 2.4)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			283	(246 - 319)	304	(282 - 326)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			97	(77 - 116)	128	(115 - 142)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			13.4%	(11.9% - 15.0%)	15.8%	(14.8% - 16.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	b		48.7%	(46.2% - 51.2%)	52.2%	(50.7% - 53.6%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			19.9%	(18.0% - 21.7%)	21.1%	(20.0% - 22.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			25.6%	(23.5% - 27.6%)	27.5%	(26.2% - 28.7%)	26.6%	(26.3% - 26.8%)
% with cataract	b	b	61.2%	(58.9% - 63.5%)	68.2%	(66.9% - 69.5%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			10.7%	(8.8% - 12.6%)	10.4%	(9.3% - 11.5%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			14.9%	(12.3% - 17.5%)	13.8%	(12.3% - 15.3%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			19.2%	(17.4% - 21.1%)	21.9%	(20.8% - 23.0%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	b	b	34.4%	(30.9% - 37.9%)	40.4%	(38.3% - 42.5%)	40.3%	(39.9% - 40.7%)
Physician visits per year			8.1	(7.7 - 8.5)	7.8	(7.6 - 8.0)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02914 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.9%	(4.2% - 5.6%)	4.9%	(4.5% - 5.2%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			3.2%	(2.4% - 4.0%)	3.8%	(3.3% - 4.3%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression		b	28.5%	(26.4% - 30.6%)	33.3%	(32.0% - 34.6%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			13.3%	(11.6% - 14.9%)	15.7%	(14.7% - 16.7%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.8%	(3.0% - 4.7%)	3.4%	(3.0% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.2%	(1.5% - 2.9%)	2.2%	(1.8% - 2.6%)	2.1%	(2.0% - 2.2%)
Home health visits per year			4.3	(3.5 - 5.2)	4.3	(3.8 - 4.7)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	b	b	4.8%	(4.1% - 5.4%)	7.1%	(6.7% - 7.6%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			12.4%	(8.5% - 16.3%)	12.3%	(9.9% - 14.6%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty	b		1.8%	(0.4% - 3.1%)	3.1%	(1.9% - 4.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty		w	24.3%	(19.5% - 29.0%)	16.6%	(14.2% - 18.9%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			8.2%	(4.8% - 11.7%)	5.6%	(4.0% - 7.1%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty	w	w	10.7%	(7.4% - 13.9%)	5.9%	(4.6% - 7.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			16.7%	(12.2% - 21.1%)	11.1%	(8.6% - 13.6%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.