

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02915 (East Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02915 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	w		81.4%	(79.7% - 83.1%)	79.3%	(78.2% - 80.4%)	78.0%	(77.7% - 78.2%)
% with diabetes	w	w	44.3%	(42.1% - 46.6%)	38.5%	(37.1% - 39.9%)	35.7%	(35.4% - 36.0%)
% with stroke			13.7%	(12.3% - 15.1%)	12.8%	(11.9% - 13.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease	w		27.1%	(25.2% - 29.0%)	26.2%	(25.0% - 27.4%)	24.1%	(23.8% - 24.3%)
% with asthma	w		16.9%	(15.3% - 18.5%)	15.9%	(14.9% - 16.9%)	14.0%	(13.8% - 14.2%)
% with hypertension	w		84.7%	(83.1% - 86.3%)	82.3%	(81.3% - 83.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			6.0%	(5.0% - 7.0%)	6.2%	(5.6% - 6.8%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			45.4%	(43.2% - 47.7%)	46.2%	(44.8% - 47.6%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			26.8%	(24.8% - 28.8%)	26.3%	(25.0% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia	w		57.4%	(55.2% - 59.5%)	55.4%	(54.1% - 56.8%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		28.9%	(26.9% - 30.9%)	27.3%	(26.1% - 28.6%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		70.5%	(68.5% - 72.5%)	68.3%	(67.0% - 69.6%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			7.1%	(6.0% - 8.3%)	8.1%	(7.4% - 8.9%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			656	(598 - 714)	616	(580 - 651)	628	(620 - 637)
Part D monthly prescription fills per person per year	w	w	58.4	(57.2 - 59.7)	56.3	(55.5 - 57.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.3	(2.1 - 2.6)	2.2	(2.1 - 2.4)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			290	(257 - 322)	304	(282 - 326)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year	w		134	(112 - 155)	128	(115 - 142)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			16.1%	(14.6% - 17.7%)	15.8%	(14.8% - 16.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis			52.3%	(50.0% - 54.6%)	52.2%	(50.7% - 53.6%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			21.3%	(19.6% - 23.1%)	21.1%	(20.0% - 22.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			28.6%	(26.6% - 30.5%)	27.5%	(26.2% - 28.7%)	26.6%	(26.3% - 26.8%)
% with cataract			70.0%	(68.0% - 72.0%)	68.2%	(66.9% - 69.5%)	67.9%	(67.6% - 68.2%)
% women with breast cancer	b		8.4%	(6.9% - 9.9%)	10.4%	(9.3% - 11.5%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			13.3%	(10.9% - 15.7%)	13.8%	(12.3% - 15.3%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism			21.3%	(19.5% - 23.0%)	21.9%	(20.8% - 23.0%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			39.2%	(35.8% - 42.6%)	40.4%	(38.3% - 42.5%)	40.3%	(39.9% - 40.7%)
Physician visits per year	b		7.4	(7.1 - 7.7)	7.8	(7.6 - 8.0)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02915 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			5.0%	(4.4% - 5.5%)	4.9%	(4.5% - 5.2%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			3.8%	(3.0% - 4.5%)	3.8%	(3.3% - 4.3%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		35.9%	(33.8% - 38.0%)	33.3%	(32.0% - 34.6%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias	w		17.9%	(16.3% - 19.6%)	15.7%	(14.7% - 16.7%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.2%	(2.5% - 4.0%)	3.4%	(3.0% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.2%	(1.6% - 2.8%)	2.2%	(1.8% - 2.6%)	2.1%	(2.0% - 2.2%)
Home health visits per year			4.3	(3.6 - 4.9)	4.3	(3.8 - 4.7)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	w	w	9.0%	(8.3% - 9.8%)	7.1%	(6.7% - 7.6%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty	w	w	21.3%	(15.8% - 26.8%)	12.3%	(9.9% - 14.6%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			6.0%	(2.4% - 9.6%)	3.1%	(1.9% - 4.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			22.1%	(17.0% - 27.2%)	16.6%	(14.2% - 18.9%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			5.3%	(2.5% - 8.0%)	5.6%	(4.0% - 7.1%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			4.6%	(2.5% - 6.6%)	5.9%	(4.6% - 7.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			13.7%	(8.0% - 19.4%)	11.1%	(8.6% - 13.6%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.