

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02916 (East Providence)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02916 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol			77.3%	(75.4% - 79.1%)	79.3%	(78.2% - 80.4%)	78.0%	(77.7% - 78.2%)
% with diabetes		b	34.4%	(32.2% - 36.6%)	38.5%	(37.1% - 39.9%)	35.7%	(35.4% - 36.0%)
% with stroke			13.3%	(11.8% - 14.8%)	12.8%	(11.9% - 13.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease	w		26.5%	(24.5% - 28.4%)	26.2%	(25.0% - 27.4%)	24.1%	(23.8% - 24.3%)
% with asthma			15.5%	(13.9% - 17.1%)	15.9%	(14.9% - 16.9%)	14.0%	(13.8% - 14.2%)
% with hypertension			80.0%	(78.2% - 81.8%)	82.3%	(81.3% - 83.4%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.9%	(4.9% - 7.0%)	6.2%	(5.6% - 6.8%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			47.0%	(44.6% - 49.3%)	46.2%	(44.8% - 47.6%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			26.6%	(24.6% - 28.6%)	26.3%	(25.0% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia			54.3%	(52.1% - 56.5%)	55.4%	(54.1% - 56.8%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		26.3%	(24.3% - 28.3%)	27.3%	(26.1% - 28.6%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w		67.5%	(65.4% - 69.5%)	68.3%	(67.0% - 69.6%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			9.0%	(7.7% - 10.3%)	8.1%	(7.4% - 8.9%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			586	(530 - 643)	616	(580 - 651)	628	(620 - 637)
Part D monthly prescription fills per person per year	w		55.9	(54.6 - 57.2)	56.3	(55.5 - 57.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year			2.2	(1.9 - 2.4)	2.2	(2.1 - 2.4)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			321	(283 - 358)	304	(282 - 326)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			122	(101 - 142)	128	(115 - 142)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			15.6%	(14.0% - 17.2%)	15.8%	(14.8% - 16.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis			52.4%	(50.1% - 54.8%)	52.2%	(50.7% - 53.6%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			22.2%	(20.4% - 24.0%)	21.1%	(20.0% - 22.2%)	21.0%	(20.8% - 21.2%)
% with glaucoma			26.1%	(24.1% - 28.0%)	27.5%	(26.2% - 28.7%)	26.6%	(26.3% - 26.8%)
% with cataract			66.6%	(64.5% - 68.7%)	68.2%	(66.9% - 69.5%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			11.5%	(9.7% - 13.3%)	10.4%	(9.3% - 11.5%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			12.9%	(10.6% - 15.2%)	13.8%	(12.3% - 15.3%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism	w		23.6%	(21.7% - 25.4%)	21.9%	(20.8% - 23.0%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			39.0%	(35.7% - 42.4%)	40.4%	(38.3% - 42.5%)	40.3%	(39.9% - 40.7%)
Physician visits per year			7.9	(7.6 - 8.2)	7.8	(7.6 - 8.0)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02916 ESTIMATE	MARGIN OF ERROR ²	EAST PROVIDENCE ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.7%	(4.1% - 5.3%)	4.9%	(4.5% - 5.2%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			4.0%	(3.2% - 4.8%)	3.8%	(3.3% - 4.3%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		32.9%	(30.8% - 35.0%)	33.3%	(32.0% - 34.6%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			14.3%	(12.8% - 15.9%)	15.7%	(14.7% - 16.7%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.5%	(2.7% - 4.3%)	3.4%	(3.0% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.4%	(1.7% - 3.0%)	2.2%	(1.8% - 2.6%)	2.1%	(2.0% - 2.2%)
Home health visits per year			4.6	(3.7 - 5.5)	4.3	(3.8 - 4.7)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports			6.4%	(5.7% - 7.1%)	7.1%	(6.7% - 7.6%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			15.0%	(9.7% - 20.4%)	12.3%	(9.9% - 14.6%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			5.8%	(2.6% - 9.1%)	3.1%	(1.9% - 4.3%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			16.5%	(10.6% - 22.4%)	16.6%	(14.2% - 18.9%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			9.7%	(4.3% - 15.1%)	5.6%	(4.0% - 7.1%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			8.0%	(3.2% - 12.9%)	5.9%	(4.6% - 7.2%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			12.8%	(7.4% - 18.2%)	11.1%	(8.6% - 13.6%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.