

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02920 (Cranston)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02920 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	w		80.5%	(79.3% - 81.6%)	78.9%	(78.0% - 79.8%)	78.0%	(77.7% - 78.2%)
% with diabetes	w		38.6%	(37.1% - 40.1%)	36.9%	(35.8% - 38.0%)	35.7%	(35.4% - 36.0%)
% with stroke			12.5%	(11.6% - 13.5%)	12.1%	(11.4% - 12.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease	w		26.5%	(25.2% - 27.8%)	25.0%	(24.1% - 26.0%)	24.1%	(23.8% - 24.3%)
% with asthma			14.3%	(13.3% - 15.3%)	14.1%	(13.3% - 14.8%)	14.0%	(13.8% - 14.2%)
% with hypertension	w		82.2%	(81.0% - 83.3%)	80.4%	(79.5% - 81.3%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			5.4%	(4.8% - 6.0%)	5.0%	(4.6% - 5.5%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease	w		48.9%	(47.3% - 50.4%)	46.4%	(45.2% - 47.5%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure	w		27.7%	(26.4% - 29.1%)	26.5%	(25.5% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia	w	w	58.5%	(57.1% - 60.0%)	54.7%	(53.6% - 55.7%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease	w		27.3%	(26.0% - 28.7%)	25.9%	(24.9% - 26.9%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions	w	w	70.0%	(68.7% - 71.3%)	66.3%	(65.3% - 67.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions	w	w	6.8%	(6.0% - 7.5%)	8.1%	(7.5% - 8.7%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year			669	(628 - 711)	627	(596 - 658)	628	(620 - 637)
Part D monthly prescription fills per person per year	b		53.1	(52.2 - 53.9)	52.6	(51.9 - 53.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year	w		2.2	(2.0 - 2.3)	2.1	(2.0 - 2.2)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year	w		323	(298 - 348)	307	(289 - 325)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			117	(102 - 131)	110	(100 - 120)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation			14.9%	(13.9% - 15.9%)	14.1%	(13.4% - 14.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	w		55.1%	(53.6% - 56.7%)	53.2%	(52.0% - 54.4%)	52.0%	(51.7% - 52.3%)
% with osteoporosis	w		24.4%	(23.2% - 25.6%)	22.4%	(21.5% - 23.3%)	21.0%	(20.8% - 21.2%)
% with glaucoma			27.9%	(26.6% - 29.2%)	26.9%	(25.9% - 27.8%)	26.6%	(26.3% - 26.8%)
% with cataract	w	w	70.9%	(69.6% - 72.2%)	67.0%	(66.0% - 68.1%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			11.3%	(10.1% - 12.5%)	11.3%	(10.4% - 12.2%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			12.7%	(11.2% - 14.2%)	13.5%	(12.4% - 14.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism	w	w	23.3%	(22.1% - 24.5%)	21.0%	(20.2% - 21.9%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia	w	w	47.0%	(44.8% - 49.3%)	43.0%	(41.3% - 44.7%)	40.3%	(39.9% - 40.7%)
Physician visits per year	w		9.0	(8.8 - 9.3)	8.6	(8.5 - 8.8)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02920 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.6%	(4.2% - 5.0%)	4.5%	(4.2% - 4.8%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture	w		4.6%	(4.0% - 5.2%)	4.2%	(3.8% - 4.7%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression	w		32.6%	(31.3% - 34.0%)	30.9%	(29.9% - 31.8%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias			15.2%	(14.2% - 16.3%)	14.0%	(13.3% - 14.8%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.4%	(2.9% - 4.0%)	3.5%	(3.1% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer			2.5%	(2.0% - 2.9%)	2.4%	(2.1% - 2.8%)	2.1%	(2.0% - 2.2%)
Home health visits per year	w		4.9	(4.4 - 5.4)	4.6	(4.2 - 5.0)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports			6.1%	(5.7% - 6.6%)	5.5%	(5.2% - 5.8%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty			13.7%	(11.1% - 16.3%)	13.6%	(11.6% - 15.7%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			4.5%	(3.2% - 5.7%)	5.5%	(4.3% - 6.6%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			21.1%	(18.1% - 24.1%)	20.2%	(17.8% - 22.6%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			6.0%	(4.3% - 7.8%)	6.0%	(4.7% - 7.3%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			6.8%	(4.8% - 8.8%)	6.7%	(5.1% - 8.3%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			12.1%	(9.8% - 14.4%)	12.2%	(10.3% - 14.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.