

RHODE ISLAND HEALTHY AGING COMMUNITY DATA PROFILE

02921 (Cranston)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02921 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
SERIOUS CHRONIC DISEASE								
% high cholesterol	w		82.0%	(79.6% - 84.3%)	78.9%	(78.0% - 79.8%)	78.0%	(77.7% - 78.2%)
% with diabetes			33.5%	(30.4% - 36.7%)	36.9%	(35.8% - 38.0%)	35.7%	(35.4% - 36.0%)
% with stroke			11.9%	(10.0% - 13.8%)	12.1%	(11.4% - 12.7%)	12.5%	(12.3% - 12.6%)
% with chronic obstructive pulmonary disease			23.1%	(20.6% - 25.6%)	25.0%	(24.1% - 26.0%)	24.1%	(23.8% - 24.3%)
% with asthma			14.0%	(11.9% - 16.0%)	14.1%	(13.3% - 14.8%)	14.0%	(13.8% - 14.2%)
% with hypertension			79.4%	(76.9% - 81.8%)	80.4%	(79.5% - 81.3%)	79.0%	(78.8% - 79.3%)
% ever had a heart attack			4.4%	(3.3% - 5.6%)	5.0%	(4.6% - 5.5%)	5.4%	(5.3% - 5.6%)
% with ischemic heart disease			44.1%	(40.8% - 47.4%)	46.4%	(45.2% - 47.5%)	45.9%	(45.5% - 46.2%)
% with congestive heart failure			24.7%	(21.9% - 27.5%)	26.5%	(25.5% - 27.5%)	24.8%	(24.5% - 25.1%)
% with anemia			51.5%	(48.6% - 54.5%)	54.7%	(53.6% - 55.7%)	52.2%	(51.9% - 52.5%)
% with chronic kidney disease			23.4%	(20.7% - 26.2%)	25.9%	(24.9% - 26.9%)	23.3%	(23.1% - 23.6%)
% with 4+ chronic conditions			62.8%	(59.9% - 65.7%)	66.3%	(65.3% - 67.4%)	63.9%	(63.6% - 64.2%)
% with 0 chronic conditions			8.3%	(6.6% - 10.1%)	8.1%	(7.5% - 8.7%)	8.4%	(8.2% - 8.6%)
Emergency room visits/1000 persons 65+ years per year	b	b	522	(450 - 593)	627	(596 - 658)	628	(620 - 637)
Part D monthly prescription fills per person per year			52.2	(50.5 - 54.0)	52.6	(51.9 - 53.2)	54.2	(54.0 - 54.4)
Durable medical equipment claims per year			1.9	(1.6 - 2.2)	2.1	(2.0 - 2.2)	2.0	(1.9 - 2.0)
Inpatient hospital stays/1000 persons 65+ years per year			275	(231 - 320)	307	(289 - 325)	284	(280 - 289)
Skilled nursing facility stays/1000 persons 65+ years per year			111	(83 - 139)	110	(100 - 120)	100	(98 - 103)
MILD CHRONIC DISEASE								
% with atrial fibrillation	b		11.8%	(10.0% - 13.7%)	14.1%	(13.4% - 14.8%)	15.2%	(15.0% - 15.4%)
% with osteoarthritis/rheumatoid arthritis	w		56.1%	(52.7% - 59.5%)	53.2%	(52.0% - 54.4%)	52.0%	(51.7% - 52.3%)
% with osteoporosis			19.7%	(17.4% - 22.0%)	22.4%	(21.5% - 23.3%)	21.0%	(20.8% - 21.2%)
% with glaucoma			27.1%	(24.5% - 29.7%)	26.9%	(25.9% - 27.8%)	26.6%	(26.3% - 26.8%)
% with cataract	b		63.3%	(60.4% - 66.3%)	67.0%	(66.0% - 68.1%)	67.9%	(67.6% - 68.2%)
% women with breast cancer			11.7%	(9.2% - 14.1%)	11.3%	(10.4% - 12.2%)	10.7%	(10.4% - 10.9%)
% men with prostate cancer			15.3%	(12.0% - 18.6%)	13.5%	(12.4% - 14.7%)	13.8%	(13.5% - 14.1%)
% with hypothyroidism	b	b	16.7%	(14.5% - 18.9%)	21.0%	(20.2% - 21.9%)	21.1%	(20.9% - 21.4%)
% with benign prostatic hyperplasia			43.2%	(38.6% - 47.8%)	43.0%	(41.3% - 44.7%)	40.3%	(39.9% - 40.7%)
Physician visits per year	w		8.5	(8.1 - 9.0)	8.6	(8.5 - 8.8)	8.0	(7.9 - 8.0)

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	BETTER / WORSE CITY RATE ¹	02921 ESTIMATE	MARGIN OF ERROR ²	CRANSTON ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
DISABILITY and LATER LIFE CHRONIC DISEASE								
Age-sex adjusted 1-year mortality rate			4.7%	(3.8% - 5.6%)	4.5%	(4.2% - 4.8%)	4.8%	(4.7% - 4.8%)
% ever had a hip fracture			4.0%	(2.9% - 5.1%)	4.2%	(3.8% - 4.7%)	3.9%	(3.8% - 4.0%)
% ever diagnosed with depression			28.7%	(26.0% - 31.4%)	30.9%	(29.9% - 31.8%)	30.0%	(29.8% - 30.3%)
% with Alzheimer's disease or related dementias	b		11.8%	(9.9% - 13.8%)	14.0%	(13.3% - 14.8%)	14.4%	(14.2% - 14.6%)
% with colon cancer			3.8%	(2.7% - 5.0%)	3.5%	(3.1% - 3.9%)	3.2%	(3.1% - 3.3%)
% with lung cancer	w		3.5%	(2.4% - 4.6%)	2.4%	(2.1% - 2.8%)	2.1%	(2.0% - 2.2%)
Home health visits per year			3.8	(3.0 - 4.6)	4.6	(4.2 - 5.0)	3.7	(3.7 - 3.8)
% of 65+ getting Medicaid long term services and supports	b		5.0%	(4.2% - 5.9%)	5.5%	(5.2% - 5.8%)	6.2%	(6.1% - 6.3%)
% 65+ with hearing difficulty	b	b	6.4%	(2.7% - 10.0%)	13.6%	(11.6% - 15.7%)	13.8%	(13.2% - 14.3%)
% 65+ with vision difficulty			1.8%	(-1.4% - 4.9%)	5.5%	(4.3% - 6.6%)	5.2%	(4.9% - 5.5%)
% 65+ with ambulatory difficulty			17.6%	(10.7% - 24.5%)	20.2%	(17.8% - 22.6%)	19.9%	(19.3% - 20.5%)
% 65+ with cognition difficulty			4.6%	(0.8% - 8.3%)	6.0%	(4.7% - 7.3%)	7.8%	(7.3% - 8.2%)
% 65+ with self-care difficulty			6.5%	(2.0% - 10.9%)	6.7%	(5.1% - 8.3%)	6.6%	(6.2% - 6.9%)
% 65+ with independent living difficulty			12.8%	(7.3% - 18.3%)	12.2%	(10.3% - 14.1%)	13.7%	(13.2% - 14.2%)

Notes.¹ Zip code-city-state differences that the margins of error do not overlap each other are noted b or w.

² All zip code, city, and state margins of error are 90% confidence intervals.