

Manchester is a city in Hillsborough County located on the Merrimack River. There are 14,552 residents age 65 or older. Compared to state average rates, older residents had a lower rate of migraine. However, they had higher rates of hip fracture, tooth loss, obesity, high cholesterol, depression, anxiety/bipolar disorders, PTSD, schizophrenia, personality disorders, substance/tobacco use disorders, Alzheimer's disease, diabetes, stroke, COPD, asthma, hypertension, ischemic heart disease, CHF, atrial fibrillation, peripheral vascular disease, arthritis, osteoporosis, colon and prostate cancers, hypothyroidism, anemia, chronic kidney disease, liver diseases, fibromyalgia, glaucoma, cataracts, ulcers, and hearing, visual, and mobility impairments. They are less likely to take the health promotion step of eating 5 or more servings of fruit/vegetables per day. Community resources to support healthy aging include 333 primary care providers, 2 hospitals, and 13 nursing homes within 5 miles, 15 home health agencies, 2 community health centers, an adult day center, 2 memory cafes, a senior center, 5 assisted living sites, 8 universities/community colleges, 2 public libraries, a YMCA, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		110,231	1,327,503
Population 60 years or older as % of total population		18.9%	22.7%
Total population 60 years or older		20,804	301,630
Population 65 years or older as % of total population		13.2%	15.8%
Total population 65 years or older		14,552	210,385
% 65-74 years		54.3%	58.5%
% 75-84 years		28.5%	28.6%
% 85 years or older		17.3%	12.9%
Gender (65+ population)			
% female		58.8%	54.7%
Race/Ethnicity (65+ population)			
% White		95.9%	97.7%
% African American		1.2%	0.5%
% Asian		1.8%	0.9%
% Other		1.1%	0.9%
% Hispanic/Latino		3.8%	0.9%
Marital Status (65+ population)			
% married		45.3%	58.5%
% divorced/separated		17.5%	14.0%
% widowed		28.0%	22.9%
% never married		9.2%	4.6%
Education (65+ population)			
% with less than high school education		20.1%	12.3%
% with high school or some college		56.3%	57.1%
% with college degree		23.6%	30.6%
% of 65+ population living alone		34.2%	26.1%
% of 65+ population who speak only English at home		74.9%	91.3%
% of 65+ population who are veterans of military service		20.5%	24.8%
Age-sex adjusted 1-year mortality rate	W	4.5%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		7.1%	3.6%
% moved from different county in New Hampshire		0.8%	1.0%
% moved from different state		1.1%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		71.0%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		22.0%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		52.7%	58.6%
% 60+ met CDC guidelines for both types of physical activities		15.5%	19.9%
% 60+ getting recommended hours of sleep		64.0%	66.4%
% 60+ injured in a fall within last 12 months		12.3%	10.4%
% 65+ had hip fracture	W	3.8%	3.3%
% 60+ with self-reported fair or poor health status		21.6%	16.5%
% 60+ with 15+ physically unhealthy days last month		15.0%	12.5%
% 60+ with physical exam/check-up in past year		90.7%	86.5%
% 60+ met CDC preventive health screening goals		34.9%	40.3%
% 60+ flu shot past year		61.0%	59.3%
% 65+ with pneumonia vaccine		77.1%	77.8%
% 60+ with cholesterol screening		95.5%	95.3%
% 60+ women with a mammogram within last 2 years		84.1%	79.3%
% 60+ with colorectal cancer screening		72.5%	77.0%
% 60+ with HIV test		13.5%	13.5%
% 60+ current smokers		9.6%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth	W	36.8%	29.0%
% 60+ with annual dental exam		72.8%	75.7%
# of dentists per 100,000 persons (all ages) (county)		77	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day	W	14.4%	20.2%
% 60+ self-reported obese		30.7%	27.2%
% 65+ clinically diagnosed obese	W	19.6%	16.7%
% 65+ with high cholesterol	W	78.0%	72.2%
% 60+ excessive drinking		8.3%	9.2%
% 65+ with poor supermarket access		35.5%	28.4%

% 65+ with depression W 33.9% 28.1 % 65+ with anxiety disorders W 27.8% 21.1 % 65+ with pipolar disorders W 4.9% 3.3 % 65+ with post-traumatic stress disorder W 2.2% 1.4 % 65+ with schizophrenia & other psychotic disorders W 2.2% 1.4 % 65+ with personality disorders W 2.0% 1.5 % 65+ with personality disorders W 2.0% 1.5 % 65+ with bersonality disorders W 2.0% 1.5 % 65+ with bersonality disorders W 13.4% 10.2 % 65+ with bersonality disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with Alzheimer's disease or related dementias W 13.4% 10.2 % 65+ with Alzheimer's disease or related dementias W 14.2% 28.2 % 65+ with Alzheimer's disease or related dementias W 13.4% 10.3 % 65+ with Alzheimer's disease or related dement	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression W 33.9% 28.1 % 65+ with anxiety disorders W 27.8% 21.1 % 65+ with pipolar disorders W 4.9% 3. % 65+ with post-traumatic stress disorder W 2.2% 1.4 % 65+ with schizophrenia & other psychotic disorders W 2.0% 1. % 65+ with schizophrenia & other psychotic disorders W 2.0% 1. % 65+ with personality disorders W 2.0% 1. # opioid deaths (all ages) (county) 491 1,2 % 65+ with busbtance use disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with busbtance use disorders W 13.4% 10. CHRONIC DISEASE W 14.2% 12.2 % 65+ with Alzheimer's disease or related dementias W 14.2% 28.2 % 65+ with Alzheimer's disease or related dementias W 14.2% 28.2 % 65+ with Alzheimer's disease or related dementias W	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders W 27.8% 21.3 % 65+ with bipolar disorders W 4.9% 3.3 % 65+ with post-traumatic stress disorder W 2.2% 1.4 % 65+ with personality disorders W 7.2% 4.3 % 65+ with personality disorders W 2.0% 1. # opioid deaths (all ages) (county) 491 1.2 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.3 % 65+ with basco use disorders W 13.4% 10. CHRONIC DISEASE W 65+ with Alzheimer's disease or related dementias W 14.2% 12.1 % 65+ with Alzheimer's disease or related dementias W 34.3% 28. % 65+ with Alzheimer's disease or related dementias W 13.4% 10.3 % 65+ with Alzheimer's disease or related dementias W 13.4% 10.3 % 65+ with Alzheimer's disease or related dementias W 13.4% 10.3 % 65+ with Alzheimer's disease W 13.4% 10.3 <td>% 60+ with 15+ days poor mental health last month</td> <td></td> <td>7.4%</td> <td>6.9%</td>	% 60+ with 15+ days poor mental health last month		7.4%	6.9%
% 65+ with bipolar disorders W 4.9% 3.3 % 65+ with post-traumatic stress disorder W 2.2% 1.4 % 65+ with schizophrenia & other psychotic disorders W 7.2% 4.4 % 65+ with schizophrenia & other psychotic disorders W 2.0% 1.5 % 65+ with personality disorders W 2.0% 1.5 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with tobacco use disorders W 13.4% 10.0 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12.1 % 65+ with substance use disorders W 34.3% 28.3 % 65+ with dabetes W 34.3% 28.3 % 65+ with bustance use disorders W 34.3% 28.3 % 65+ with dabetes W 34.3% 28.3 % 65+ with obsecutes W 34.3% 28.3 % 65+ with chronic obstructive pulmonary disease W 17.4% 4.3 % 65+ with hypertension	% 65+ with depression	W	33.9%	28.8%
% 65+ with post-traumatic stress disorder W 2.2% 1.4 % 65+ with schizophrenia & other psychotic disorders W 7.2% 4.3 % 65+ with personality disorders W 2.0% 1. # opioid deaths (all ages) (county) 491 1,2 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.3 % 65+ with bobacco use disorders W 13.4% 10. CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12. % 65+ with diabetes W 34.3% 28. % 65+ with stroke W 13.4% 10. % 65+ with stroke W 13.4% 10. % 65+ with stroke W 17.4% 13. % 65+ with stroke W 17.4% 13. % 65+ with stroke W 25.6% 20. % 65+ with stroke W 74.6% 70. % 65+ with hypertension W 74.6% 70. % 65+ with hypertension	% 65+ with anxiety disorders	W	27.8%	21.9%
% 65+ with schizophrenia & other psychotic disorders W 7.2% 4.8 % 65+ with personality disorders W 2.0% 1. # opioid deaths (all ages) (county) 491 1.2 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.1 % 65+ with tobacco use disorders W 13.4% 10. CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12.4 % 65+ with diabetes W 34.3% 28.3 % 65+ with broke W 13.4% 10.3 % 65+ with chronic cobstructive pulmonary disease W 25.6% 20.2 % 65+ with hypertension W 74.6% 70.2 % 65+ with bischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17. % 65+ with peripheral vascular disease W 18.0% 14.4 % 65+ with osteoarthritis/rheumatoid arthritis W 21.8% 22.4 % 65+ with lung cancer W	% 65+ with bipolar disorders	W	4.9%	3.1%
% 65+ with personality disorders W 2.0% 1. # opioid deaths (all ages) (county) 491 1,2 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7,3% 5.3 % 65+ with tobacco use disorders W 13,4% 10. CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14,2% 12. % 65+ with stroke W 13,4% 10. % 65+ with obrionic obstructive pulmonary disease W 13,4% 10. % 65+ with asthma W 17,4% 13. % 65+ with asthma W 17,4% 13. % 65+ with hypertension W 74,6% 70. % 65+ with hypertension W 74,6% 70. % 65+ with obreath disease W 38,8% 34. % 65+ with congestive heart failure W 21,8% 17. % 65+ with obsteoporosis W 18,0% 14. % 65+ with osteoporosis W 18,0% 14. % 65+ with lung	% 65+ with post-traumatic stress disorder	W	2.2%	1.4%
# opioid deaths (all ages) (county) 491 1,2 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.9 % 65+ with tobacco use disorders W 13.4% 10.6 CHRONIC DISEASE ***65+ with Alzheimer's disease or related dementias W 14.2% 12.9 % 65+ with diabetes W 34.3% 28.3 % 65+ with chronic obstructive pulmonary disease W 15.6% 20.9 % 65+ with stroke W 17.4% 13.4 % 65+ with stroma W 17.4% 13.4 % 65+ with hypertension W 74.6% 70.3 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with ongestive heart failure W 21.8% 17.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoarthritis/rheumatoid arthritis W 22.1% 17.9 % 65+ with osteoarthritis/rheumatoid arthritis W 22.1% 17.9 % 65+ with lung cancer W 22.8% 2.0 % 65+ with osteoarthritis/rheumatoid arthritis W 23.9% 2.0 % 65+ with heniq cancer W 23.8% 2.0 % 65+ with hong prostatic hyperplasia 38.1% 36.1 % 65+ with hong prostatic hyperplasia 38.1% 36.1 % 65+ with hong prostatic hyperplasia W 26.3% 20.0 % 65+ with hypothyroidism W 26.3% 20.0 % 65+ with hypothyroidism W 26.3% 20.0 % 65+ with hypothyroidism W 26.3% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ with honic kidney disease W 26.5% 20.0 % 65+ wi	% 65+ with schizophrenia & other psychotic disorders	W	7.2%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.3% 5.5 % 65+ with tobacco use disorders W 13.4% 10.0 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12.1 % 65+ with diabetes W 34.3% 28.3 % 65+ with stroke W 13.4% 10.1 % 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with chronic obstructive pulmonary disease W 17.4% 13.1 % 65+ with periphersion W 74.6% 70.2 % 65+ with hypertension W 74.6% 70.2 % 65+ with hypertension heart disease W 38.8% 34.1 % 65+ with ischemic heart disease W 38.8% 34.1 % 65+ with congestive heart failure W 21.8% 17.7 % 65+ with peripheral vascular disease W 18.0% 14.2 % 65+ with peripheral vascular disease W 18.0% 14.2 % 65+ with osteoporosis W 22.1% 17.7 % 65+ with lung cancer	% 65+ with personality disorders	W	2.0%	1.1%
% 65+ with tobacco use disorders W 13.4% 10.4 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12.1 % 65+ with diabetes W 34.3% 28.3 % 65+ with stroke W 13.4% 10.3 % 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with hypertension W 17.4% 13.1 % 65+ with hypertension W 74.6% 70.3 % 65+ with hypertension W 74.6% 70.3 % 65+ ever had a heart attack 4.7% 4.3 % 65+ ever had a heart attack 4.7% 4.3 % 65+ with conjective heart failure W 38.8% 34.3 % 65+ with peripheral vascular disease W 18.0% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with leukemias and lymphomas 2.1% 2.1% 2.1% % 65+ with	# opioid deaths (all ages) (county)		491	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 14.2% 12.4 % 65+ with diabetes W 34.3% 28.3 % 65+ with stroke W 13.4% 10.4 % 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with chronic obstructive pulmonary disease W 17.4% 13.4 % 65+ with asthma W 17.4% 13.4 % 65+ with hypertension W 74.6% 70.3 % 65+ with hypertension W 74.6% 70.3 % 65+ with hypertension W 38.8% 34.4 % 65+ with ischemic heart disease W 38.8% 34.4 % 65+ with congestive heart failure W 21.8% 17. % 65+ with thrall fibrillation W 15.8% 14.4 % 65+ with osteoarthritis/rheumatoid arthritis W 18.0% 14.4 % 65+ with osteoarthritis/rheumatoid arthritis W 22.1% 17.4 % 65+ with leukemias and lymphomas 21.7% 17.4	% 65+ with substance use disorders (drug use +/or alcohol abuse)	W	7.3%	5.5%
% 65+ with Alzheimer's disease or related dementias W 14.2% 12.0 % 65+ with diabetes W 34.3% 28.3 % 65+ with stroke W 13.4% 10.0 % 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with asthma W 17.4% 13.1 % 65+ with hypertension W 74.6% 70.2 % 65+ ever had a heart attack 4.7% 4.4 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.3 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoarthritis/rheumatoid arthritis W 22.1% 17.4 % 65+ with lung cancer U 1.7% 11.4 % 65+ with lung cancer U 1.7% 11.4 % 65+ with lung cancer W 2.8% 2.2 % 65+ with lung cancer W <	% 65+ with tobacco use disorders	W	13.4%	10.4%
% 65+ with diabetes W 34,3% 28.3 % 65+ with stroke W 13,4% 10.0 % 65+ with chronic obstructive pulmonary disease W 25,6% 20.3 % 65+ with asthma W 17,4% 13.0 % 65+ with hypertension W 74,6% 70.3 % 65+ ever had a heart attack 4,7% 4.4 % 65+ with ischemic heart disease W 38,8% 34.3 % 65+ with congestive heart failure W 21,8% 17.3 % 65+ with atrial fibrillation W 15,8% 14.3 % 65+ with peripheral vascular disease W 18,0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53,9% 49.3 % 65+ with leukemias and lymphomas 2,1% 20.3 % 65+ with lung cancer 1,7% 11.4 % 65+ with colon cancer W 2,8% 2.4 % 65+ women with preast cancer W 2,8% 2.4 % 65+ men with prostate cancer W 12,7% 11.3 % 65+ with benign prostatic hyperplasia 38,1% 36,3 <	CHRONIC DISEASE			
% 65+ with stroke W 13.4% 10.4 % 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with asthma W 17.4% 13.4 % 65+ with hypertension W 74.6% 70.3 % 65+ ever had a heart attack 4.7% 4.4 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.4 % 65+ with osteoporosis W 53.9% 49.4 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with lung cancer 1.7% 1.1 % 65+ with lung cancer 1.7% 1.1 % 65+ women with breast cancer W 2.8% 2.4 % 65+ women with prostate cancer W 12.7% 11. % 65+ women with prostate cancer W 12.7% 11. % 65+ with benign prostatic hyperplasia 38.1% 36.3 % 65+ with hypothyr	% 65+ with Alzheimer's disease or related dementias	W	14.2%	12.0%
% 65+ with chronic obstructive pulmonary disease W 25.6% 20.9 % 65+ with asthma W 17.4% 13.1 % 65+ with hypertension W 74.6% 70.2 % 65+ ever had a heart attack 4.7% 4.4 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.3 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with lung cancer 1.7% 1.4 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ wornen with prostate cancer W 12.7% 11.5 % 65+ men with prostate cancer W 12.7% 11.4 % 65+ with henign prostatic hyperplasia 38.1% 36.3 % 65+ with hypothyroidism W 26.3% 20.3	% 65+ with diabetes	W	34.3%	28.2%
% 65+ with asthma W 17.4% 13.0 % 65+ with hypertension W 74.6% 70.0 % 65+ ever had a heart attack 4.7% 4.8 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with congestive heart failure W 15.8% 14.3 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.1 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.1 % 65+ women with prostate cancer W 12.7% 11.1 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with hypothyroidism W 26.3% 20.0 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney di	% 65+ with stroke	W	13.4%	10.8%
% 65+ with hypertension W 74.6% 70.2 % 65+ ever had a heart attack 4.7% 4.4 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with ischemic heart disease W 21.8% 17.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.1 % 65+ with osteoparthritis/rheumatoid arthritis W 53.9% 49.2 % 65+ with osteoparosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with lung cancer W 2.8% 2.4 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with prostate cancer W 1.7% 11.5 % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.3 % 65+ with hypothyroidism W 26.3% 20.4 % 65+ with hypothyroidism W 26.3% 20.4	% 65+ with chronic obstructive pulmonary disease	W	25.6%	20.5%
% 65+ ever had a heart attack 4.7% 4.8 % 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.1 % 65+ with osteoparthritis/rheumatoid arthritis W 53.9% 49.1 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.1 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.2 % 65+ women with breast cancer 9.1% 9.1 % 65+ women with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.3 % 65+ with hypothyroidism W 26.3% 20.4 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with asthma	W	17.4%	13.0%
% 65+ with ischemic heart disease W 38.8% 34.3 % 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with colon cancer W 2.8% 2.4 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with preast cancer 9.1% 9.1 % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.5	% 65+ with hypertension	W	74.6%	70.2%
% 65+ with congestive heart failure W 21.8% 17.3 % 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.4 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer W 2.8% 2.4 % 65+ women with prostate cancer W 1.7% 1.1 % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.5	% 65+ ever had a heart attack		4.7%	4.5%
% 65+ with atrial fibrillation W 15.8% 14.4 % 65+ with peripheral vascular disease W 18.0% 14.4 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.5 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.4 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1. % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with hypothyroidism W 26.3% 20.0 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.5	% 65+ with ischemic heart disease	W	38.8%	34.3%
% 65+ with peripheral vascular disease W 18.0% 14.3 % 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1. % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with congestive heart failure	W	21.8%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis W 53.9% 49.3 % 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.1 % 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with hypothyroidism W 26.3% 20.4 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with atrial fibrillation	W	15.8%	14.4%
% 65+ with osteoporosis W 22.1% 17.4 % 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.7 % 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.4 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with peripheral vascular disease	W	18.0%	14.7%
% 65+ with leukemias and lymphomas 2.1% 2.0 % 65+ with lung cancer 1.7% 1.0 % 65+ with colon cancer W 2.8% 2.0 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.1 % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.05 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with osteoarthritis/rheumatoid arthritis	W	53.9%	49.1%
% 65+ with lung cancer 1.7% 1.6 % 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.7 % 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with osteoporosis	W	22.1%	17.4%
% 65+ with colon cancer W 2.8% 2.4 % 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.7 % 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with leukemias and lymphomas		2.1%	2.0%
% 65+ women with breast cancer 9.1% 9.8 % 65+ women with endometrial cancer 1.9% 1.7 % 65+ men with prostate cancer W 12.7% 11.5 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.05 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with lung cancer		1.7%	1.6%
% 65+ women with endometrial cancer 1.9% 1.7 % 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with colon cancer	W	2.8%	2.4%
% 65+ men with prostate cancer W 12.7% 11.9 % 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ women with breast cancer		9.1%	9.8%
% 65+ with benign prostatic hyperplasia 38.1% 36.8 % 65+ with HIV/AIDS 0.07% 0.09 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ women with endometrial cancer		1.9%	1.7%
% 65+ with HIV/AIDS 0.07% 0.08 % 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ men with prostate cancer	W	12.7%	11.5%
% 65+ with hypothyroidism W 26.3% 20.8 % 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with benign prostatic hyperplasia		38.1%	36.8%
% 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3	% 65+ with HIV/AIDS		0.07%	0.05%
% 65+ with anemia W 43.7% 37.3 % 65+ with chronic kidney disease W 28.0% 22.3		W		20.8%
% 65+ with chronic kidney disease W 28.0% 22.3		W		37.3%
·		W		22.3%
% 8.8% $%$ 6.9 with liver diseases $%$ 8.8%	% 65+ with liver diseases	W	8.8%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	W		18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache	В	3.5%	4.0%
% 65+ with epilepsy		2.4%	2.1%
% 65+ with traumatic brain injury		1.1%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	W	27.1%	22.9%
% 65+ with cataract	W	63.0%	61.2%
% 65+ with pressure ulcer or chronic ulcer	W	8.6%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	W	62.6%	54.4%
% 65+ with 0 chronic conditions	W	8.2%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		16.9%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment	W	15.6%	14.4%
% 65+ with self-reported vision difficulty		7.0%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment	W	1.3%	0.9%
% 65+ with self-reported cognition difficulty	W	10.1%	6.9%
% 65+ with self-reported ambulatory difficulty	W	24.3%	18.8%
% 65+ with clinical diagnosis of mobility impairments	W	3.8%	3.2%
% 65+ with self-reported self-care difficulty	W	8.2%	5.6%
% 65+ with self-reported independent living difficulty	W	15.0%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		8.4%	7.9%
% dually eligible for Medicare and Medicaid	*	12.7%	7.5%
% 60+ with a regular doctor		95.0%	95.8%
% 60+ who did not see doctor when needed due to cost		7.1%	5.4%
# of primary care providers within 5 miles		333	2,961
# of hospitals within 5 miles		2	26
# of nursing homes within 5 miles		13	74
# of home health agencies		15	49
# of community health centers		2	22
# of adult day health centers		1	21
# of memory cafes		2	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	7.3	6.3
Emergency room visits/1000 persons 65+ years per year	*	613	572
Part D monthly prescription fills per person per year	*	54.3	49.1
Home health visits per year	*	3.3	2.5
Durable medical equipment claims per year	*	2.4	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	307	237
Medicare inpatient hospital readmissions (as % of admissions)		16.6%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year	*	113	76
# skilled nursing home Medicare beds/1000 persons 65+ years		48	33
% 65+ getting Medicaid long term services and supports	*	5.9%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		1	44
Air pollution: annual # of unhealthy days for 65+ (county)		1	N/A
% of grandparents raising grandchildren		0.9%	0.8%
% of grandparents who live with grandchildren		3.0%	2.5%
# of assisted living sites		5	134
% of vacant homes in community		7.5%	16.0%
# of universities and community colleges		8	41
# of public libraries		2	234
# of YMCAs		1	12
% in county with access to broadband (all ages)		94.0%	93.0%
% 60+ who used Internet in last month		63.7%	77.6%
Voter participation rate in 2018 election (age 18+)		42.6%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		655	207
Homicide rate /100,000 persons (county)		2	1
# firearm fatalities (county)		196	586
Property crime rate /100,000 persons		3,701	2,012
% 65+ who own a motor vehicle		81.1%	91.0%
% 60+ who always drive wearing a seatbelt		77.1%	77.1%
# of fatal crashes involving adult age 60+/town		9	151
# of fatal crashes involving adult age 60+/county		31	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		8.0%	5.4%
% 60+ receiving food stamps past year		10.9%	5.7%
% 65+ employed past year		22.7%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		27.1%	18.2%
% households with annual income \$20,000-\$49,999		37.1%	36.5%
% households with annual income > \$50,000		35.9%	45.3%
% 60+ own home		61.6%	79.9%
% 60+ have mortgage on home		30.4%	35.3%
% 65+ households spend >35% of income on housing (renter)		19.2%	8.7%
% 65+ households spend >35% of income on housing (owner)		18.4%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$26,196	\$25,284	1.04
Single, renter, good health	\$27,264	\$26,400	1.03
Couple, homeowner without mortgage, good health	\$38,160	\$37,128	1.03
Couple, renter, good health	\$39,228	\$38,244	1.03

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are not due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019