

Somersworth is a city located in Strafford County that is bordered to the northeast by the Salmon Falls River. There are 1,392 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of prostate cancer, and cataract. However, they had higher rates of obesity, depression, anxiety disorders, post-traumatic stress disorder, tobacco use disorders, diabetes, chronic obstructive pulmonary disease, asthma, ischemic heart disease, congestive heart failure, chronic kidney disease, liver diseases, and fibromyalgia/chronic pain/fatigue. They are more likely to take the health promotion steps of getting screened for high cholesterol and colorectal cancer and always wearing a seatbelt. Community resources to support healthy aging include 131 primary care providers, a hospital, and a nursing home within 5 miles, 9 home health agencies, an assisted living site, a public library, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		11,756	1,327,503
Population 60 years or older as % of total population		17.9%	22.7%
Total population 60 years or older		2,110	301,630
Population 65 years or older as % of total population		11.8%	15.8%
Total population 65 years or older		1,392	210,385
% 65-74 years		59.1%	58.5%
% 75-84 years		27.2%	28.6%
% 85 years or older		13.8%	12.9%
Gender (65+ population)			
% female		59.3%	54.7%
Race/Ethnicity (65+ population)			
% White		94.0%	97.7%
% African American		1.4%	0.5%
% Asian		2.4%	0.9%
% Other		2.2%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		60.1%	58.5%
% divorced/separated		10.1%	14.0%
% widowed		22.6%	22.9%
% never married		7.3%	4.6%
Education (65+ population)			
% with less than high school education		13.3%	12.3%
% with high school or some college		72.0%	57.1%
% with college degree		14.7%	30.6%
% of 65+ population living alone		26.0%	26.1%
% of 65+ population who speak only English at home		83.6%	91.3%
% of 65+ population who are veterans of military service		26.4%	24.8%
Age-sex adjusted 1-year mortality rate	W	5.6%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		2.0%	3.6%
% moved from different county in New Hampshire		0.0%	1.0%
% moved from different state		1.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		77.2%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		25.4%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		61.9%	58.6%
% 60+ met CDC guidelines for both types of physical activities		21.6%	19.9%
% 60+ getting recommended hours of sleep		69.3%	66.4%
% 60+ injured in a fall within last 12 months		12.2%	10.4%
% 65+ had hip fracture		2.7%	3.3%
% 60+ with self-reported fair or poor health status		13.4%	16.5%
% 60+ with 15+ physically unhealthy days last month		9.8%	12.5%
% 60+ with physical exam/check-up in past year		90.0%	86.5%
% 60+ met CDC preventive health screening goals		44.3%	40.3%
% 60+ flu shot past year		61.6%	59.3%
% 65+ with pneumonia vaccine		79.8%	77.8%
% 60+ with cholesterol screening	В	98.6%	95.3%
% 60+ women with a mammogram within last 2 years		78.2%	79.3%
% 60+ with colorectal cancer screening	В	83.6%	77.0%
% 60+ with HIV test		10.9%	13.5%
% 60+ current smokers		6.4%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		26.9%	29.0%
% 60+ with annual dental exam		80.0%	75.7%
# of dentists per 100,000 persons (all ages) (county)		68	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	20.2%
% 60+ self-reported obese		24.4%	27.2%
% 65+ clinically diagnosed obese	W	21.5%	16.7%
% 65+ with high cholesterol		73.1%	72.2%
% 60+ excessive drinking		10.6%	9.2%
% 65+ with poor supermarket access		39.0%	28.4%

% 60+ with 15+ days poor mental health last month 6.9% 6.9% % 65+ with depression W 32.3% 28.8% % 65+ with anxiety disorders W 26.0% 21.9% % 65+ with bipolar disorders 3.8% 3.11% % 65+ with bipost-traumatic stress disorder W 2.7% 1.4% % 65+ with bipost-traumatic stress disorders I.0% 1.1% % 65+ with post-traumatic stress disorders I.0% 1.1% % 65+ with post-traumatic stress disorders I.0% 1.1% % 65+ with schizophrenia & other psychotic disorders I.0% 1.1% % 65+ with schizophrenia & other psychotic disorders I.0% 1.1% % 65+ with substance use disorders W 14.3 10.4% CHRONIC Discrass W 14.3 10.4% CHRONIC DISCrass W 36.0% 22.2% % 65+ with Alzheimer's disease or related dementias 12.6% 12.6% 65+ with disbetes W 36.0% 22.2% 65+ with hit disbetes W 27.0% 20.5% 65+	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression W 32.3% 28.8% % 65+ with anxiety disorders W 26.0% 21.9% % 65+ with bipolar disorders J.8% 3.1% % 65+ with post-traumatic stress disorder W 2.7% 4.4% % 65+ with post-traumatic stress disorders I.0% 1.1% % 65+ with post-portal & other psychotic disorders I.0% 1.1% % 65+ with schizophrenia & other psychotic disorders I.0% 1.1% % 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with biscace use disorders W 14.3% 10.4% CHRONIC DISEASE W 14.3% 10.4% CHRONIC DISEASE W 36.0% 28.2% 65+ with disbetes W 36.0% 28.2% 65+ with stroke 10.5% 10.8% 65+ with stroke 10.5% 10.8% 65+ with stroke W 27.0% 20.5% 65+ with hypertension 72.5% 70.2% 65+ with hypertension 72.5% 70.2%	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders W 26.0% 21.9% % 65+ with bipolar disorders 3.8% 3.1% % 65+ with post-traumatic stress disorder W 2.7% 1.4% % 65+ with schizophrenia & other psychotic disorders 6.1% 4.9% % 65+ with personality disorders 1.0% 1.1% # opioid deaths (all ages) (county) 142 1.279 % 65+ with bubstance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with batchologed 8.0 1.1.0% 1.2.0% 65+ with disorders W 36.0% 28.2% 65+ with batchologed W 21.0% 20.5% 65+ with batchologed W 21.0%<	% 60+ with 15+ days poor mental health last month		6.9%	6.9%
% 65+ with bipolar disorders 3.8% 3.1% % 65+ with post-traumatic stress disorder W 2.7% 1.4% % 65+ with personality disorders 6.1% 4.9% % 65+ with personality disorders 1.0% 1.1% # oploid deaths (all ages) (county) 142 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% % 65+ with bush substance use disorders W 14.3% 10.4% CHRONIC DISEASE *** *** 12.6% 12.0% % 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.8% % 65+ with stroke 10.5% 10.8% <	% 65+ with depression	W	32.3%	28.8%
% 65+ with post-traumatic stress disorder W 2.7% 1.4% % 65+ with schizophrenia & other psychotic disorders 6.1% 4.9% % 65+ with personality disorders 1.0% 1.1% 4 opioid deaths (all ages) (county) 142 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% % 65+ with thobacco use disorders W 14.3% 10.4% CHRONIC DISEASE W 12.6% 12.0% 65+ with diabetes W 36.0% 28.2% 65+ with stroke 10.5% 10.8% 65+ with stroke 10.5% 10.8% 65+ with hasthma W 27.0% 20.5% 65+ with asthma W 15.8% 13.0% 65+ with pertension 72.5% 70.2% 65+ with asthma W 15.8% 13.0% 65+ with ischemic heart disease W 38.1% 34.3% 65+ with congestive heart failure W 20.4% 17.7% 65+ with congestive heart failure W 20.4% 17.7% 65+ with osteoporrosis 17.6%<	% 65+ with anxiety disorders	W	26.0%	21.9%
% 65+ with schizophrenia & other psychotic disorders 6.1% 4.9% % 65+ with personality disorders 1.0% 1.1% # opioid deaths (all ages) (county) 142 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% 65+ with tobacco use disorders W 14.3% 10.4% CHRONIC DISEASE 8 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with stroke 10.5% 10.8% % 65+ with stroke 10.5% 10.8% % 65+ with asthma W 27.0% 20.5% % 65+ with obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with asthma W 15.8% 13.0% % 65+ with phypertension 72.5% 70.2% % 65+ with in schemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with peripheral vascular disease W 38.1% 14.7% % 65+ with beripheral vascular disease 17.6% <	% 65+ with bipolar disorders		3.8%	3.1%
% 65+ with personality disorders 1.0% 1.1% # opioid deaths (all ages) (county) 142 1,279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% % 65+ with tobacco use disorders W 14.3% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with diabetes W 36.0% 28.2% % 65+ with diabetes W 36.0% 28.2% % 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.8% % 65+ with toroic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with pypertension 72.5% 70.2% % 65+ with pypertension 72.5% 70.2% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with osteoparbritis/rheumatoid arthritis 50.0% 49.1% % 65+ with peripheral vascular disease 16.4% 14.7	% 65+ with post-traumatic stress disorder	W	2.7%	1.4%
# opioid deaths (all ages) (county) % 65+ with substance use disorders (drug use +/or alcohol abuse) % 65+ with tobacco use disorders W 14.3% 10.4% CHRONIC DISEASE W 36.0% 28.2% % 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.5% % 65+ with stroke % 65+ with orionic obstructive pulmonary disease W 27.0% 20.5% % 65+ with hypertension % 65+ with hypertension % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with stroke indiabetes W 38.1% 34.3% % 65+ with peripheral vascular disease W 38.1% 34.3% % 65+ with osteoarthritis/rheumatoid arthritis % 65+ with osteoarthritis/rheumatoid arthritis % 65+ with lung cancer \$ 2.1% 16.6% % 65+ with lung cancer \$ 2.7% 2.4% % 65+ with lung cancer \$ 3.6% 9.8% % 65+ with lung cancer \$ 3.6% 9.8% % 65+ with propriate cancer \$ 6.6 with propriate cancer \$ 6.6 with propriate cancer \$ 6.7 with propriate can	% 65+ with schizophrenia & other psychotic disorders		6.1%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 7.0% 5.5% % 65+ with tobacco use disorders W 14.3% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with stroke 10.5% 10.8% % 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with chronic obstructive pulmonary disease W 15.8% 13.0% % 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ with pypertension 72.5% 70.2% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with torial librillation 14.4% 14.4% % 65+ with osteoparthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoparosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with leukemias and lymphomas <	% 65+ with personality disorders		1.0%	1.1%
% 65+ with tobacco use disorders W 14.3% 10.4% CHRONIC DISEASE ************************************	# opioid deaths (all ages) (county)		142	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.8% % 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with pypertension 72.5% 70.2% % 65+ with hypertension 72.5% 70.2% % 65+ with pypertension 72.5% 70.2% % 65+ with loybertension 72.5% 70.2% % 65+ with pypertension 72.5% 70.2% % 65+ with loybertension 72.5% 70.2% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoparthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoparthritis/rheumatoid arthritis	% 65+ with substance use disorders (drug use +/or alcohol abuse)		7.0%	5.5%
% 65+ with Alzheimer's disease or related dementias 12.6% 12.0% % 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.8% % 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with orgestive heart failure W 20.4% 17.7% % 65+ with orgestive heart failure W 20.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoporosis 17.6% 17.4% % 65+ with osteoporosis 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4%	% 65+ with tobacco use disorders	W	14.3%	10.4%
% 65+ with diabetes W 36.0% 28.2% % 65+ with stroke 10.5% 10.8% % 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with lung cancer 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ with prostate cancer 8.6% 9.8% % 65+ women with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypot	CHRONIC DISEASE			
% 65+ with stroke 10.5% 10.8% % 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with preast cancer 8.6% 9.8% % 65+ women with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypothyroidism 20.3% 20.3% % 65+ with	% 65+ with Alzheimer's disease or related dementias		12.6%	12.0%
% 65+ with chronic obstructive pulmonary disease W 27.0% 20.5% % 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.1% 1.6% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with nemia 35.7% 37.3% % 65+ with c	% 65+ with diabetes	W	36.0%	28.2%
% 65+ with asthma W 15.8% 13.0% % 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoprosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with bolon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver di	% 65+ with stroke		10.5%	10.8%
% 65+ with hypertension 72.5% 70.2% % 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoprosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ mith benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypothyroidism 20.3% 20.3% % 65+ with nemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with chronic obstructive pulmonary disease	W	27.0%	20.5%
% 65+ ever had a heart attack 5.2% 4.5% % 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with long cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with liver diseases W 27.6% 22.3% % 65+ with liver diseases W 29.0% 6.9%	% 65+ with asthma	W	15.8%	13.0%
% 65+ with ischemic heart disease W 38.1% 34.3% % 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 36.8% 36.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with hypertension		72.5%	70.2%
% 65+ with congestive heart failure W 20.4% 17.7% % 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ ever had a heart attack		5.2%	4.5%
% 65+ with atrial fibrillation 14.4% 14.4% % 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with nemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with ischemic heart disease	W	38.1%	34.3%
% 65+ with peripheral vascular disease 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with congestive heart failure	W	20.4%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis 50.0% 49.1% % 65+ with osteoporosis 17.6% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with atrial fibrillation		14.4%	14.4%
% 65+ with osteoporosis 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with peripheral vascular disease		16.4%	14.7%
% 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with osteoarthritis/rheumatoid arthritis		50.0%	49.1%
% 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with osteoporosis		17.6%	17.4%
% 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with leukemias and lymphomas		2.0%	2.0%
% 65+ women with breast cancer 8.6% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with lung cancer		2.1%	1.6%
% 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with colon cancer		2.7%	2.4%
% 65+ men with prostate cancer B 8.7% 11.5% % 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ women with breast cancer		8.6%	9.8%
% 65+ with benign prostatic hyperplasia 38.6% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ women with endometrial cancer		1.7%	1.7%
% 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ men with prostate cancer	В	8.7%	11.5%
% 65+ with hypothyroidism 20.3% 20.8% % 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with benign prostatic hyperplasia		38.6%	36.8%
% 65+ with anemia 35.7% 37.3% % 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with HIV/AIDS		0.04%	0.05%
% 65+ with chronic kidney disease W 27.6% 22.3% % 65+ with liver diseases W 9.0% 6.9%	% 65+ with hypothyroidism		20.3%	20.8%
% 65+ with liver diseases W 9.0% 6.9%	% 65+ with anemia		35.7%	37.3%
% 65+ with liver diseases W 9.0% 6.9%	% 65+ with chronic kidney disease	W	27.6%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue W 26.6% 18.6%	% 65+ with liver diseases	W	9.0%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	W	26.6%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		4.4%	4.0%
% 65+ with epilepsy		2.7%	2.1%
% 65+ with traumatic brain injury		1.6%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma		22.6%	22.9%
% 65+ with cataract	В	54.2%	61.2%
% 65+ with pressure ulcer or chronic ulcer		8.8%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	W	60.2%	54.4%
% 65+ with 0 chronic conditions	В	12.8%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		18.7%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		15.7%	14.4%
% 65+ with self-reported vision difficulty		6.8%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		1.2%	0.9%
% 65+ with self-reported cognition difficulty		6.8%	6.9%
% 65+ with self-reported ambulatory difficulty		22.5%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.4%	3.2%
% 65+ with self-reported self-care difficulty		5.1%	5.6%
% 65+ with self-reported independent living difficulty		9.2%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	3.7%	7.9%
% dually eligible for Medicare and Medicaid	*	10.0%	7.5%
% 60+ with a regular doctor	В	98.2%	95.8%
% 60+ who did not see doctor when needed due to cost		4.2%	5.4%
# of primary care providers within 5 miles		131	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		1	74
# of home health agencies		9	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	7.0	6.3
Emergency room visits/1000 persons 65+ years per year	*	666	572
Part D monthly prescription fills per person per year	*	54.5	49.1
Home health visits per year	*	3.3	2.5
Durable medical equipment claims per year	*	3.1	2.0
Inpatient hospital stays/1000 persons 65+ years per year		244	237
Medicare inpatient hospital readmissions (as % of admissions)		14.9%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		88	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports		3.5%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		N/A	N/A
% of grandparents raising grandchildren		1.8%	0.8%
% of grandparents who live with grandchildren		3.6%	2.5%
# of assisted living sites		1	134
% of vacant homes in community		6.9%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		97.0%	93.0%
% 60+ who used Internet in last month		80.1%	77.6%
Voter participation rate in 2018 election (age 18+)		45.6%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		420	207
Homicide rate /100,000 persons (county)		1	1
# firearm fatalities (county)		52	586
Property crime rate /100,000 persons		3,622	2,012
% 65+ who own a motor vehicle		82.9%	91.0%
% 60+ who always drive wearing a seatbelt	В	83.1%	77.1%
# of fatal crashes involving adult age 60+/town		1	151
# of fatal crashes involving adult age 60+/county		17	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		7.7%	5.4%
% 60+ receiving food stamps past year		6.7%	5.7%
% 65+ employed past year		19.2%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		19.5%	18.2%
% households with annual income \$20,000-\$49,999		42.6%	36.5%
% households with annual income > \$50,000		37.9%	45.3%
% 60+ own home		65.7%	79.9%
% 60+ have mortgage on home		30.2%	35.3%
% 65+ households spend >35% of income on housing (renter)		5.6%	8.7%
% 65+ households spend >35% of income on housing (owner)		21.8%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$24,936	\$25,284	0.99
Single, renter, good health	\$26,112	\$26,400	0.99
Couple, homeowner without mortgage, good health	\$36,504	\$37,128	0.98
Couple, renter, good health	\$37,680	\$38,244	0.99

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are not due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019