## TUFTS Health Plan

## **Derry (Rockingham)**

Derry is an urban town located in southeastern New Hampshire, nicknamed "Spacetown." There are 3,776 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of prostate cancer, benign prostatic hyperplasia, and cataracts. However, they had higher rates of obesity, anxiety disorders, personality disorders, substance and tobacco use disorders, diabetes, chronic obstructive pulmonary disease, asthma, hypertension, congestive heart failure, chronic kidney disease, and glaucoma. They are more likely to take the health promotion step of having a regular mammogram. Community resources to support healthy aging include 2 senior centers, a hospital within 5 miles, 2 nursing homes within 5 miles, 15 home health agencies, 2 public libraries, an adult day health center, a assisted living site, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		33,255	1,327,503
Population 60 years or older as % of total population		17.1%	22.7%
Total population 60 years or older		5,680	301,630
Population 65 years or older as % of total population		11.4%	15.8%
Total population 65 years or older		3,776	210,385
% 65-74 years		60.1%	58.5%
% 75-84 years		29.6%	28.6%
% 85 years or older		10.3%	12.9%
Gender (65+ population)			
% female		54.5%	54.7%
Race/Ethnicity (65+ population)			
% White		99.5%	97.7%
% African American		0.0%	0.5%
% Asian		0.0%	0.9%
% Other		0.5%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		56.4%	58.5%
% divorced/separated		13.5%	14.0%
% widowed		24.1%	22.9%
% never married		6.0%	4.6%
Education (65+ population)			
% with less than high school education		15.5%	12.3%
% with high school or some college		59.8%	57.1%
% with college degree		24.7%	30.6%
% of 65+ population living alone		26.2%	26.1%
% of 65+ population who speak only English at home		96.4%	91.3%
% of 65+ population who are veterans of military service		22.1%	24.8%
Age-sex adjusted 1-year mortality rate		4.4%	4.1%
			54654

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		2.3%	3.6%
% moved from different county in New Hampshire		1.4%	1.0%
% moved from different state		1.2%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		70.1%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		27.2%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		52.0%	58.6%
% 60+ met CDC guidelines for both types of physical activities		20.4%	19.9%
% 60+ getting recommended hours of sleep		67.5%	66.4%
% 60+ injured in a fall within last 12 months		12.3%	10.4%
% 65+ had hip fracture		2.7%	3.3%
% 60+ with self-reported fair or poor health status		13.2%	16.5%
% 60+ with 15+ physically unhealthy days last month		12.1%	12.5%
% 60+ with physical exam/check-up in past year		87.9%	86.5%
% 60+ met CDC preventive health screening goals		47.8%	40.3%
% 60+ flu shot past year		67.1%	59.3%
% 65+ with pneumonia vaccine		79.7%	77.8%
% 60+ with cholesterol screening		96.4%	95.3%
% 60+ women with a mammogram within last 2 years	В	88.9%	79.3%
% 60+ with colorectal cancer screening		82.5%	77.0%
% 60+ with HIV test		16.1%	13.5%
% 60+ current smokers		6.1%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		30.8%	29.0%
% 60+ with annual dental exam		80.9%	75.7%
# of dentists per 100,000 persons (all ages) (county)		72	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		21.6%	20.2%
% 60+ self-reported obese		29.0%	27.2%
% 65+ clinically diagnosed obese	W	20.7%	16.7%
% 65+ with high cholesterol		73.1%	72.2%
% 60+ excessive drinking		7.8%	9.2%
% 65+ with poor supermarket access		42.3%	28.4%

BEHAVIORAL HEALTH           % 60+ with 15+ days poor mental health last month         9.4%           % 65+ with depression         30.1%           % 65+ with anxiety disorders         W 24.9%           % 65+ with bipolar disorders         3.8%           % 65+ with post-traumatic stress disorder         1.7%           % 65+ with personality disorders         W 1.7%           # opioid deaths (all ages) (county)         276           % 65+ with substance use disorders (drug use +/or alcohol abuse)         W 7.0%           % 65+ with tobacco use disorders         W 13.6%           CHRONIC DISEASE         W 33.2%           % 65+ with Alzheimer's disease or related dementias         12.8%           % 65+ with diabetes         W 33.2%           % 65+ with chronic obstructive pulmonary disease         W 25.2%           % 65+ with sathma         W 17.1%           % 65+ with apart attack         5.3%           % 65+ with pypertension         W 73.3%           % 65+ with porpheral dae heart attack         5.3%           % 65+ with pypertension heart disease         35.8%           % 65+ with osteopersive heart failure         W 20.0%           % 65+ with osteopersive heart failure         W 20.0%           % 65+ with osteoporosis         16.2%	6.9%
% 65+ with depression         30.1%           % 65+ with anxiety disorders         W         24.9%           % 65+ with bipolar disorders         3.8%           % 65+ with post-traumatic stress disorder         1.7%           % 65+ with post-traumatic stress disorders         5.7%           % 65+ with personality disorders         W         1.7%           # opioid deaths (all ages) (county)         276           % 65+ with substance use disorders (drug use +/or alcohol abuse)         W         7.0%           % 65+ with tobacco use disorders         W         13.6%           CHRONIC DISEASE         W         33.2%           % 65+ with Alzheimer's disease or related dementias         12.8%           % 65+ with stroke         11.5%           % 65+ with stroke         11.5%           % 65+ with hypertension         W         25.2%           % 65+ with hypertension         W         73.3%           % 65+ ever had a heart attack         5.3%           % 65+ with ischemic heart disease         35.8%           % 65+ with peripheral vascular disease         35.8%           % 65+ with peripheral vascular disease         14.8%           % 65+ with osteoporosis         16.2%           % 65+ with leukemias and lymphomas         16.6%     <	
% 65+ with anxiety disorders         W         24.9%           % 65+ with bipolar disorders         3.8%           % 65+ with post-traumatic stress disorder         1.7%           % 65+ with schizophrenia & other psychotic disorders         5.7%           % 65+ with personality disorders         W         1.7%           # opioid deaths (all ages) (county)         276           % 65+ with substance use disorders (drug use +/or alcohol abuse)         W         7.0%           % 65+ with tobacco use disorders         W         13.6%           CHRONIC DISEASE         W         33.2%           % 65+ with Alzheimer's disease or related dementias         12.8%           % 65+ with diabetes         W         33.2%           % 65+ with othoronic obstructive pulmonary disease         W         25.2%           % 65+ with asthma         W         17.1%           % 65+ with hypertension         W         73.3%           % 65+ with hypertension         W         73.3%           % 65+ with ischemic heart disease         35.8%           % 65+ with congestive heart failure         W         20.0%           % 65+ with peripheral vascular disease         14.8%           % 65+ with osteoporosis         16.2%           % 65+ with lung cancer         1.	28.8%
% 65+ with bipolar disorders       3.8%         % 65+ with post-traumatic stress disorder       1.7%         % 65+ with schizophrenia & other psychotic disorders       5.7%         % 65+ with personality disorders       W       1.7%         # opioid deaths (all ages) (county)       276         % 65+ with substance use disorders (drug use +/or alcohol abuse)       W       7.0%         % 65+ with tobacco use disorders       W       13.6%         CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with hypertension       W       73.3%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with orgestive heart failure       W       20.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with luurg cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer	_0.0,0
% 65+ with post-traumatic stress disorder         1.7%           % 65+ with schizophrenia & other psychotic disorders         5.7%           % 65+ with personality disorders         W         1.7%           # opioid deaths (all ages) (county)         276           % 65+ with substance use disorders (drug use +/or alcohol abuse)         W         7.0%           % 65+ with tobacco use disorders         W         13.6%           CHRONIC DISEASE           % 65+ with Alzheimer's disease or related dementias         12.8%           % 65+ with stroke         11.5%           % 65+ with chronic obstructive pulmonary disease         W         25.2%           % 65+ with hypertension         W         17.1%           % 65+ ever had a heart attack         5.3%           % 65+ ever had a heart disease         35.8%           % 65+ with congestive heart disease         35.8%           % 65+ with peripheral vascular disease         14.0%           % 65+ with osteoporosis         16.2%           % 65+ with leukemias and lymphomas         1.6%           % 65+ with lung cancer         1.9%           % 65+ with colon cancer         2.0%           % 65+ women with breast cancer         9.9%	21.9%
% 65+ with schizophrenia & other psychotic disorders       5.7%         % 65+ with personality disorders       W       1.7%         # opioid deaths (all ages) (county)       276         % 65+ with substance use disorders (drug use +/or alcohol abuse)       W       7.0%         % 65+ with tobacco use disorders       W       13.6%         CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with long cancer       1.9%         % 65+ women with breast cancer       9.9%	3.1%
% 65+ with personality disorders       W       1.7%         # opioid deaths (all ages) (county)       276         % 65+ with substance use disorders (drug use +/or alcohol abuse)       W       7.0%         % 65+ with bobacco use disorders       W       13.6%         CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with lung cancer       1.9%         % 65+ with lung cancer       1.9%         % 65+ women with breast cancer       9.9%	1.4%
# opioid deaths (all ages) (county)  % 65+ with substance use disorders (drug use +/or alcohol abuse)  W 7.0%  % 65+ with tobacco use disorders  W 13.6%  CHRONIC DISEASE  % 65+ with Alzheimer's disease or related dementias  12.8%  % 65+ with diabetes  W 33.2%  % 65+ with stroke  11.5%  % 65+ with chronic obstructive pulmonary disease  W 25.2%  % 65+ with asthma  W 17.1%  % 65+ with hypertension  W 73.3%  % 65+ ever had a heart attack  5.3%  % 65+ with ischemic heart disease  % 65+ with congestive heart failure  W 20.0%  % 65+ with atrial fibrillation  14.0%  % 65+ with peripheral vascular disease  14.8%  % 65+ with osteoarthritis/rheumatoid arthritis  47.1%  % 65+ with osteoporosis  16.2%  % 65+ with lung cancer  1.9%  % 65+ with lung cancer  9.9%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse)       W       7.0%         % 65+ with tobacco use disorders       W       13.6%         CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with osteoporosis       14.8%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	1.1%
% 65+ with tobacco use disorders       W       13.6%         CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ ever had a heart disease       35.8%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	1,279
CHRONIC DISEASE         % 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	5.5%
% 65+ with Alzheimer's disease or related dementias       12.8%         % 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoporosis       16.2%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	10.4%
% 65+ with diabetes       W       33.2%         % 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoarthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	
% 65+ with stroke       11.5%         % 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoarthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	12.0%
% 65+ with chronic obstructive pulmonary disease       W       25.2%         % 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoarthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	28.2%
% 65+ with asthma       W       17.1%         % 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoparthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	10.8%
% 65+ with hypertension       W       73.3%         % 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoarthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	20.5%
% 65+ ever had a heart attack       5.3%         % 65+ with ischemic heart disease       35.8%         % 65+ with congestive heart failure       W       20.0%         % 65+ with atrial fibrillation       14.0%         % 65+ with peripheral vascular disease       14.8%         % 65+ with osteoarthritis/rheumatoid arthritis       47.1%         % 65+ with osteoporosis       16.2%         % 65+ with leukemias and lymphomas       1.6%         % 65+ with lung cancer       1.9%         % 65+ with colon cancer       2.0%         % 65+ women with breast cancer       9.9%	13.0%
% 65+ with ischemic heart disease35.8%% 65+ with congestive heart failureW20.0%% 65+ with atrial fibrillation14.0%% 65+ with peripheral vascular disease14.8%% 65+ with osteoarthritis/rheumatoid arthritis47.1%% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	70.2%
% 65+ with congestive heart failureW20.0%% 65+ with atrial fibrillation14.0%% 65+ with peripheral vascular disease14.8%% 65+ with osteoarthritis/rheumatoid arthritis47.1%% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	4.5%
% 65+ with atrial fibrillation14.0%% 65+ with peripheral vascular disease14.8%% 65+ with osteoarthritis/rheumatoid arthritis47.1%% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	34.3%
% 65+ with peripheral vascular disease14.8%% 65+ with osteoarthritis/rheumatoid arthritis47.1%% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis47.1%% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	14.4%
% 65+ with osteoporosis16.2%% 65+ with leukemias and lymphomas1.6%% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	14.7%
% 65+ with leukemias and lymphomas  1.6%  % 65+ with lung cancer  1.9%  % 65+ with colon cancer  2.0%  % 65+ women with breast cancer  9.9%	49.1%
% 65+ with lung cancer1.9%% 65+ with colon cancer2.0%% 65+ women with breast cancer9.9%	17.4%
% 65+ with colon cancer 2.0% % 65+ women with breast cancer 9.9%	2.0%
% 65+ women with breast cancer 9.9%	1.6%
	2.4%
% 65+ women with endometrial cancer 1.8%	9.8%
	1.7%
% 65+ men with prostate cancer B 9.5%	11.5%
% 65+ with benign prostatic hyperplasia B 29.2%	36.8%
% 65+ with HIV/AIDS 0.04%	0.05%
% 65+ with hypothyroidism 22.2%	20.8%
% 65+ with anemia 39.1%	37.3%
% 65+ with chronic kidney disease W 27.8%	22.3%
% 65+ with liver diseases 6.8%	6.9%
% 65+ with fibromyalgia, chronic pain and fatigue 17.8%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		4.2%	4.0%
% 65+ with epilepsy		2.1%	2.1%
% 65+ with traumatic brain injury		1.4%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	W	25.6%	22.9%
% 65+ with cataract	В	58.9%	61.2%
% 65+ with pressure ulcer or chronic ulcer		7.2%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	W	57.5%	54.4%
% 65+ with 0 chronic conditions		11.2%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty	В	10.7%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		14.3%	14.4%
% 65+ with self-reported vision difficulty		3.8%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		1.1%	0.9%
% 65+ with self-reported cognition difficulty		5.6%	6.9%
% 65+ with self-reported ambulatory difficulty		18.9%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.0%	3.2%
% 65+ with self-reported self-care difficulty		6.3%	5.6%
% 65+ with self-reported independent living difficulty		14.7%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		8.8%	7.9%
% dually eligible for Medicare and Medicaid		8.3%	7.5%
% 60+ with a regular doctor		97.2%	95.8%
% 60+ who did not see doctor when needed due to cost		2.7%	5.4%
# of primary care providers within 5 miles		95	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		2	74
# of home health agencies		15	49
# of community health centers		0	22
# of adult day health centers		1	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	7.9	6.3
Emergency room visits/1000 persons 65+ years per year	*	645	572
Part D monthly prescription fills per person per year	*	52.1	49.1
Home health visits per year		2.3	2.5
Durable medical equipment claims per year		2.1	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	279	237
Medicare inpatient hospital readmissions (as % of admissions)	*	20.2%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		91	76
# skilled nursing home Medicare beds/1000 persons 65+ years		43	33
% 65+ getting Medicaid long term services and supports	*	4.8%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		2	44
Air pollution: annual # of unhealthy days for 65+ (county)		2	N/A
% of grandparents raising grandchildren		0.5%	0.8%
% of grandparents who live with grandchildren		2.4%	2.5%
# of assisted living sites		1	134
% of vacant homes in community		4.8%	16.0%
# of universities and community colleges		0	41
# of public libraries		2	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		99.0%	93.0%
% 60+ who used Internet in last month		80.1%	77.6%
Voter participation rate in 2018 election (age 18+)		44.1%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		160	207
Homicide rate /100,000 persons (county)		1	1
# firearm fatalities (county)		92	586
Property crime rate /100,000 persons		1,906	2,012
% 65+ who own a motor vehicle		88.9%	91.0%
% 60+ who always drive wearing a seatbelt		83.0%	77.1%
# of fatal crashes involving adult age 60+/town		3	151
# of fatal crashes involving adult age 60+/county		31	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		4.2%	5.4%
% 60+ receiving food stamps past year		7.1%	5.7%
% 65+ employed past year		24.1%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		17.7%	18.2%
% households with annual income \$20,000-\$49,999		43.1%	36.5%
% households with annual income > \$50,000		39.3%	45.3%
% 60+ own home		70.3%	79.9%
% 60+ have mortgage on home		40.8%	35.3%
% 65+ households spend >35% of income on housing (renter)		17.1%	8.7%
% 65+ households spend >35% of income on housing (owner)		23.8%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,536	\$25,284	1.01
Single, renter, good health	\$26,712	\$26,400	1.01
Couple, homeowner without mortgage, good health	\$37,092	\$37,128	1.00
Couple, renter, good health	\$38,268	\$38,244	1.00

## TECHNICAL NOTES

\*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are not due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an \*. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

## Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019