Lee (Strafford)



Lee is a rural town located in Sullivan County and home to the notable Bedrock Gardens. There are 570 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of high cholesterol, depression, Alzheimer's disease or related dementias, chronic obstructive pulmonary disease, hypertension, ischemic heart disease, congestive heart failure, peripheral vascular disease, osteoarthritis/rheumatoid arthritis, osteoporosis, hypothyroidism, anemia, chronic kidney disease, liver diseases, and cataracts. However, they had a higher rate of schizophrenia and other psychotic disorders. They are more likely to take the health promotion steps of having a cholesterol screening or colorectal cancer screening. Community resources to support healthy aging include 3 primary care providers within 5 miles, 10 home health agencies, an assisted living site, a public library, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		4,388	1,327,503
Population 60 years or older as % of total population		20.1%	22.7%
Total population 60 years or older		884	301,630
Population 65 years or older as % of total population		13.0%	15.8%
Total population 65 years or older		570	210,385
% 65-74 years		64.2%	58.5%
% 75-84 years		35.8%	28.6%
% 85 years or older		0.0%	12.9%
Gender (65+ population)			
% female		54.9%	54.7%
Race/Ethnicity (65+ population)			
% White		100.0%	97.7%
% African American		0.0%	0.5%
% Asian		0.0%	0.9%
% Other		0.0%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		75.1%	58.5%
% divorced/separated		8.9%	14.0%
% widowed		13.2%	22.9%
% never married		2.8%	4.6%
Education (65+ population)			
% with less than high school education		4.9%	12.3%
% with high school or some college		43.0%	57.1%
% with college degree		52.1%	30.6%
% of 65+ population living alone		18.4%	26.1%
% of 65+ population who speak only English at home		93.2%	91.3%
% of 65+ population who are veterans of military service		20.4%	24.8%
Age-sex adjusted 1-year mortality rate		3.1%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		0.0%	3.6%
% moved from different county in New Hampshire		0.0%	1.0%
% moved from different state		0.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		77.2%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		25.4%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		61.9%	58.6%
% 60+ met CDC guidelines for both types of physical activities		21.6%	19.9%
% 60+ getting recommended hours of sleep		69.3%	66.4%
% 60+ injured in a fall within last 12 months		12.2%	10.4%
% 65+ had hip fracture		2.9%	3.3%
% 60+ with self-reported fair or poor health status		13.4%	16.5%
% 60+ with 15+ physically unhealthy days last month		9.8%	12.5%
% 60+ with physical exam/check-up in past year		90.0%	86.5%
% 60+ met CDC preventive health screening goals		44.3%	40.3%
% 60+ flu shot past year		61.6%	59.3%
% 65+ with pneumonia vaccine		79.8%	77.8%
% 60+ with cholesterol screening	В	98.6%	95.3%
% 60+ women with a mammogram within last 2 years		78.2%	79.3%
% 60+ with colorectal cancer screening	В	83.6%	77.0%
% 60+ with HIV test		10.9%	13.5%
% 60+ current smokers		6.4%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		26.9%	29.0%
% 60+ with annual dental exam		80.0%	75.7%
# of dentists per 100,000 persons (all ages) (county)		68	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	20.2%
% 60+ self-reported obese		24.4%	27.2%
% 65+ clinically diagnosed obese		15.6%	16.7%
% 65+ with high cholesterol	В	66.3%	72.2%
% 60+ excessive drinking		10.6%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

AVIORAL HEALTH + with 15+ days poor mental health last month + with depression F with anxiety disorders + with bipolar disorders + with post-traumatic stress disorder + with schizophrenia & other psychotic disorders + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with diabetes + with stroke + with hypertension + with hypertension + ever had a heart attack + with congestive heart failure + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with lool cancer + women with breast cancer	18.6% 2.7% 1.1% 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	28.8% 21.9% 3.1% 1.4% 4.9% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.1% 2 1.279 5 5.5% 10.4% 12.0% 28.2% 10.8% 20.5% 13.0% 70.2% 4.5%
+ with depression E + with anxiety disorders + + with bipolar disorders + + with post-traumatic stress disorder + + with schizophrenia & other psychotic disorders W + with personality disorders W + with substance use disorders (drug use +/or alcohol abuse) + + with substance use disorders (drug use +/or alcohol abuse) + + with tobacco use disorders ONIC DISEASE + with diabetes + + with diabetes + + with stroke + + with hypertension E + with ischemic heart disease E + with congestive heart failure E + with congestive heart failure E + with peripheral vascular disease E + with osteoporosis E + with leukemias and lymphomas + + with colon cancer + + with colon cancer +	23.7% 18.6% 2.7% 1.1% 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	28.8% 21.9% 3.1% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.4% 1.1% 2 1.279 5 5.5% 10.4% 12.0% 2 28.2% 10.8% 20.5% 13.0% 20.5% 4.5%
 + with anxiety disorders + with bipolar disorders + with post-traumatic stress disorder + with schizophrenia & other psychotic disorders W + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with diabetes + with diabetes + with stroke + with stroke + with hypertension + ever had a heart attack + with congestive heart failure + with atrial fibrillation + with osteoorosis E + with lung cancer + with colon cancer + women with breast cancer 	18.6% 2.7% 1.1% 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	21.9% 3.1% 1.4% 1.4% 1.4% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 2.1,279 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1% 1.1%
 + with bipolar disorders + with post-traumatic stress disorder + with schizophrenia & other psychotic disorders W + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias + with diabetes + with stroke + with stroke + with hypertension + ever had a heart attack + with congestive heart failure + with peripheral vascular disease + with osteoorthritis/rheumatoid arthritis + with osteoortorsis + with lung cancer + with colon cancer + women with breast cancer 	2.7% 1.1% 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	3.1% 1.4% 4.9% 1.1% <t< td=""></t<>
 + with post-traumatic stress disorder + with schizophrenia & other psychotic disorders + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias + with diabetes + with stroke + with stroke + with asthma + ever had a heart attack + with congestive heart failure + with peripheral vascular disease + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with lung cancer + women with breast cancer 	1.1% 2 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	a 1.4% b 4.9% b 1.1% c 1,279 b 5.5% b 10.4% b 12.0% b 28.2% b 10.8% b 10.8% b 13.0% b 70.2% b 4.5%
 + with schizophrenia & other psychotic disorders + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with substance use disorders ONIC DISEASE + with Alzheimer's disease or related dementias E + with diabetes + with stroke + with stroke + with asthma + with hypertension + ever had a heart attack + with congestive heart failure + with peripheral vascular disease + with osteoorthritis/rheumatoid arthritis + with osteoporosis + with lung cancer + with colon cancer + women with breast cancer 	2 6.3% 0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	4.9% 1.1% 1.1% 1.279 5.5% 10.4% 11.1% 12.0% 22.12.0% 10.4% 10.4% 10.4% 10.4% 10.4% 10.4% 10.4% 11.0% 10.4% 10.4% 10.4% 11.0% 10.4% 10.8% 10.8% 10.2% 13.0% 14.5%
 + with personality disorders bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias E + with diabetes + with stroke + with chronic obstructive pulmonary disease E + with hypertension + ever had a heart attack + with congestive heart failure + with osteoporosis + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with colon cancer + women with breast cancer 	0.9% 142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	1.1% 1,279 5.5% 10.4% 112.0% 28.2% 10.8% 10.8% 113.0%
bid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias + with diabetes + with diabetes + with stroke + with chronic obstructive pulmonary disease + with asthma + with hypertension E + with asthma + with hypertension E + with ischemic heart disease + with ischemic heart disease + with congestive heart failure + with congestive heart failure + with peripheral vascular disease + with osteoporosis E + with osteoporosis E + with leukemias and lymphomas + with colon cancer + women with breast cancer E	142 4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	2 1,279 5.5% 5 10.4% 5 12.0% 5 28.2% 5 28.2% 5 10.8% 5 20.5% 5 13.0% 5 70.2% 5 4.5%
 + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias + with diabetes + with stroke + with chronic obstructive pulmonary disease + with asthma + with asthma + ever had a heart attack + with ischemic heart disease + with atrial fibrillation + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with colon cancer + women with breast cancer 	4.7% 7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	5.5% 10.4% 10.4% 28.2% 28.2% 10.8% 20.5% 13.0% 5 70.2% 5 4.5%
+ with tobacco use disorders ONIC DISEASE + with Alzheimer's disease or related dementias F with Alzheimer's disease or related dementias F with diabetes + with stroke + with chronic obstructive pulmonary disease F with asthma + with hypertension F ever had a heart attack + with ischemic heart disease F with congestive heart failure F with congestive heart failure F with peripheral vascular disease F with osteoporosis F with leukemias and lymphomas F with colon cancer F women with breast cancer F women with breast cancer F women with breast cancer F with osteoporosis F S S S S S S S S S S S S S S S S S S S	7.8% 9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	a 10.4% b 12.0% b 28.2% b 10.8% b 20.5% b 13.0% b 70.2% b 4.5%
ONIC DISEASE+ with Alzheimer's disease or related dementiasE+ with diabetes+ with stroke+ with chronic obstructive pulmonary diseaseE+ with asthma+ with hypertensionE+ ever had a heart attack+ with ischemic heart diseaseE+ with congestive heart failureE+ with atrial fibrillationE+ with osteoarthritis/rheumatoid arthritisE+ with leukemias and lymphomasE+ with loon cancerE+ women with breast cancerE	9.5% 25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	12.0% 28.2% 10.8% 20.5% 13.0% 70.2% 4.5%
 + with Alzheimer's disease or related dementias + with diabetes + with stroke + with chronic obstructive pulmonary disease + with asthma + with asthma + with hypertension + ever had a heart attack + with ischemic heart disease + with congestive heart failure + with atrial fibrillation + with osteoarthritis/rheumatoid arthritis + with osteoporosis + with leukemias and lymphomas + with colon cancer + women with breast cancer 	25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	28.2% 10.8% 20.5% 13.0% 570.2% 54.5%
 + with diabetes + with stroke + with chronic obstructive pulmonary disease + with asthma + with asthma + with hypertension + ever had a heart attack + with ischemic heart disease + with congestive heart failure + with congestive heart failure + with peripheral vascular disease + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with lung cancer + women with breast cancer 	25.8% 8.6% 15.7% 11.1% 63.8% 3.5%	28.2% 10.8% 20.5% 13.0% 570.2% 54.5%
 + with stroke + with chronic obstructive pulmonary disease + with asthma + with hypertension + ever had a heart attack + with ischemic heart disease + with ischemic heart disease + with congestive heart failure + with atrial fibrillation + with peripheral vascular disease + with osteoporosis + with osteoporosis + with leukemias and lymphomas + with colon cancer + women with breast cancer 	8.6% 15.7% 11.1% 63.8% 3.5%	10.8% 20.5% 13.0% 70.2% 4.5%
+ with chronic obstructive pulmonary diseaseE+ with asthmaE+ with hypertensionE+ ever had a heart attackE+ with ischemic heart diseaseE+ with congestive heart failureE+ with atrial fibrillationE+ with osteoarthritis/rheumatoid arthritisE+ with osteoporosisE+ with leukemias and lymphomasE+ with colon cancerE+ women with breast cancerE	15.7% 11.1% 63.8% 3.5%	20.5% 13.0% 70.2% 4.5%
+ with asthma+ with hypertension+ ever had a heart attack+ with ischemic heart disease+ with ischemic heart disease+ with congestive heart failure+ with congestive heart failure+ with atrial fibrillation+ with peripheral vascular disease+ with osteoarthritis/rheumatoid arthritis+ with osteoporosis+ with leukemias and lymphomas+ with lung cancer+ with colon cancer+ women with breast cancer	11.1% 63.8% 3.5%	13.0% 70.2% 4.5%
+ with hypertensionE+ ever had a heart attack+ with ischemic heart diseaseE+ with congestive heart failureE+ with atrial fibrillationE+ with peripheral vascular diseaseE+ with osteoarthritis/rheumatoid arthritisE+ with osteoporosisE+ with leukemias and lymphomasE+ with colon cancerE+ women with breast cancerE	63.8% 3.5%	5 70.2% 5 4.5%
+ ever had a heart attack + with ischemic heart disease E + with congestive heart failure E + with atrial fibrillation + with peripheral vascular disease E + with osteoarthritis/rheumatoid arthritis E + with osteoporosis E + with leukemias and lymphomas + with leukemias and lymphomas + with colon cancer + women with breast cancer E	3.5%	4.5%
+ with ischemic heart diseaseE+ with congestive heart failureE+ with congestive heart failureE+ with atrial fibrillationE+ with peripheral vascular diseaseE+ with osteoarthritis/rheumatoid arthritisE+ with osteoporosisE+ with leukemias and lymphomas+ with lung cancer+ with colon cancer+ women with breast cancerE		
+ with congestive heart failureE+ with atrial fibrillation+ with peripheral vascular disease+ with osteoarthritis/rheumatoid arthritis+ with osteoporosis+ with osteoporosis+ with leukemias and lymphomas+ with lung cancer+ with colon cancer+ women with breast cancerE		
 + with atrial fibrillation + with peripheral vascular disease + with osteoarthritis/rheumatoid arthritis + with osteoporosis + with leukemias and lymphomas + with lung cancer + with colon cancer + women with breast cancer 	29.3%	34.3%
+ with peripheral vascular diseaseE+ with osteoarthritis/rheumatoid arthritisE+ with osteoporosisE+ with leukemias and lymphomas+ with lung cancer+ with colon cancer+ women with breast cancerE	13.3%	5 17.7%
+ with osteoarthritis/rheumatoid arthritis E + with osteoporosis E + with leukemias and lymphomas E + with lung cancer E + with colon cancer E + women with breast cancer E	11.8%	5 14.4%
+ with osteoporosis E + with leukemias and lymphomas + with lung cancer + with colon cancer + women with breast cancer E	10.2%	5 14.7%
+ with leukemias and lymphomas + with lung cancer + with colon cancer + women with breast cancer E	43.3%	49.1%
+ with lung cancer + with colon cancer + women with breast cancer E	12.5%	5 17.4%
+ with colon cancer + women with breast cancer E	1.7%	2.0%
+ women with breast cancer E	1.2%	5 1.6%
	2.9%	2.4%
+ women with endometrial cancer	5.4%	9.8%
	1.2%	5 1.7%
+ men with prostate cancer	12.9%	5 11.5%
+ with benign prostatic hyperplasia	36.5%	36.8%
+ with HIV/AIDS	0.04%	0.05%
+ with hypothyroidism E	16.8%	5 20.8%
+ with anemia E	00.001	37.3%
+ with chronic kidney disease E	28.6%	5 22.3%
+ with liver diseases E		
+ with fibromyalgia, chronic pain and fatigue	17.9%	6.9%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		4.4%	4.0%
% 65+ with epilepsy		2.0%	2.1%
% 65+ with traumatic brain injury		1.3%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma		23.0%	22.9%
% 65+ with cataract	В	54.4%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.1%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	В	46.8%	54.4%
% 65+ with 0 chronic conditions	В	16.4%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		7.5%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		17.0%	14.4%
% 65+ with self-reported vision difficulty	В	0.0%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	0.9%
% 65+ with self-reported cognition difficulty		5.6%	6.9%
% 65+ with self-reported ambulatory difficulty		12.3%	18.8%
% 65+ with clinical diagnosis of mobility impairments		2.8%	3.2%
% 65+ with self-reported self-care difficulty		2.8%	5.6%
% 65+ with self-reported independent living difficulty	В	2.8%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	4.4%	7.9%
% dually eligible for Medicare and Medicaid	*	4.1%	7.5%
% 60+ with a regular doctor	В	98.2%	95.8%
% 60+ who did not see doctor when needed due to cost		4.2%	5.4%
# of primary care providers within 5 miles		3	2,961
# of hospitals within 5 miles		0	26
# of nursing homes within 5 miles		0	74
# of home health agencies		10	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year		5.9	6.3
Emergency room visits/1000 persons 65+ years per year	*	403	572
Part D monthly prescription fills per person per year	*	44.6	49.1
Home health visits per year	*	1.8	2.5
Durable medical equipment claims per year	*	1.3	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	175	237
Medicare inpatient hospital readmissions (as % of admissions)		13.5%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		57	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports	*	1.3%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		N/A	N/A
% of grandparents raising grandchildren		1.5%	0.8%
% of grandparents who live with grandchildren		3.7%	2.5%
# of assisted living sites		1	134
% of vacant homes in community		0.0%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		97.0%	93.0%
% 60+ who used Internet in last month		80.1%	77.6%
Voter participation rate in 2018 election (age 18+)		67.7%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		115	207
Homicide rate /100,000 persons (county)		1	1
# firearm fatalities (county)		52	586
Property crime rate /100,000 persons		640	2,012
% 65+ who own a motor vehicle		95.5%	91.0%
% 60+ who always drive wearing a seatbelt	В	83.1%	77.1%
# of fatal crashes involving adult age 60+/town		1	151
# of fatal crashes involving adult age 60+/county		17	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		0.0%	5.4%
% 60+ receiving food stamps past year		0.0%	5.7%
% 65+ employed past year		42.8%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		15.3%	18.2%
% households with annual income \$20,000-\$49,999		13.0%	36.5%
% households with annual income > \$50,000		71.8%	45.3%
% 60+ own home		87.2%	79.9%
% 60+ have mortgage on home		53.4%	35.3%
% 65+ households spend >35% of income on housing (renter)		15.3%	8.7%
% 65+ households spend >35% of income on housing (owner)		9.0%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$24,936	\$25,284	0.99
Single, renter, good health	\$26,112	\$26,400	0.99
Couple, homeowner without mortgage, good health	\$36,504	\$37,128	0.98
Couple, renter, good health	\$37,680	\$38,244	0.99

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019