
FOUNDATION

Milan (Coös)

Milan is a rural town in Coös County that is home to Milan Hill State Park. There are 216 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of obesity, depression, anxiety and bipolar disorders, personality disorders, Alzheimer's disease or related dementias, chronic obstructive pulmonary disease, asthma, hypertension, osteoarthritis/rheumatoid arthritis, osteoporosis, benign prostatic hyperplasia, anemia, chronic kidney disease, fibromyalgia/chronic pain/fatigue, migraine, traumatic brain injury, glaucoma, cataracts, pressure ulcers, and visual impairment. However, they had a higher rate of tooth loss. They are less likely to take the health promotion steps of always wearing a seatbelt when driving, engaging in any physical activity within the last month, meeting the CDC guidelines for aerobic and muscle-strengthening physical activities (relative to the state estimate), meeting the CDC preventive health screening goals, or having an annual dental exam. Community resources to support healthy aging include 2 home health agencies and a public library.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		1,396	1,327,503
Population 60 years or older as % of total population		23.2%	22.7%
Total population 60 years or older		324	301,630
Population 65 years or older as % of total population		15.5%	15.8%
Total population 65 years or older		216	210,385
% 65-74 years		72.2%	58.5%
% 75-84 years		21.3%	28.6%
% 85 years or older		6.5%	12.9%
Gender (65+ population)			
% female		42.6%	54.7%
Race/Ethnicity (65+ population)			
% White		100.0%	97.7%
% African American		0.0%	0.5%
% Asian		0.0%	0.9%
% Other		0.0%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		69.0%	58.5%
% divorced/separated		6.9%	14.0%
% widowed		21.3%	22.9%
% never married		2.8%	4.6%
Education (65+ population)			
% with less than high school education		15.7%	12.3%
% with high school or some college		54.6%	57.1%
% with college degree		29.6%	30.6%
% of 65+ population living alone		18.5%	26.1%
% of 65+ population who speak only English at home		69.4%	91.3%
% of 65+ population who are veterans of military service		32.9%	24.8%
Age-sex adjusted 1-year mortality rate	В	2.3%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		1.9%	3.6%
% moved from different county in New Hampshire		1.4%	1.0%
% moved from different state		2.8%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month	W	64.7%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity	W	17.7%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		61.2%	58.6%
% 60+ met CDC guidelines for both types of physical activities	W	12.7%	19.9%
% 60+ getting recommended hours of sleep		64.2%	66.4%
% 60+ injured in a fall within last 12 months		11.3%	10.4%
% 65+ had hip fracture		3.7%	3.3%
% 60+ with self-reported fair or poor health status		24.8%	16.5%
% 60+ with 15+ physically unhealthy days last month		15.2%	12.5%
% 60+ with physical exam/check-up in past year		90.7%	86.5%
% 60+ met CDC preventive health screening goals	W	26.9%	40.3%
% 60+ flu shot past year		63.0%	59.3%
% 65+ with pneumonia vaccine		80.2%	77.8%
% 60+ with cholesterol screening		92.3%	95.3%
% 60+ women with a mammogram within last 2 years		75.9%	79.3%
% 60+ with colorectal cancer screening		69.1%	77.0%
% 60+ with HIV test		9.9%	13.5%
% 60+ current smokers		11.1%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth	W	43.3%	29.0%
% 60+ with annual dental exam	W	60.2%	75.7%
# of dentists per 100,000 persons (all ages) (county)		59	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		13.9%	20.2%
% 60+ self-reported obese		26.2%	27.2%
% 65+ clinically diagnosed obese	В	10.4%	16.7%
% 65+ with high cholesterol		70.1%	72.2%
% 60+ excessive drinking		7.2%	9.2%
% 65+ with poor supermarket access		45.5%	28.4%

% 65+ with post-traumatic stress disorder 1.3% 1.4% % 65+ with schizophrenia & other psychotic disorders 4.5% 4.9% % 65+ with personality disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with substance use disorders 10.6% 10.4% CHRONIC DISEASE 20.57% 28.2% % 65+ with Alzheimer's disease or related dementias B 8.0% 12.0% % 65+ with stroke 11.6% 10.8% 654 with stroke 11.6% 10.8% % 65+ with diabetes 25.7% 28.2% 34.3% 20.5% % 65+ with stroke 11.6% 10.8% 20.5% 654 with stroke 11.6% 10.8% 654 with stroke 13.0% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 365 30.2% 36	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression B 18.3% 28.8% % 65+ with anxiety disorders B 12.5% 21.9% % 65+ with biolar disorders B 1.7% 3.1% % 65+ with schizophrenia & other psychotic disorders 4.5% 4.9% % 65+ with schizophrenia & other psychotic disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with schizophrenia & other psychotic disorders 10.6% 5.5% % 65+ with bactsco use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with tobacco use disorders (drug use +/or alcohol abuse) 5.6% 5.5% % 65+ with tobacco use disorders (drug use +/or alcohol abuse) 5.6% 5.5% % 65+ with bachener's disease or related dementias B 8.0% 12.0% % 65+ with dribetes 25.7% 28.2% 65* % 65+ with hypertension B 16.7% 20.5% % 65+ with hypertension B 65.4% 70.2% % 65+ with chronic obstructive pulmonary disease 32.2% 34.3% % 65+ with ustharia	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders B 12.5% 21.9% % 65+ with bolar disorders B 1.7% 3.1% % 65+ with bolar disorders 4.5% 4.9% % 65+ with bersonality disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with bersonality disorders B 0.5% 1.1% % 65+ with bersonality disorders 10.6% 10.4% CHRONC DISEASE 10.6% 10.2% 28.2% % 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with stroke 11.6% 10.8% % 65+ with stroke 11.6% 10.8% % 65+ with stroma 8 9.4% 10.3% % 65+ with stroma 8 9.4% 10.8% % 65+ with stroma 8 9.4% 10.8% % 65+ with stroma 11.5%	% 60+ with 15+ days poor mental health last month		8.1%	6.9%
% 65+ with bipolar disorders B 1.7% 3.1% % 65+ with post-traumatic stress disorder 1.3% 1.4% % 65+ with personality disorders 4.5% 4.9% % 65+ with personality disorders B 0.5% 1.1% % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with substance use disorders 10.6% 10.4% CHRONIC DISEASE 25.7% 28.2% % 65+ with diabetes 25.7% 28.2% % 65+ with stoke 11.6% 10.8% % 65+ with diabetes 25.7% 28.2% % 65+ with asthma B 9.4% 13.0% % 65+ with diabetes 22.2% 34.3% % 65+ with diabetes 32.2% 34.3% % 65+ with stoke <td< td=""><td>% 65+ with depression</td><td>В</td><td>18.3%</td><td>28.8%</td></td<>	% 65+ with depression	В	18.3%	28.8%
% 65+ with post-traumatic stress disorder 1.3% 1.4% % 65+ with schizophrenia & other psychotic disorders 4.5% 4.9% % 65+ with personality disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with substance use disorders 10.6% 10.4% CHRONIC DISEASE 10.6% 10.4% % 65+ with Alzheimer's disease or related dementias B 8.0% 12.0% % 65+ with stroke 11.6% 10.8% 654 % 65+ with schemic heart disease 32.2% 34.3% 70.2% % 65+ with congestive heart disease 32.2% 34.3% 70.2% % 65+ with congestive heart disease 32.2% 34.3% 70.2% % 65+ with congestive heart disease 32.2% 14.7% 65+ with osteoporosis	% 65+ with anxiety disorders	В	12.5%	21.9%
% 65+ with schizophrenia & other psychotic disorders 4.5% 4.9% % 65+ with personality disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with tobacco use disorders 10.6% 10.4% CHRONIC DISEASE 25.7% 28.2% % 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with stroke 11.6% 10.8% % 65+ with torkonic obstructive pulmonary disease B 16.7% 20.5% % 65+ with stroke 11.6% 10.2% 4.3% % 65+ with astima B 9.4% 13.0% % 65+ with astima B 65.4% 70.2% % 65+ with congestive heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with osteoarthritis/rheumatoid arthritis B 11.2% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 11.2%<	% 65+ with bipolar disorders	В	1.7%	3.1%
% 65+ with personality disorders B 0.5% 1.1% # opioid deaths (all ages) (county) 37 1.279 % 65+ with substance use disorders 10.6% 10.4% CHRONIC DISEASE 10.6% 10.4% % 65+ with dabetes 25.7% 28.2% % 65+ with dabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with asthma B 9.4% 13.0% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with osteoarthritis/rheumatoid arthritis B 11.5% 49.1% % 65+ with osteoarthritis/rheumatoid arthritis	% 65+ with post-traumatic stress disorder		1.3%	1.4%
# opioid deaths (all ages) (county) 37 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with tobacco use disorders 10.6% 10.4% CHRONIC DISEASE	% 65+ with schizophrenia & other psychotic disorders		4.5%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 5.5% % 65+ with tobacco use disorders 10.6% 10.4% CHRONIC DISEASE ************************************	% 65+ with personality disorders	В	0.5%	1.1%
% 65+ with tobacco use disorders 10.6% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 8.0% 12.0% % 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with stroke 11.6% 10.8% % 65+ with stroke 11.6% 20.5% % 65+ with obstructive pulmonary disease B 9.4% 13.0% % 65+ with stroke 30.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with congestive heart disease 32.2% 34.3% % 65+ with obsteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoorosis B 11.2% 17.4% % 65+ with lung cancer 1.5% 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% 3.1% 2.4% % 65+ with lung cancer 1.5% 1.6% 3.8% 3.8% 3.8% <td># opioid deaths (all ages) (county)</td> <td></td> <td>37</td> <td>1,279</td>	# opioid deaths (all ages) (county)		37	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 8.0% 12.0% % 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with chronic obstructive pulmonary disease B 16.7% 20.5% % 65+ with asthma B 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ with congestive heart disease 32.2% 34.3% % 65+ with actial fibrillation 12.4% 14.4% % 65+ with dried fibrillation 12.4% 14.4% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with osteoporosis B 11.2% 1.7% % 65+ with locolon cancer	% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.5%	5.5%
% 65+ with Alzheimer's disease or related dementias B 8.0% 12.0% % 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with chronic obstructive pulmonary disease B 16.7% 20.5% % 65+ with asthma B 9.4% 13.0% % 65+ with asthma B 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ with asthma B 9.4% 13.0% % 65+ with asthma is chemic heart disease 32.2% 34.3% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with paripheral vascular disease 13.2% 14.4% % 65+ with paripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoarthritis/rheumatoid arthritis B 11.2% 17.4% % 65+ with log cancer 1.5% 1.6% 2.0% % 65+ with log cancer 1.5% 1.6% 2.0% % 65+ with loc	% 65+ with tobacco use disorders		10.6%	10.4%
% 65+ with diabetes 25.7% 28.2% % 65+ with stroke 11.6% 10.8% % 65+ with chronic obstructive pulmonary disease B 16.7% 20.5% % 65+ with asthma B 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ ever had a heart attack 4.0% 4.5% % 65+ with congestive heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoarthritis/rheumatoid arthritis B 11.2% 17.4% % 65+ with losteoarthritis/rheumatoid arthritis B	CHRONIC DISEASE			
% 65+ with stroke 11.6% 10.8% % 65+ with chronic obstructive pulmonary disease B 16.7% 20.5% % 65+ with asthma B 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ ever had a heart attack 4.0% 4.5% % 65+ ever had a heart attack 4.0% 4.5% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with loug cancer 1.5% 1.6% 2.0% % 65+ women with prestat cancer 8.8% 9.8% 9.8% % 65+ women with prostate cancer 9.3% 11.5% 1.5% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 22.3% <td>% 65+ with Alzheimer's disease or related dementias</td> <td>В</td> <td>8.0%</td> <td>12.0%</td>	% 65+ with Alzheimer's disease or related dementias	В	8.0%	12.0%
% 65+ with chronic obstructive pulmonary disease B 16.7% 20.5% % 65+ with asthma B 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ ever had a heart attack 4.0% 4.5% % 65+ ever had a heart attack 4.0% 4.5% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with loug cancer 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% % 65+ women with prostate cancer 8.8% 9.8% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with hulping prostatic hyperplasia B 27.1% 36.8% % 65+ with hulping prostatic hyperplasia B 29.1% 37.3% % 65+ with hypothyroidism 19.7% 20.8%<	% 65+ with diabetes		25.7%	28.2%
b 9.4% 13.0% % 65+ with hypertension B 65.4% 70.2% % 65+ ever had a heart attack 4.0% 4.5% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% % 65+ women with prostate cancer 9.3% 11.5% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with heign prostatic hyperplasia B 27.1% 36.8% % 65+ with hypothyroidism 19.7% 20.8% 65+ with anemia B 29.1% 37.3% % 65+ with hypothyroidism 19.7% 20.8% 65+ with chronic kidney disease B 15.5%	% 65+ with stroke		11.6%	10.8%
% 65+ with hypertension B 65.4% 70.2% % 65+ ever had a heart attack 4.0% 4.5% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% % 65+ wornen with prostate cancer 8.8% 9.8% % 65+ wornen with prostate cancer 9.3% 11.5% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease <t< td=""><td>% 65+ with chronic obstructive pulmonary disease</td><td>В</td><td>16.7%</td><td>20.5%</td></t<>	% 65+ with chronic obstructive pulmonary disease	В	16.7%	20.5%
% 65+ ever had a heart attack 4.0% 4.5% % 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% % 65+ with colon cancer 3.1% 2.4% % 65+ women with prostate cancer 8.8% 9.8% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with hypothyroidism 19.7% 20.8% 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% 37.3% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with asthma	В	9.4%	13.0%
% 65+ with ischemic heart disease 32.2% 34.3% % 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoarthritis/rheumatoid arthritis B 11.2% 17.4% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with long cancer 1.6% 2.0% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with hulp/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with hypertension	В	65.4%	70.2%
% 65+ with congestive heart failure 15.2% 17.7% % 65+ with atrial fibrillation 12.4% 14.4% % 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with colon cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 9.3% 11.5% % 65+ with benign prostatic cancer 9.3% 11.5% % 65+ with hypothyroidism 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease	% 65+ ever had a heart attack		4.0%	4.5%
% 65+ with atrial fibrillation 12.4% 14.4% % 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteopartsi B 11.2% 17.4% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with long cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ with benign prostatic cancer 9.3% 11.5% % 65+ with hypothyroidism 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease B 29.1% 37.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with ischemic heart disease		32.2%	34.3%
% 65+ with peripheral vascular disease 13.2% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with lung cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with chronic kidney disease 5.1% 6.9%	% 65+ with congestive heart failure		15.2%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis B 41.5% 49.1% % 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with lung cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% 37.3% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with atrial fibrillation		12.4%	14.4%
% 65+ with osteoporosis B 11.2% 17.4% % 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with lung cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with peripheral vascular disease		13.2%	14.7%
% 65+ with leukemias and lymphomas 1.6% 2.0% % 65+ with lung cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with hypothyroidism 0.05% 0.05% % 65+ with namenia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with osteoarthritis/rheumatoid arthritis	В	41.5%	49.1%
% 65+ with lung cancer 1.5% 1.6% % 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with hypothyroidism 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with osteoporosis	В	11.2%	17.4%
% 65+ with colon cancer 3.1% 2.4% % 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with leukemias and lymphomas		1.6%	2.0%
% 65+ women with breast cancer 8.8% 9.8% % 65+ women with endometrial cancer 1.2% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with lung cancer		1.5%	1.6%
% 65+ women with endometrial cancer 1.2% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with colon cancer		3.1%	2.4%
% 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ women with breast cancer		8.8%	9.8%
% 65+ with benign prostatic hyperplasia B 27.1% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ women with endometrial cancer		1.2%	1.7%
% 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ men with prostate cancer		9.3%	11.5%
% 65+ with hypothyroidism 19.7% 20.8% % 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with benign prostatic hyperplasia	В	27.1%	36.8%
% 65+ with anemia B 29.1% 37.3% % 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with HIV/AIDS		0.05%	0.05%
% 65+ with chronic kidney disease B 15.5% 22.3% % 65+ with liver diseases 5.1% 6.9%	% 65+ with hypothyroidism		19.7%	20.8%
% 65+ with liver diseases 5.1% 6.9%	% 65+ with anemia	В	29.1%	37.3%
	% 65+ with chronic kidney disease	В	15.5%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue B 13.8% 18.6%	-		5.1%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	В	13.8%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache	В	3.0%	4.0%
% 65+ with epilepsy		2.5%	2.1%
% 65+ with traumatic brain injury	В	0.6%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	В	17.7%	22.9%
% 65+ with cataract	В	54.6%	61.2%
% 65+ with pressure ulcer or chronic ulcer	В	3.7%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	В	44.2%	54.4%
% 65+ with 0 chronic conditions		13.1%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty	W	25.0%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		16.1%	14.4%
% 65+ with self-reported vision difficulty		3.7%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment	В	0.4%	0.9%
% 65+ with self-reported cognition difficulty		1.4%	6.9%
% 65+ with self-reported ambulatory difficulty		22.7%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.8%	3.2%
% 65+ with self-reported self-care difficulty		1.9%	5.6%
% 65+ with self-reported independent living difficulty		8.3%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	3.5%	7.9%
% dually eligible for Medicare and Medicaid		6.6%	7.5%
% 60+ with a regular doctor		93.8%	95.8%
% 60+ who did not see doctor when needed due to cost		8.2%	5.4%
# of primary care providers within 5 miles		0	2,961
# of hospitals within 5 miles		0	26
# of nursing homes within 5 miles		0	74
# of home health agencies		2	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	2.1	6.3
Emergency room visits/1000 persons 65+ years per year	*	459	572
Part D monthly prescription fills per person per year		47.6	49.1
Home health visits per year		2.0	2.5
Durable medical equipment claims per year		2.1	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	142	237
Medicare inpatient hospital readmissions (as % of admissions)		11.5%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year	*	46	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports		2.5%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		3	N/A
% of grandparents raising grandchildren		1.8%	0.8%
% of grandparents who live with grandchildren		2.4%	2.5%
# of assisted living sites		0	134
% of vacant homes in community		30.2%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		70.0%	93.0%
% 60+ who used Internet in last month		61.3%	77.6%
Voter participation rate in 2018 election (age 18+)		48.7%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		N/A	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		31	586
Property crime rate /100,000 persons		N/A	2,012
% 65+ who own a motor vehicle		93.9%	91.0%
% 60+ who always drive wearing a seatbelt	W	60.7%	77.1%
# of fatal crashes involving adult age 60+/town		1	151
# of fatal crashes involving adult age 60+/county		11	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		5.1%	5.4%
% 60+ receiving food stamps past year		5.7%	5.7%
% 65+ employed past year		14.8%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		20.9%	18.2%
% households with annual income \$20,000-\$49,999		40.0%	36.5%
% households with annual income > \$50,000		39.1%	45.3%
% 60+ own home		85.9%	79.9%
% 60+ have mortgage on home		26.6%	35.3%
% 65+ households spend >35% of income on housing (renter)		3.5%	8.7%
% 65+ households spend >35% of income on housing (owner)		15.7%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$23,496	\$25,284	0.93
Single, renter, good health	\$23,100	\$26,400	0.88
Couple, homeowner without mortgage, good health	\$35,436	\$37,128	0.95
Couple, renter, good health	\$35,040	\$38,244	0.92

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019