

New London is a town located in Merrimack County and is home to Colby-Sawyer College. There are 1,532 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of obesity, depression, anxiety disorders, post-traumatic stress disorder, schizophrenia and other psychotic disorders, tobacco use disorders, Alzheimer's disease or related dementias, diabetes, asthma, congestive heart failure, lung cancer, chronic kidney disease, and fibromyalgia/chronic pain/fatigue, and mobility impairments. However, they had higher rates of osteoarthritis/rheumatoid arthritis, osteoporosis, leukemias and lymphomas, prostate cancer, benign prostatic hyperplasia, hypothyroidism, glaucoma, and cataracts. Community resources to support healthy aging include 21 primary care providers and a hospital within 5 miles, 2 home health agencies, a senior center, an adult day center, an assisted living site, a university or community college, a public library, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		4,594	1,327,503
Population 60 years or older as % of total population		36.7%	22.7%
Total population 60 years or older		1,688	301,630
Population 65 years or older as % of total population		33.3%	15.8%
Total population 65 years or older		1,532	210,385
% 65-74 years		40.0%	58.5%
% 75-84 years		33.6%	28.6%
% 85 years or older		26.4%	12.9%
Gender (65+ population)			
% female		56.1%	54.7%
Race/Ethnicity (65+ population)			
% White		99.7%	97.7%
% African American		0.0%	0.5%
% Asian		0.0%	0.9%
% Other		0.3%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		58.9%	58.5%
% divorced/separated		12.8%	14.0%
% widowed		25.7%	22.9%
% never married		2.6%	4.6%
Education (65+ population)			
% with less than high school education		4.0%	12.3%
% with high school or some college		33.4%	57.1%
% with college degree		62.7%	30.6%
% of 65+ population living alone		33.3%	26.1%
% of 65+ population who speak only English at home		95.3%	91.3%
% of 65+ population who are veterans of military service		27.3%	24.8%
Age-sex adjusted 1-year mortality rate	В	2.7%	4.1%
Now London (Marrimack)			DACE 1

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		5.1%	3.6%
% moved from different county in New Hampshire		1.1%	1.0%
% moved from different state		0.3%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		71.7%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		26.4%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		59.6%	58.6%
% 60+ met CDC guidelines for both types of physical activities		20.1%	19.9%
% 60+ getting recommended hours of sleep		70.6%	66.4%
% 60+ injured in a fall within last 12 months		11.2%	10.4%
% 65+ had hip fracture		4.0%	3.3%
% 60+ with self-reported fair or poor health status		18.1%	16.5%
% 60+ with 15+ physically unhealthy days last month		13.5%	12.5%
% 60+ with physical exam/check-up in past year		84.8%	86.5%
% 60+ met CDC preventive health screening goals		40.7%	40.3%
% 60+ flu shot past year		60.6%	59.3%
% 65+ with pneumonia vaccine		78.0%	77.8%
% 60+ with cholesterol screening		94.1%	95.3%
% 60+ women with a mammogram within last 2 years		76.7%	79.3%
% 60+ with colorectal cancer screening		73.1%	77.0%
% 60+ with HIV test		13.1%	13.5%
% 60+ current smokers		7.4%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		32.8%	29.0%
% 60+ with annual dental exam		73.2%	75.7%
# of dentists per 100,000 persons (all ages) (county)		78	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		18.2%	20.2%
% 60+ self-reported obese		31.5%	27.2%
% 65+ clinically diagnosed obese	В	8.2%	16.7%
% 65+ with high cholesterol		72.2%	72.2%
% 60+ excessive drinking		8.2%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

% 60+ with 15+ days poor mental health last month 5.6% 6.9% % 65+ with depression B 25.5% 28.8% % 65+ with polar disorders B 17.9% 21.9% % 65+ with post-traumatic stress disorder B 0.7% 1.4% % 65+ with post-traumatic stress disorder B 0.7% 1.4% % 65+ with post-traumatic stress disorders B 2.8% 4.9% % 65+ with post-traumatic stress disorders B 2.8% 4.9% % 65+ with post-traumatic stress disorders B 2.8% 4.9% % 65+ with schizophrenia & other psychotic disorders B 2.8% 4.9% % 65+ with substance use disorders B 2.8% 4.9% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.5% % 65+ with bubacco use disorders (drug use +/or alcohol abuse) 4.7% 5.5% % 65+ with bubacco use disorders (drug use +/or alcohol abuse) 4.7% 5.5% % 65+ with bubacco use disorders B 4.4% 10.4% CHRONIC DISCASE B 4.1% 4.20	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression B 25.5% 28.8% % 65+ with anxiety disorders B 17.9% 21.9% % 65+ with post-traumatic stress disorder B 0.7% 1.4% % 65+ with post-traumatic stress disorder B 0.7% 1.4% % 65+ with post-traumatic stress disorders B 2.8% 4.9% % 65+ with schizophrenia & other psychotic disorders B 2.8% 4.9% % 65+ with post-traumatic stress disorders B 2.8% 4.9% % 65+ with schizophrenia & other psychotic disorders B 2.8% 4.9% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.5% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.5% % 65+ with bubbacco use disorders B 4.4% 10.4% CHRONIC DISEASE B 4.4% 10.4% CHRONIC DISEASE B 16.4% 28.2° % 65+ with Alzheimer's disease or related dementias B 9.8% 12.0° % 65+ with chronic obstructive pulmonary disease B 16.4%<	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders B 17.9% 21.9% % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with personality disorders B 2.8% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1,27 % 65+ with bubstance use disorders (drug use +/or alcohol abuse) 4.7% 5.59 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE B 4.4% 10.49 CHRONIC DISEASE B 16.4% 28.29 65+ with disbetes B 16.4% 28.29 65+ with disbetes B 16.4% 28.29 65+ with bub fronic obstructive pulmonary disease B 16.4% 28.29 65+ with bub fronic obstructive pulmonary disease B 10.7% 13.09 65+ with bub fronic obstructive pulmonary disease B 10.7% 13.09 65+ with hub poperation at a track <th< td=""><td>% 60+ with 15+ days poor mental health last month</td><td></td><td>5.6%</td><td>6.9%</td></th<>	% 60+ with 15+ days poor mental health last month		5.6%	6.9%
% 65+ with bipolar disorders 2.3% 3.1% % 65+ with post-traumatic stress disorder B 0.7% 1.4% % 65+ with post-traumatic stress disorders B 2.8% 4.99 % 65+ with personality disorders D.8% 1.17 # opioid deaths (all ages) (county) 1118 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.59 % 65+ with substance use disorders B 4.4% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with stroke 9.7% 10.89 % 65+ with hypertension 6.8% 70.29 % 65+ with hypertens	% 65+ with depression	В	25.5%	28.8%
% 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 2.8% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.59 % 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.7% 20.59 % 65+ with stroke 9.7% 6.88 70.29 % 65+ with stroke 19.7% 13.09 % 65+ with stroke 9.7% 6.5 % 65+ with stroke 9.7% 6.5 % 65+ with stroke 9.7% 6.5 % 65+ with stroke 9.1% 4.5 % 65+ with stroke 9.1	% 65+ with anxiety disorders	В	17.9%	21.9%
% 65+ with schizophrenia & other psychotic disorders B 2.8% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1,27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.5% 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with disbetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with stroke 9.7% 10.89 % 65+ with stroke with dispersension 68.8% 70.29 % 65+ with stromic obstructive pulmonary disease 18.7% 20.59 % 65+ with stromic heart disease 34.0% 34.39 % 65+ with in thronic obstructive pulmonary disease 34.0% 34.39 % 65+ with hypertension 68.8% 70.29 % 65+ with strain disease 34.0% 34.39 % 65+ with strain fibrillation 16.5% 14.79 % 65+	% 65+ with bipolar disorders		2.3%	3.1%
% 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1,27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.59 % 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with diabetes B 16.4% 28.29 % 65+ with diabetes B 16.4% 28.29 % 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.79 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with pertension 68.8% 70.22 % 65+ with pertension 68.8% 70.22 % 65+ with ischemic heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.3% % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with peripheral vascular disease 13.4% 14.79 % 65	% 65+ with post-traumatic stress disorder	В	0.7%	1.4%
# opioid deaths (all ages) (county) # opioid deaths (all ages) # opioid deaths # opioid deaths (all ages) # opioid deaths	% 65+ with schizophrenia & other psychotic disorders	В	2.8%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 4.7% 5.59 % 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with Alzheimer's disease or related dementias B 16.4% 28.2° % 65+ with Alzheimer's disease or related dementias B 16.4% 28.2° % 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.7% 20.5° % 65+ with asthma B 10.7% 13.0° % 65+ with obstructive pulmonary disease 8 10.7% 13.0° % 65+ with bypertension 68.8% 70.2° % 65+ with hypertension 68.8% 70.2° % 65+ with bypertension 68.8% 70.2° % 65+ with progestive heart disease 34.0% 34.3° % 65+ with congestive heart disease 34.0% 34.3° % 65+ with toral disease 13.4% 14.7° % 65+ with pripheral vascular disease 13.4% 14.7° % 65+ with osteoparthritis/rheumatoid arthritis	% 65+ with personality disorders		0.8%	1.1%
% 65+ with tobacco use disorders B 4.4% 10.49 CHRONIC DISEASE *** *** 12.09 % 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with Alzheimer's disease or related dementias B 16.4% 28.29 % 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with rocobstructive pulmonary disease 18.7% 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with hypertension 68.8% 70.29 % 65+ with hypertension 68.8% 70.29 % 65+ were had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with orgestive heart failure B 13.3% 17.79 % 65+ with osteoporosis W 21.7% 14.49 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with lung cancer B 0.9% 1.69 % 65+ with posteoporosis	# opioid deaths (all ages) (county)		118	1,279
CHRONIC DISEASE 8 65+ with Alzheimer's disease or related dementias B 9.8% 12.09 % 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.7% 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with pypertension 68.8% 70.29 % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.5% % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with loug cancer B 0.9% 1.69 % 65+ with loug cancer B 0.9% 1.69 % 65+ with prostate cancer <t< td=""><td>% 65+ with substance use disorders (drug use +/or alcohol abuse)</td><td></td><td>4.7%</td><td>5.5%</td></t<>	% 65+ with substance use disorders (drug use +/or alcohol abuse)		4.7%	5.5%
% 65+ with Alzheimer's disease or related dementias B 9.8% 12.0% % 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.8% % 65+ with chronic obstructive pulmonary disease 18.7% 20.5% % 65+ with asthma B 10.7% 13.0% % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.5% % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with lukemias and lymphomas W 4.6% 2.09 % 65+ with brit colon cancer B 0.9% 1.6% % 65+ with olon cancer 12.0% 9.8% % 65+ women with prostate cancer W 17.3% 1.5% % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with hy	% 65+ with tobacco use disorders	В	4.4%	10.4%
% 65+ with diabetes B 16.4% 28.29 % 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.7% 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis//rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.4% % 65+ with lung cancer B 0.9% 1.69 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ with prostate cancer W 17.3% 1.59 % 65+ women with prostate cancer W 17.3% 1.59 % 65+ with benign prostatic hyperplasia W	CHRONIC DISEASE			
% 65+ with stroke 9.7% 10.89 % 65+ with chronic obstructive pulmonary disease 18.7% 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with lung cancer B 0.9% 1.69 % 65+ with lung cancer B 0.9% 1.69 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with prostate cancer W 17.3% 11.59 % 65+ women with prostatic hyperplasia W 56.4% 36.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with hypothyroidism W	% 65+ with Alzheimer's disease or related dementias	В	9.8%	12.0%
% 65+ with chronic obstructive pulmonary disease 18.7% 20.59 % 65+ with asthma B 10.7% 13.09 % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with prostate cancer W 17.3% 11.59 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ men with pengap prostatic hyperplasia W 56.4% 36.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with ph	% 65+ with diabetes	В	16.4%	28.2%
% 65+ with asthma B 10.7% 13.09 % 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with leukemias and lymphomas W 21.7% 17.49 % 65+ with lung cancer B 0.9% 1.69 % 65+ with lung cancer B 0.9% 1.69 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ with benign prostatic hyperplasia W 17.3% 11.59 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases <td>% 65+ with stroke</td> <td></td> <td>9.7%</td> <td>10.8%</td>	% 65+ with stroke		9.7%	10.8%
% 65+ with hypertension 68.8% 70.29 % 65+ ever had a heart attack 4.1% 4.59 % 65+ with ischemic heart disease 34.0% 34.33 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with preast cancer 12.0% 9.89 % 65+ women with endometrial cancer 15.5% 1.79 % 65+ mith benign prostatic hyperplasia W 56.4% 36.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with nemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99<	% 65+ with chronic obstructive pulmonary disease		18.7%	20.5%
% 65+ ever had a heart attack 4.1% 4.5% % 65+ with ischemic heart disease 34.0% 34.3% % 65+ with congestive heart failure B 13.3% 17.7% % 65+ with atrial fibrillation 16.5% 14.4% % 65+ with peripheral vascular disease 13.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.1% % 65+ with osteoporosis W 21.7% 17.4% % 65+ with leukemias and lymphomas W 4.6% 2.0% % 65+ with colon cancer B 0.9% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer 12.0% 9.8% % 65+ men with prostate cancer W 17.3% 11.5% % 65+ with benign prostatic hyperplasia W 56.4% 36.8% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with asthma	В	10.7%	13.0%
% 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 12.0% 9.89 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.5%	% 65+ with hypertension		68.8%	70.2%
% 65+ with congestive heart failure B 13.3% 17.79 % 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ ever had a heart attack		4.1%	4.5%
% 65+ with atrial fibrillation 16.5% 14.49 % 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 12.0% 9.89 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with ischemic heart disease		34.0%	34.3%
% 65+ with peripheral vascular disease 13.4% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with liver diseases B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with congestive heart failure	В	13.3%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis W 59.0% 49.19 % 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.059 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with atrial fibrillation		16.5%	14.4%
% 65+ with osteoporosis W 21.7% 17.49 % 65+ with leukemias and lymphomas W 4.6% 2.09 % 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.49 % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with peripheral vascular disease		13.4%	14.7%
% 65+ with leukemias and lymphomas W 4.6% 2.0% % 65+ with lung cancer B 0.9% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer 12.0% 9.8% % 65+ women with endometrial cancer 1.5% 1.7% % 65+ men with prostate cancer W 17.3% 11.5% % 65+ with benign prostatic hyperplasia W 56.4% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with anemia 40.2% 37.3% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with osteoarthritis/rheumatoid arthritis	W	59.0%	49.1%
% 65+ with lung cancer B 0.9% 1.69 % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with osteoporosis	W	21.7%	17.4%
% 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer 12.0% 9.8% % 65+ women with endometrial cancer 1.5% 1.7% % 65+ men with prostate cancer W 17.3% 11.5% % 65+ with benign prostatic hyperplasia W 56.4% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with anemia 40.2% 37.3% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with leukemias and lymphomas	W	4.6%	2.0%
% 65+ women with breast cancer 12.0% 9.89 % 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% 0.059 % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with lung cancer	В	0.9%	1.6%
% 65+ women with endometrial cancer 1.5% 1.79 % 65+ men with prostate cancer W 17.3% 11.59 % 65+ with benign prostatic hyperplasia W 56.4% 36.89 % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with colon cancer		2.3%	2.4%
% 65+ men with prostate cancer W 17.3% 11.5% % 65+ with benign prostatic hyperplasia W 56.4% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with anemia 40.2% 37.3% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ women with breast cancer		12.0%	9.8%
% 65+ with benign prostatic hyperplasia W 56.4% 36.8% % 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.8% % 65+ with anemia 40.2% 37.3% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ women with endometrial cancer		1.5%	1.7%
% 65+ with HIV/AIDS 0.05% 0.05% % 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ men with prostate cancer	W	17.3%	11.5%
% 65+ with hypothyroidism W 24.5% 20.89 % 65+ with anemia 40.2% 37.39 % 65+ with chronic kidney disease B 14.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with benign prostatic hyperplasia	W	56.4%	36.8%
% 65+ with anemia 40.2% 37.3% % 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with HIV/AIDS		0.05%	0.05%
% 65+ with chronic kidney disease B 14.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with hypothyroidism	W	24.5%	20.8%
% 65+ with liver diseases 6.5% 6.9%	% 65+ with anemia		40.2%	37.3%
	% 65+ with chronic kidney disease	В	14.6%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue B 14.4% 18.69	% 65+ with liver diseases		6.5%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	В	14.4%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		4.9%	4.0%
% 65+ with epilepsy		1.5%	2.1%
% 65+ with traumatic brain injury		1.1%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	W	27.4%	22.9%
% 65+ with cataract	W	76.3%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.0%	7.1%
% 65+ with 4+ (out of 15) chronic conditions		52.5%	54.4%
% 65+ with 0 chronic conditions	W	6.1%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		10.6%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		16.0%	14.4%
% 65+ with self-reported vision difficulty		3.8%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.6%	0.9%
% 65+ with self-reported cognition difficulty		6.6%	6.9%
% 65+ with self-reported ambulatory difficulty		12.9%	18.8%
% 65+ with clinical diagnosis of mobility impairments	В	2.1%	3.2%
% 65+ with self-reported self-care difficulty		4.4%	5.6%
% 65+ with self-reported independent living difficulty		9.5%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		8.3%	7.9%
% dually eligible for Medicare and Medicaid	*	2.8%	7.5%
% 60+ with a regular doctor		94.3%	95.8%
% 60+ who did not see doctor when needed due to cost		4.5%	5.4%
# of primary care providers within 5 miles		21	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		0	74
# of home health agencies		2	49
# of community health centers		0	22
# of adult day health centers		1	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	5.1	6.3
Emergency room visits/1000 persons 65+ years per year		577	572
Part D monthly prescription fills per person per year	*	40.7	49.1
Home health visits per year	*	1.8	2.5
Durable medical equipment claims per year	*	1.4	2.0
Inpatient hospital stays/1000 persons 65+ years per year		233	237
Medicare inpatient hospital readmissions (as % of admissions)		14.1%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		85	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports	*	1.7%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		1	44
Air pollution: annual # of unhealthy days for 65+ (county)		0	N/A
% of grandparents raising grandchildren		0.2%	0.8%
% of grandparents who live with grandchildren		1.1%	2.5%
# of assisted living sites		1	134
% of vacant homes in community		25.4%	16.0%
# of universities and community colleges		1	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		92.0%	93.0%
% 60+ who used Internet in last month		76.5%	77.6%
Voter participation rate in 2018 election (age 18+)		64.1%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		28	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		51	586
Property crime rate /100,000 persons		626	2,012
% 65+ who own a motor vehicle		87.4%	91.0%
% 60+ who always drive wearing a seatbelt		72.2%	77.1%
# of fatal crashes involving adult age 60+/town		0	151
# of fatal crashes involving adult age 60+/county		18	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		0.9%	5.4%
% 60+ receiving food stamps past year		3.2%	5.7%
% 65+ employed past year		25.3%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		11.2%	18.2%
% households with annual income \$20,000-\$49,999		30.9%	36.5%
% households with annual income > \$50,000		57.9%	45.3%
% 60+ own home		79.7%	79.9%
% 60+ have mortgage on home		24.4%	35.3%
% 65+ households spend >35% of income on housing (renter)		3.6%	8.7%
% 65+ households spend >35% of income on housing (owner)		25.0%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,860	\$25,284	1.02
Single, renter, good health	\$26,376	\$26,400	1.00
Couple, homeowner without mortgage, good health	\$37,800	\$37,128	1.02
Couple, renter, good health	\$38,316	\$38,244	1.00

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are not due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019