North Hampton (Rockingham)

North Hampton is a seaside town in Rockingham County and home to the Fuller Gardens. There are 915 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of hip fracture, tooth loss, depression, bipolar disorders, schizophrenia and other psychotic disorders, substance use disorders, Alzheimer's disease or related dementias, diabetes, chronic obstructive pulmonary disease, asthma, glaucoma, and hearing impairment. They are more likely to take the health promotion steps of engaging in any physical activity within the past month, meeting the CDC guidelines for muscle-strengthening and aerobic physical activities (relative to the state estimate), and having an annual dental exam. Community resources to support healthy aging include 27 primary care providers and 2 nursing homes within 5 miles, 8 home health agencies, a public library and access to broadband.

POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		4,384	1,327,503
Population 60 years or older as % of total population		28.4%	22.7%
Total population 60 years or older		1,244	301,630
Population 65 years or older as % of total population		20.9%	15.8%
Total population 65 years or older		915	210,385
% 65-74 years		55.8%	58.5%
% 75-84 years		38.1%	28.6%
% 85 years or older		6.0%	12.9%
Gender (65+ population)			
% female		53.0%	54.7%
Race/Ethnicity (65+ population)			
% White		97.9%	97.7%
% African American		0.0%	0.5%
% Asian		2.1%	0.9%
% Other		0.0%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		60.9%	58.5%
% divorced/separated		16.9%	14.0%
% widowed		19.3%	22.9%
% never married		2.8%	4.6%
Education (65+ population)			
% with less than high school education		2.8%	12.3%
% with high school or some college		64.2%	57.1%
% with college degree		33.0%	30.6%
% of 65+ population living alone		17.8%	26.1%
% of 65+ population who speak only English at home		93.6%	91.3%
% of 65+ population who are veterans of military service		25.9%	24.8%
Age-sex adjusted 1-year mortality rate		3.6%	4.1%





HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		0.9%	3.6%
% moved from different county in New Hampshire		2.5%	1.0%
% moved from different state		1.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month	В	84.5%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity	В	37.2%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		60.6%	58.6%
% 60+ met CDC guidelines for both types of physical activities	В	29.6%	19.9%
% 60+ getting recommended hours of sleep		65.1%	66.4%
% 60+ injured in a fall within last 12 months		9.0%	10.4%
% 65+ had hip fracture	В	2.6%	3.3%
% 60+ with self-reported fair or poor health status		12.8%	16.5%
% 60+ with 15+ physically unhealthy days last month		9.6%	12.5%
% 60+ with physical exam/check-up in past year		90.9%	86.5%
% 60+ met CDC preventive health screening goals		45.0%	40.3%
% 60+ flu shot past year		59.3%	59.3%
% 65+ with pneumonia vaccine		79.5%	77.8%
% 60+ with cholesterol screening		96.9%	95.3%
% 60+ women with a mammogram within last 2 years		80.6%	79.3%
% 60+ with colorectal cancer screening		82.4%	77.0%
% 60+ with HIV test		12.6%	13.5%
% 60+ current smokers		6.4%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth	В	20.4%	29.0%
% 60+ with annual dental exam	В	85.1%	75.7%
# of dentists per 100,000 persons (all ages) (county)		72	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		26.9%	20.2%
% 60+ self-reported obese		21.1%	27.2%
% 65+ clinically diagnosed obese		15.7%	16.7%
% 65+ with high cholesterol		71.1%	72.2%
% 60+ excessive drinking		9.5%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

HAVIORAL HEALTH 50+ with 15+ days poor mental health last month 55+ with depression 55+ with anxiety disorders 55+ with post-traumatic stress disorder 55+ with post-traumatic stress disorder 55+ with schizophrenia & other psychotic disorders 55+ with personality disorders pioid deaths (all ages) (county) 55+ with substance use disorders (drug use +/or alcohol abuse) 55+ with tobacco use disorders RONIC DISEASE 55+ with diabetes 55+ with diabetes 55+ with stroke 55+ with hypertension 55+ with hypertension 55+ with congestive heart failure 55+ with atrial fibrillation 55+ with atrial fibrillation 55+ with osteoarthritis/rheumatoid arthritis	B B B B B B B B B B B	5.8% 24.9% 21.8% 1.9% 1.3% 3.8% 0.8% 276 3.8% 9.8% 9.8% 23.9% 9.8% 17.5% 9.1%	
35+ with depression 35+ with anxiety disorders 35+ with bipolar disorders 35+ with post-traumatic stress disorder 35+ with post-traumatic stress disorder 35+ with schizophrenia & other psychotic disorders 35+ with personality disorders pioid deaths (all ages) (county) 35+ with substance use disorders (drug use +/or alcohol abuse) 35+ with tobacco use disorders RONIC DISEASE 35+ with diabetes 35+ with diabetes 35+ with chronic obstructive pulmonary disease 35+ with stroke 35+ with hypertension 35+ with ischemic heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with peripheral vascular disease 35+ with osteoarthritis/rheumatoid arthritis	B B B B B B	24.9% 21.8% 1.9% 1.3% 3.8% 0.8% 276 3.8% 9.8% 9.8% 23.9% 9.8% 17.5%	28.8% 21.9% 3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4% 28.2% 10.8% 20.5%
55+ with anxiety disorders 55+ with bipolar disorders 55+ with post-traumatic stress disorder 55+ with schizophrenia & other psychotic disorders 55+ with personality disorders pioid deaths (all ages) (county) 55+ with substance use disorders (drug use +/or alcohol abuse) 55+ with tobacco use disorders RONIC DISEASE 55+ with diabetes 55+ with hypertension 55+ with hypertension 55+ with hypertension 55+ with ischemic heart disease 55+ with congestive heart failure 55+ with atrial fibrillation 55+ with osteoarthritis/rheumatoid arthritis	B B B B B B	21.8% 1.9% 1.3% 3.8% 0.8% 276 3.8% 9.8% 8.0% 23.9% 9.8% 17.5%	21.9% 3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4% 28.2% 10.8% 20.5%
35+ with bipolar disorders 35+ with post-traumatic stress disorder 35+ with schizophrenia & other psychotic disorders 35+ with personality disorders pioid deaths (all ages) (county) 35+ with substance use disorders (drug use +/or alcohol abuse) 35+ with tobacco use disorders RONIC DISEASE 35+ with diabetes 35+ with stroke 35+ with chronic obstructive pulmonary disease 35+ with asthma 35+ with ischemic heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with osteoarthritis/rheumatoid arthritis	B B B B B	1.9% 1.3% 3.8% 0.8% 276 3.8% 9.8% 23.9% 9.8% 17.5%	3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4% 12.0% 28.2% 10.8% 20.5%
55+ with post-traumatic stress disorder 55+ with schizophrenia & other psychotic disorders 55+ with personality disorders pioid deaths (all ages) (county) 55+ with substance use disorders (drug use +/or alcohol abuse) 55+ with substance use disorders RONIC DISEASE 55+ with Alzheimer's disease or related dementias 55+ with diabetes 55+ with stroke 55+ with chronic obstructive pulmonary disease 55+ with hypertension 55+ ever had a heart attack 55+ with ischemic heart disease 55+ with atrial fibrillation 55+ with osteoarthritis/rheumatoid arthritis	B B B B B	1.3% 3.8% 0.8% 276 3.8% 9.8% 23.9% 9.8% 17.5%	1.4% 4.9% 1.1% 1,279 5.5% 10.4% 28.2% 10.8% 20.5%
35+ with schizophrenia & other psychotic disorders 35+ with personality disorders pioid deaths (all ages) (county) 35+ with substance use disorders (drug use +/or alcohol abuse) 35+ with substance use disorders RONIC DISEASE 35+ with Alzheimer's disease or related dementias 35+ with diabetes 35+ with stroke 35+ with stroke 35+ with asthma 35+ with hypertension 35+ with ischemic heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with peripheral vascular disease 35+ with osteoarthritis/rheumatoid arthritis	B B B B	3.8% 0.8% 276 3.8% 9.8% 8.0% 23.9% 9.8% 17.5%	4.9% 1.1% 1,279 5.5% 10.4% 12.0% 28.2% 10.8% 20.5%
35+ with personality disorders pioid deaths (all ages) (county) 35+ with substance use disorders (drug use +/or alcohol abuse) 35+ with tobacco use disorders RONIC DISEASE 35+ with Alzheimer's disease or related dementias 35+ with diabetes 35+ with stroke 35+ with chronic obstructive pulmonary disease 35+ with asthma 35+ with asthma 35+ with ischemic heart disease 35+ with ischemic heart disease 35+ with asthma 35+ with asthma 35+ with asthma 35+ with asthma 35+ with ongestive heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with peripheral vascular disease 35+ with osteoarthritis/rheumatoid arthritis	B B B B	0.8% 276 3.8% 9.8% 8.0% 23.9% 9.8% 17.5%	1.1% 1,279 5.5% 10.4% 12.0% 28.2% 10.8% 20.5%
pioid deaths (all ages) (county) 35+ with substance use disorders (drug use +/or alcohol abuse) 35+ with tobacco use disorders RONIC DISEASE 35+ with Alzheimer's disease or related dementias 35+ with diabetes 35+ with diabetes 35+ with stroke 35+ with stroke 35+ with chronic obstructive pulmonary disease 35+ with asthma 35+ with asthma 35+ with hypertension 35+ ever had a heart attack 35+ with ischemic heart disease 35+ with ischemic heart disease 35+ with atrial fibrillation 35+ with peripheral vascular disease 35+ with osteoarthritis/rheumatoid arthritis	B B B	276 3.8% 9.8% 8.0% 23.9% 9.8% 17.5%	1,279 5.5% 10.4% 12.0% 28.2% 10.8% 20.5%
65+ with substance use disorders (drug use +/or alcohol abuse) 65+ with tobacco use disorders RONIC DISEASE 65+ with Alzheimer's disease or related dementias 65+ with Alzheimer's disease or related dementias 65+ with diabetes 65+ with stroke 65+ with chronic obstructive pulmonary disease 65+ with asthma 65+ with hypertension 65+ with ischemic heart disease 65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis	B B B	3.8% 9.8% 8.0% 23.9% 9.8% 17.5%	5.5% 10.4% 12.0% 28.2% 10.8% 20.5%
35+ with tobacco use disorders RONIC DISEASE 35+ with Alzheimer's disease or related dementias 35+ with diabetes 35+ with diabetes 35+ with stroke 35+ with chronic obstructive pulmonary disease 35+ with asthma 35+ with hypertension 35+ ever had a heart attack 35+ with ischemic heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with osteoarthritis/rheumatoid arthritis	B B B	9.8% 8.0% 23.9% 9.8% 17.5%	10.4% 12.0% 28.2% 10.8% 20.5%
RONIC DISEASE 65+ with Alzheimer's disease or related dementias 65+ with diabetes 65+ with stroke 65+ with chronic obstructive pulmonary disease 65+ with asthma 65+ with hypertension 65+ ever had a heart attack 65+ with ischemic heart disease 65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis	B	8.0% 23.9% 9.8% 17.5%	12.0% 28.2% 10.8% 20.5%
35+ with Alzheimer's disease or related dementias35+ with diabetes35+ with stroke35+ with chronic obstructive pulmonary disease35+ with asthma35+ with asthma35+ with hypertension35+ ever had a heart attack35+ with ischemic heart disease35+ with congestive heart failure35+ with atrial fibrillation35+ with peripheral vascular disease35+ with osteoarthritis/rheumatoid arthritis	B	23.9% 9.8% 17.5%	28.2% 10.8% 20.5%
35+ with diabetes35+ with stroke35+ with chronic obstructive pulmonary disease35+ with asthma35+ with hypertension35+ ever had a heart attack35+ with ischemic heart disease35+ with congestive heart failure35+ with atrial fibrillation35+ with peripheral vascular disease35+ with osteoarthritis/rheumatoid arthritis	B	23.9% 9.8% 17.5%	28.2% 10.8% 20.5%
65+ with stroke65+ with chronic obstructive pulmonary disease65+ with asthma65+ with hypertension65+ ever had a heart attack65+ with ischemic heart disease65+ with congestive heart failure65+ with atrial fibrillation65+ with peripheral vascular disease65+ with osteoarthritis/rheumatoid arthritis	В	9.8% 17.5%	10.8% 20.5%
 35+ with chronic obstructive pulmonary disease 35+ with asthma 35+ with hypertension 35+ ever had a heart attack 35+ with ischemic heart disease 35+ with congestive heart failure 35+ with atrial fibrillation 35+ with peripheral vascular disease 35+ with osteoarthritis/rheumatoid arthritis 		17.5%	20.5%
65+ with asthma 65+ with hypertension 65+ ever had a heart attack 65+ with ischemic heart disease 65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis			
65+ with hypertension65+ ever had a heart attack65+ with ischemic heart disease65+ with congestive heart failure65+ with atrial fibrillation65+ with peripheral vascular disease65+ with osteoarthritis/rheumatoid arthritis	В	9.1%	12 00/
 65+ ever had a heart attack 65+ with ischemic heart disease 65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis 			13.0%
65+ with ischemic heart disease 65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis		70.3%	70.2%
65+ with congestive heart failure 65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis		3.4%	4.5%
65+ with atrial fibrillation 65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis		35.5%	34.3%
65+ with peripheral vascular disease 65+ with osteoarthritis/rheumatoid arthritis		16.7%	17.7%
5+ with osteoarthritis/rheumatoid arthritis		13.9%	14.4%
		13.9%	14.7%
i5+ with osteoporosis		46.2%	49.1%
		16.5%	17.4%
65+ with leukemias and lymphomas		1.9%	2.0%
65+ with lung cancer		1.4%	1.6%
65+ with colon cancer		2.2%	2.4%
65+ women with breast cancer		10.1%	9.8%
65+ women with endometrial cancer		1.6%	1.7%
65+ men with prostate cancer		12.2%	11.5%
65+ with benign prostatic hyperplasia		35.5%	36.8%
65+ with HIV/AIDS		0.04%	0.05%
65+ with hypothyroidism		20.0%	20.8%
55+ with anemia		35.5%	37.3%
5+ with chronic kidney disease		19.5%	22.3%
5+ with liver diseases		6.9%	6.9%
5+ with fibromyalgia, chronic pain and fatigue		19.8%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		3.9%	4.0%
% 65+ with epilepsy		2.0%	2.1%
% 65+ with traumatic brain injury		0.8%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	В	19.1%	22.9%
% 65+ with cataract		60.8%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.8%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	В	48.4%	54.4%
% 65+ with 0 chronic conditions		9.8%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		15.2%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment	В	11.9%	14.4%
% 65+ with self-reported vision difficulty		3.6%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.5%	0.9%
% 65+ with self-reported cognition difficulty		7.9%	6.9%
% 65+ with self-reported ambulatory difficulty		18.9%	18.8%
% 65+ with clinical diagnosis of mobility impairments		2.8%	3.2%
% 65+ with self-reported self-care difficulty		3.0%	5.6%
% 65+ with self-reported independent living difficulty		9.9%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	4.6%	7.9%
% dually eligible for Medicare and Medicaid	*	2.9%	7.5%
% 60+ with a regular doctor		97.4%	95.8%
% 60+ who did not see doctor when needed due to cost		2.6%	5.4%
# of primary care providers within 5 miles		27	2,961
# of hospitals within 5 miles		0	26
# of nursing homes within 5 miles		2	74
# of home health agencies		8	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	7.5	6.3
Emergency room visits/1000 persons 65+ years per year	*	491	572
Part D monthly prescription fills per person per year	*	43.9	49.1
Home health visits per year		2.3	2.5
Durable medical equipment claims per year		1.7	2.0
Inpatient hospital stays/1000 persons 65+ years per year		203	237
Medicare inpatient hospital readmissions (as % of admissions)		14.1%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		59	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports	*	1.2%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		2	N/A
% of grandparents raising grandchildren		0.0%	0.8%
% of grandparents who live with grandchildren		0.0%	2.5%
# of assisted living sites		0	134
% of vacant homes in community		10.5%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		99.0%	93.0%
% 60+ who used Internet in last month		86.3%	77.6%
Voter participation rate in 2018 election (age 18+)		73.3%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		38	207
Homicide rate /100,000 persons (county)		1	1
# firearm fatalities (county)		92	586
Property crime rate /100,000 persons		986	2,012
% 65+ who own a motor vehicle		94.5%	91.0%
% 60+ who always drive wearing a seatbelt		83.7%	77.1%
# of fatal crashes involving adult age 60+/town		1	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		2.7%	5.4%
% 60+ receiving food stamps past year		5.6%	5.7%
% 65+ employed past year		30.7%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		7.5%	18.2%
% households with annual income \$20,000-\$49,999		24.5%	36.5%
% households with annual income > \$50,000		68.0%	45.3%
% 60+ own home		95.6%	79.9%
% 60+ have mortgage on home		39.0%	35.3%
% 65+ households spend >35% of income on housing (renter)		4.0%	8.7%
% 65+ households spend >35% of income on housing (owner)		14.8%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,536	\$25,284	1.01
Single, renter, good health	\$26,712	\$26,400	1.01
Couple, homeowner without mortgage, good health	\$37,092	\$37,128	1.00
Couple, renter, good health	\$38,268	\$38,244	1.00

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019