Northfield (Merrimack)

Northfield is a town located in Merrimack County that is home to the Tilton Memorial Arch. There are 582 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of osteoarthritis/rheumatoid arthritis, osteoporosis, hypothyroidism, glaucoma, and cataracts. However, they had higher rates of depression, tobacco use disorders, diabetes, chronic obstructive pulmonary disease, hypertension, ischemic heart disease, congestive heart failure, and peripheral vascular disease. They are less likely to take the health promotion steps of getting screened for colorectal cancer and always wearing a seatbelt while driving. Community resources to support healthy aging include 10 primary care providers, a hospital, and a nursing home within 5 miles, 4 home health agencies, 2 assisted living sites, a public library, and access to broadband.

POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		4,821	1,327,503
Population 60 years or older as % of total population		19.8%	22.7%
Total population 60 years or older		953	301,630
Population 65 years or older as % of total population		12.1%	15.8%
Total population 65 years or older		582	210,385
% 65-74 years		63.7%	58.5%
% 75-84 years		25.6%	28.6%
% 85 years or older		10.7%	12.9%
Gender (65+ population)			
% female		47.1%	54.7%
Race/Ethnicity (65+ population)			
% White		100.0%	97.7%
% African American		0.0%	0.5%
% Asian		0.0%	0.9%
% Other		0.0%	0.9%
% Hispanic/Latino		0.0%	0.9%
Marital Status (65+ population)			
% married		64.8%	58.5%
% divorced/separated		20.6%	14.0%
% widowed		13.2%	22.9%
% never married		1.4%	4.6%
Education (65+ population)			
% with less than high school education		24.2%	12.3%
% with high school or some college		54.3%	57.1%
% with college degree		21.5%	30.6%
% of 65+ population living alone		9.6%	26.1%
% of 65+ population who speak only English at home		100.0%	91.3%
% of 65+ population who are veterans of military service		28.7%	24.8%
Age-sex adjusted 1-year mortality rate		4.3%	4.1%





HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		1.2%	3.6%
% moved from different county in New Hampshire		0.0%	1.0%
% moved from different state		0.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		71.7%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		21.8%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		56.3%	58.6%
% 60+ met CDC guidelines for both types of physical activities		16.7%	19.9%
% 60+ getting recommended hours of sleep		58.8%	66.4%
% 60+ injured in a fall within last 12 months		9.6%	10.4%
% 65+ had hip fracture		3.8%	3.3%
% 60+ with self-reported fair or poor health status		17.8%	16.5%
% 60+ with 15+ physically unhealthy days last month		12.4%	12.5%
% 60+ with physical exam/check-up in past year		90.8%	86.5%
% 60+ met CDC preventive health screening goals		38.0%	40.3%
% 60+ flu shot past year		58.8%	59.3%
% 65+ with pneumonia vaccine		77.1%	77.8%
% 60+ with cholesterol screening		93.2%	95.3%
% 60+ women with a mammogram within last 2 years		76.5%	79.3%
% 60+ with colorectal cancer screening	W	69.3%	77.0%
% 60+ with HIV test		9.2%	13.5%
% 60+ current smokers		11.5%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		31.2%	29.0%
% 60+ with annual dental exam		68.9%	75.7%
# of dentists per 100,000 persons (all ages) (county)		78	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		16.8%	20.2%
% 60+ self-reported obese		27.5%	27.2%
% 65+ clinically diagnosed obese		14.8%	16.7%
% 65+ with high cholesterol		74.0%	72.2%
% 60+ excessive drinking		11.0%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

% 60+ with 15+ days poor mental health last month 6.9% 6.9% 6.9% % 65+ with depression W 31.8% 28.89 % 65+ with biolar disorders 3.4% 3.19 % 65+ with post-traumatic stress disorder 1.5% 1.44 % 65+ with post-traumatic stress disorders 5.8% 4.99 % 65+ with post-traumatic stress disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with busbtance use disorders (drug use +/or alcohol abuse) 6.8% 5.55 % 65+ with bubatcace use disorders (drug use +/or alcohol abuse) 6.8% 5.55 % 65+ with abubtance use disorders W 13.4% 10.49 CHEONIC DISEASE V 22.7% 28.27 % 65+ with diabetes W 32.7% 28.27 % 65+ with abtrane 12.1% 10.89 % 65+ with abtrane 12.8% 13.09 % 65+ with stroke 41.7% 14.2% % 65+ with stroke 14.3% 14.4% % 65+ with stroke 12.8% 13.4%	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression W 31.8% 28.8% % 65+ with anxiety disorders 20.6% 21.9% % 65+ with polar disorders 3.4% 3.19 % 65+ with polar disorders 3.4% 3.19 % 65+ with schizophrenia & other psychotic disorders 5.8% 4.99 % 65+ with schizophrenia & other psychotic disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with botstance use disorders (drug use +/or alcohol abuse) 6.8% 5.5% % 65+ with botstance use disorders W 13.4% 10.4% CHRONC DISEASE W 32.7% 28.27 % 65+ with disbetes W 32.7% 28.27 % 65+ with asthma 12.8% 13.00 % 65+ with hopertension W 70.29 % 65+ with hopertension W 74.2% 70.29 % 65+ with asthma 12.8% 13.00 % 65+ with congestive heart failure W 22.0% 17.7% % 65+ with setomaritis//heumatoid arthritis B 44.2% 49.19 <td>BEHAVIORAL HEALTH</td> <td></td> <td></td> <td></td>	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders 20.6% 21.99 % 65+ with bipolar disorders 3.4% 3.1% % 65+ with post-traumatic stress disorder 1.5% 1.49 % 65+ with schizophrenia & other psychotic disorders 5.8% 4.99 % 65+ with bersonality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with bersonality disorders W 13.4% 10.49 CHRONC DISEASE W 32.7% 28.29 % 65+ with diabetes W 32.7% 28.29 % 65+ with strike 12.1% 10.89 % 65+ with strike 12.4% 13.09 % 65+ with strike 12.1% 10.89 % 65+ with strike 12.8% 13.10 % 65+ with strike 12.8% 13.09 % 65+ with strike 12.8% 13.9% % 65+ with strike 12.8%	% 60+ with 15+ days poor mental health last month		6.9%	6.9%
% 65+ with bipolar disorders 3.4% 3.19 % 65+ with post-traumatic stress disorder 1.5% 1.49 % 65+ with personality disorders 5.8% 4.99 % 65+ with personality disorders 0.8% 1.17 % 65+ with personality disorders 0.8% 1.17 % 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with substance use disorders W 13.4% 10.49 CHEONIC DISEASE V 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with otroke 12.1% 10.88 % 65+ with stroke 12.1% 10.88 % 65+ with stroke 12.1% 10.88 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 12.8% 13.09 % 65+ with schernic heart disease W 39.4% 34.39 % 65+ with schernic heart disease W 39.4% 34.39 % 65+ with schernic heart disease W 39.4% 34.39 % 65+ with schernic heart dis	% 65+ with depression	W	31.8%	28.8%
% 65+ with post-traumatic stress disorder 1.5% 1.49 % 65+ with schizophrenia & other psychotic disorders 5.8% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with tobacco use disorders W 13.4% 10.49 CHRONIC DISEASE W 32.7% 28.29 % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 5.6% 4.59 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 5.6% 4.59 % 65+ with scheart disease<	% 65+ with anxiety disorders		20.6%	21.9%
% 65+ with schizophrenia & other psychotic disorders 5.8% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 1118 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with tobacco use disorders W 13.4% 10.49 CHRONC DISEASE	% 65+ with bipolar disorders		3.4%	3.1%
% 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 118 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with tobacco use disorders W 13.4% 10.44 CHRONIC DISEASE V 13.4% 12.4% % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.1% 10.89 % 65+ with stroke 12.8% 13.09 % 65+ with stroke 12.8% 13.09 % 65+ with asthma 12.8% 13.09 % 65+ with operstore heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with congestive heart failure W 28.2% 49.19 % 65+ with congestive heart failure W 28.0% 2.0%	% 65+ with post-traumatic stress disorder		1.5%	1.4%
# opioid deaths (all ages) (county) 118 1,27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with tobacco use disorders W 13.4% 10.49 CHRONIC DISEASE 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.88 % 65+ with stroke 12.1% 10.88 % 65+ with stroke 12.8% 13.09 % 65+ with asthma 12.8% 13.09 % 65+ with operstree heart disease W 34.43 % 65+ with colemative heart disease W 22.0% 17.7% % 65+ with osteopartoxitis/rheumatoid arthritis B 44.2% 49.1% % 65+ with osteopartoxitis/rheumatoid arthritis B 42.2% 20.0% % 65+ with colon cancer 2.7	% 65+ with schizophrenia & other psychotic disorders		5.8%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 6.8% 5.59 % 65+ with tobacco use disorders W 13.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with droke 12.1% 10.89 % 65+ with stroke 12.1% 10.89 % 65+ with obstructive pulmonary disease W 24.5% 20.5% % 65+ with asthma 12.8% 13.0% 65.4 13.0% % 65+ with asthma 12.8% 13.0% 70.29 70.29 70.29 70.29 70.29 70.29 70.29 70.29 70.29 70.45 43.0% 70.29 70.45 43.0% 70.29 70.45 43.0% 70.29 70.43 70.29 70.44 70.29 70.44 70.29 70.44 70.29 70.49 70.49 70.44 70.29 71.7% 7.49 70.29 71.49 76.54 41.7% 41.47 </td <td>% 65+ with personality disorders</td> <td></td> <td>0.8%</td> <td>1.1%</td>	% 65+ with personality disorders		0.8%	1.1%
% 65+ with tobacco use disorders W 13.4% 10.49 CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.8% % 65+ with stroke 12.1% 10.8% % 65+ with obstructive pulmonary disease W 24.5% 20.5% % 65+ with hypertension W 74.2% 70.29 % 65+ with hypertension W 39.4% 34.39 % 65+ with congestive heart disease W 39.4% 34.39 % 65+ with congestive heart disease W 39.4% 34.39 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoorosis B 12.9% 17.4% % 65+ with loukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with loug cancer 1.7% 1.6% 34.0% 36.8% % 65+ with loug cancer 2.1% 1.7% 4.8% 34.0% 36.8%	# opioid deaths (all ages) (county)		118	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.8% % 65+ with stroke 12.1% 10.8% % 65+ with asthma 12.8% 20.5% % 65+ with asthma 12.8% 13.0% % 65+ with chronic heart disease W 74.2% 70.2% % 65+ with actical fibrillation W 22.0% 17.7% % 65+ with astoporosis B 12.9% 14.4% % 65+ with osteoporosis B 12.9% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with locion cancer 2.7% 2.4% 9.8% % 65+ with locion cancer	% 65+ with substance use disorders (drug use +/or alcohol abuse)		6.8%	5.5%
% 65+ with Alzheimer's disease or related dementias 12.4% 12.0% % 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.89 % 65+ with chronic obstructive pulmonary disease W 24.5% 20.59 % 65+ with asthma 12.8% 13.09 % 65+ with schemic heart disease W 70.29 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with osteoparthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with colon cancer 2.1% 1.7% 1.6% % 65+ with loug cancer 1.7% 2.4% 9.1% % 65+ with nelondometrial cancer 2.1% <td< td=""><td>% 65+ with tobacco use disorders</td><td>W</td><td>13.4%</td><td>10.4%</td></td<>	% 65+ with tobacco use disorders	W	13.4%	10.4%
% 65+ with diabetes W 32.7% 28.29 % 65+ with stroke 12.1% 10.89 % 65+ with chronic obstructive pulmonary disease W 24.5% 20.59 % 65+ with asthma 12.8% 13.09 % 65+ with hypertension W 74.2% 70.29 % 65+ with hypertension W 74.2% 70.29 % 65+ with congestive heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with osteoparthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with colon cancer 2.7% 2.4% 39.8% % 65+ with lung cancer 1.7% 1.69 36.89 % 65+ women with prostate cancer 9.3% 11.5% 34.0% 36.89 % 65+ women with prostate cancer 9.3% 11.5% 34.0% 36.89 % 65+ with hologn prostatic hyperplasia 34.0%	CHRONIC DISEASE			
% 65+ with stroke 12.1% 10.89 % 65+ with chronic obstructive pulmonary disease W 24.5% 20.59 % 65+ with asthma 12.8% 13.09 % 65+ with hypertension W 74.2% 70.29 % 65+ ever had a heart attack 5.6% 4.59 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ women with prostate cancer 7.8% 9.89 9.89 % 65+ women with prostate cancer 9.3% 11.5% 5.6% % 65+ with hypothyroidism B 17.2% 20.89	% 65+ with Alzheimer's disease or related dementias		12.4%	12.0%
% 65+ with chronic obstructive pulmonary disease W 24.5% 20.59 % 65+ with asthma 12.8% 13.09 % 65+ with hypertension W 74.2% 70.29 % 65+ ever had a heart attack 5.6% 4.59 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.0% 2.09 % 65+ woren with perister cancer 7.8% 9.89 9.89 65+ woren with prostate cancer 2.1% 1.7% % 65+ woren with prostate cancer 9.3% 11.59 34.0% 36.89 % 65+ woren with prostate cancer 9.3% 11.59 36.89 % 65+ woren with prostate cancer 9.3% 11.59 36.89 % 65+ with hulpong prostatic hyperplasia 34.0% 3	% 65+ with diabetes	W	32.7%	28.2%
% 65+ with asthma 12.8% 13.09 % 65+ with hypertension W 74.2% 70.29 % 65+ with hypertension S.6% 4.59 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with peripheral vascular disease W 18.0% 14.79 % 65+ with osteoorthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with colon cancer 2.7% 2.49 9.89 % 65+ women with prostate cancer 7.8% 9.89 9.3% 11.59 % 65+ with henign prostatic hyperplasia 34.0% 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 36.89 <td< td=""><td>% 65+ with stroke</td><td></td><td>12.1%</td><td>10.8%</td></td<>	% 65+ with stroke		12.1%	10.8%
% 65+ with hypertension W 74.2% 70.29 % 65+ ever had a heart attack 5.6% 4.59 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with peripheral vascular disease W 18.0% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with colon cancer 2.0% 2.0% 2.0% % 65+ with lung cancer 1.7% 1.69 38.89 % 65+ women with breast cancer 7.8% 9.89 % 65+ women with prostate cancer 9.3% 11.59 % 65+ with hypothyroidism B 17.2% 20.89	% 65+ with chronic obstructive pulmonary disease	W	24.5%	20.5%
% 65+ ever had a heart attack 5.6% 4.59 % 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with ischemic heart disease W 22.0% 17.79 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with congestive heart failure W 22.0% 17.79 % 65+ with atrial fibrillation 14.3% 14.49 % 65+ with peripheral vascular disease W 18.0% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.0% 2.09 % 65+ with long cancer 1.7% 1.69 2.0% 2.09 % 65+ women with breast cancer 7.8% 9.89 9.89 % 65+ women with prostate cancer 9.3% 11.59 65+ % 65+ with HIV/AIDS 0.04% 0.059 0.04% 0.059 % 65+ with hypothyroidism B 17.2% 20.89 0.64 0.059 0.04% 0.059 0.04% 0.059 0.65+ 0	% 65+ with asthma		12.8%	13.0%
% 65+ with ischemic heart disease W 39.4% 34.39 % 65+ with congestive heart failure W 22.0% 17.7% % 65+ with atrial fibrillation 14.3% 14.4% % 65+ with peripheral vascular disease W 18.0% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.1% % 65+ with osteoarthritis/rheumatoid arthritis B 12.9% 17.4% % 65+ with osteoporosis B 12.9% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with long cancer 1.7% 1.6% 4.4% 49.1% % 65+ women with breast cancer 2.7% 2.4% 4.4% 49.1% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.6% 4.6% 4.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.0% 2.1% 1.7% 1.6% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4% 4.4%	% 65+ with hypertension	W	74.2%	70.2%
% 65+ with congestive heart failure W 22.0% 17.7% % 65+ with atrial fibrillation 14.3% 14.4% % 65+ with peripheral vascular disease W 18.0% 14.7% % 65+ with osteoparthritis/rheumatoid arthritis B 44.2% 49.1% % 65+ with osteoparthritis/rheumatoid arthritis B 12.9% 17.4% % 65+ with osteoparthritis/rheumatoid arthritis B 12.9% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% 2.0% % 65+ with long cancer 1.7% 1.69 2.6% 2.4% % 65+ with colon cancer 2.7% 2.4% 9.8% % 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with chronic kidney disease	% 65+ ever had a heart attack		5.6%	4.5%
% 65+ with atrial fibrillation 14.3% 14.4% % 65+ with peripheral vascular disease W 18.0% 14.7% % 65+ with peripheral vascular disease W 18.0% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.1% % 65+ with osteoporosis B 12.9% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with colon cancer 1.7% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ with benign prostate cancer 9.3% 11.5% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with chronic kidney disease 21.5%	% 65+ with ischemic heart disease	W	39.4%	34.3%
% 65+ with peripheral vascular disease W 18.0% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.09 % 65+ with leukemias and lymphomas 2.0% 2.09 % 65+ with lung cancer 1.7% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 7.8% 9.89 % 65+ women with endometrial cancer 2.1% 1.79 % 65+ women with prostate cancer 9.3% 11.59 % 65+ with benign prostatic hyperplasia 34.0% 36.89 % 65+ with hypothyroidism B 17.2% 20.89 % 65+ with hypothyroidism B 17.2% 20.89 % 65+ with chronic kidney disease 21.5% 22.39 37.39 % 65+ with liver diseases 8.1% 6.99	% 65+ with congestive heart failure	W	22.0%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis B 44.2% 49.19 % 65+ with osteoporosis B 12.9% 17.4% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with leukemias and lymphomas 2.0% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ women with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with atrial fibrillation		14.3%	14.4%
% 65+ with osteoporosis B 12.9% 17.49 % 65+ with leukemias and lymphomas 2.0% 2.09 % 65+ with lung cancer 1.7% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 2.7% 2.49 % 65+ women with breast cancer 2.7% 2.49 % 65+ women with endometrial cancer 2.1% 1.79 % 65+ men with prostate cancer 9.3% 11.59 % 65+ with benign prostatic hyperplasia 34.0% 36.89 % 65+ with hypothyroidism B 17.2% 20.89 % 65+ with nemia 37.8% 37.39 % 65+ with chronic kidney disease 21.5% 22.39 % 65+ with liver diseases 8.1% 6.99	% 65+ with peripheral vascular disease	W	18.0%	14.7%
% 65+ with leukemias and lymphomas 2.0% 2.09 % 65+ with lung cancer 1.7% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 7.8% 9.89 % 65+ women with endometrial cancer 2.1% 1.79 % 65+ women with prostate cancer 9.3% 11.59 % 65+ with benign prostatic hyperplasia 34.0% 36.89 % 65+ with hypothyroidism B 17.2% 20.89 % 65+ with nemia 37.8% 37.39 % 65+ with chronic kidney disease 21.5% 22.39 % 65+ with liver diseases 8.1% 6.99	% 65+ with osteoarthritis/rheumatoid arthritis	В	44.2%	49.1%
% 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with osteoporosis	В	12.9%	17.4%
% 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with leukemias and lymphomas		2.0%	2.0%
% 65+ women with breast cancer 7.8% 9.8% % 65+ women with endometrial cancer 2.1% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with lung cancer		1.7%	1.6%
% 65+ women with endometrial cancer 2.1% 1.7% % 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with colon cancer		2.7%	2.4%
% 65+ men with prostate cancer 9.3% 11.5% % 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ women with breast cancer		7.8%	9.8%
% 65+ with benign prostatic hyperplasia 34.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ women with endometrial cancer		2.1%	1.7%
% 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ men with prostate cancer		9.3%	11.5%
% 65+ with hypothyroidism B 17.2% 20.8% % 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with benign prostatic hyperplasia		34.0%	36.8%
% 65+ with anemia 37.8% 37.3% % 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with HIV/AIDS		0.04%	0.05%
% 65+ with chronic kidney disease 21.5% 22.3% % 65+ with liver diseases 8.1% 6.9%	% 65+ with hypothyroidism	В	17.2%	20.8%
% 65+ with liver diseases 8.1% 6.9%	% 65+ with anemia		37.8%	37.3%
	% 65+ with chronic kidney disease		21.5%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue 17.1% 18.6%	% 65+ with liver diseases		8.1%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue		17.1%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		3.5%	4.0%
% 65+ with epilepsy		2.8%	2.1%
% 65+ with traumatic brain injury		0.9%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	В	19.3%	22.9%
% 65+ with cataract	В	54.7%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.4%	7.1%
% 65+ with 4+ (out of 15) chronic conditions		57.3%	54.4%
% 65+ with 0 chronic conditions		9.7%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		13.1%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		13.4%	14.4%
% 65+ with self-reported vision difficulty		7.4%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	0.9%
% 65+ with self-reported cognition difficulty		6.4%	6.9%
% 65+ with self-reported ambulatory difficulty		18.4%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.8%	3.2%
% 65+ with self-reported self-care difficulty		7.2%	5.6%
% 65+ with self-reported independent living difficulty		7.0%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		8.3%	7.9%
% dually eligible for Medicare and Medicaid	*	9.4%	7.5%
% 60+ with a regular doctor		93.5%	95.8%
% 60+ who did not see doctor when needed due to cost		6.5%	5.4%
# of primary care providers within 5 miles		10	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		1	74
# of home health agencies		4	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	4.9	6.3
Emergency room visits/1000 persons 65+ years per year		606	572
Part D monthly prescription fills per person per year		49.9	49.1
Home health visits per year		2.3	2.5
Durable medical equipment claims per year		2.2	2.0
Inpatient hospital stays/1000 persons 65+ years per year		243	237
Medicare inpatient hospital readmissions (as % of admissions)		12.5%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		97	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports		3.7%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		0	N/A
% of grandparents raising grandchildren		1.3%	0.8%
% of grandparents who live with grandchildren		2.8%	2.5%
# of assisted living sites		2	134
% of vacant homes in community		3.2%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		92.0%	93.0%
% 60+ who used Internet in last month		70.4%	77.6%
Voter participation rate in 2018 election (age 18+)		46.7%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		236	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		51	586
Property crime rate /100,000 persons		2,535	2,012
% 65+ who own a motor vehicle		100.0%	91.0%
% 60+ who always drive wearing a seatbelt	W	66.1%	77.1%
# of fatal crashes involving adult age 60+/town		0	151
# of fatal crashes involving adult age 60+/county		18	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		1.2%	5.4%
% 60+ receiving food stamps past year		6.7%	5.7%
% 65+ employed past year		21.5%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		12.9%	18.2%
% households with annual income \$20,000-\$49,999		54.9%	36.5%
% households with annual income > \$50,000		32.3%	45.3%
% 60+ own home		84.2%	79.9%
% 60+ have mortgage on home		41.2%	35.3%
% 65+ households spend >35% of income on housing (renter)		5.3%	8.7%
% 65+ households spend >35% of income on housing (owner)		17.2%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,860	\$25,284	1.02
Single, renter, good health	\$26,376	\$26,400	1.00
Couple, homeowner without mortgage, good health	\$37,800	\$37,128	1.02
Couple, renter, good health	\$38,316	\$38,244	1.00

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019