

Portsmouth is a city in Rockingham County home to a historic seaport and a series of historic house museums. There are 3,740 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of obesity, high cholesterol, cataracts, osteoarthritis/rheumatoid arthritis, and endometrial cancer. However, they had higher rates of excessive drinking, schizophrenia, ischemic heart disease, peripheral vascular disease, osteoporosis, breast cancer, benign prostatic hyperplasia, hypothyroidism, and fibromyalgia. Relative to the state, they are more likely to take the health promotion steps of meeting CDC guidelines for both muscle-strengthening and aerobic physical activity and getting screened for colorectal cancer. Community resources to support healthy aging include 105 primary care providers, a hospital, and 2 nursing homes within 5 miles, 11 home health agencies, a community health center, a memory cafe, a dementia support group, a senior center, 5 assisted living sites, 3 universities or community colleges, a public library, a YMCA, and access to broadband. Portsmouth city is a designated Age-Friendly community.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		21,458	1,327,503
Population 60 years or older as % of total population		23.2%	22.7%
Total population 60 years or older		4,975	301,630
Population 65 years or older as % of total population		17.4%	15.8%
Total population 65 years or older		3,740	210,385
% 65-74 years		51.1%	58.5%
% 75-84 years		32.4%	28.6%
% 85 years or older		16.5%	12.9%
Gender (65+ population)			
% female		59.5%	54.7%
Race/Ethnicity (65+ population)			
% White		96.0%	97.7%
% African American		2.1%	0.5%
% Asian		1.0%	0.9%
% Other		0.9%	0.9%
% Hispanic/Latino		1.9%	0.9%
Marital Status (65+ population)			
% married		44.3%	58.5%
% divorced/separated		21.4%	14.0%
% widowed		26.1%	22.9%
% never married		8.2%	4.6%
Education (65+ population)			
% with less than high school education		9.1%	12.3%
% with high school or some college		44.1%	57.1%
% with college degree		46.9%	30.6%
% of 65+ population living alone		39.6%	26.1%
% of 65+ population who speak only English at home		93.8%	91.3%
% of 65+ population who are veterans of military service		24.1%	24.8%
Age-sex adjusted 1-year mortality rate		4.2%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		6.0%	3.6%
% moved from different county in New Hampshire		1.2%	1.0%
% moved from different state		3.6%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		81.4%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity	В	35.7%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		66.6%	58.6%
% 60+ met CDC guidelines for both types of physical activities	В	29.8%	19.9%
% 60+ getting recommended hours of sleep		72.7%	66.4%
% 60+ injured in a fall within last 12 months		10.4%	10.4%
% 65+ had hip fracture		3.8%	3.3%
% 60+ with self-reported fair or poor health status		16.1%	16.5%
% 60+ with 15+ physically unhealthy days last month		10.4%	12.5%
% 60+ with physical exam/check-up in past year		88.8%	86.5%
% 60+ met CDC preventive health screening goals		40.6%	40.3%
% 60+ flu shot past year		57.9%	59.3%
% 65+ with pneumonia vaccine		78.2%	77.8%
% 60+ with cholesterol screening		97.3%	95.3%
% 60+ women with a mammogram within last 2 years		85.6%	79.3%
% 60+ with colorectal cancer screening	В	84.7%	77.0%
% 60+ with HIV test		14.9%	13.5%
% 60+ current smokers		7.3%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		23.1%	29.0%
% 60+ with annual dental exam		81.8%	75.7%
# of dentists per 100,000 persons (all ages) (county)		72	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		23.7%	20.2%
% 60+ self-reported obese		20.1%	27.2%
% 65+ clinically diagnosed obese	В	14.2%	16.7%
% 65+ with high cholesterol	В	69.8%	72.2%
% 60+ excessive drinking	W	16.2%	9.2%
% 65+ with poor supermarket access		6.1%	28.4%

BEHAVIORAL HEALTH 6.6+ with 15+ days poor mental health last month 6.4% 6.98 6.65+ with depression 29.4% 22.88 6.65+ with depression 23.4% 21.9% 6.65+ with bipolar disorders 3.4% 3.1% 6.65+ with bipolar disorders W 6.4% 4.9% 6.65+ with personality disorders W 6.4% 4.9% 6.65+ with personality disorders W 6.4% 4.9% 6.65+ with personality disorders W 6.4% 1.5% 1.1% # opioid deaths (all ages) (county) 276 1.279 6.5% 6.5% 5.5% 6.5% <	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression 29.4% 28.8% % 65+ with anxiety disorders 23.4% 21.9% % 65+ with bipolar disorders 3.4% 3.1% % 65+ with post-traumatic stress disorder 1.8% 4.4% % 65+ with post-traumatic stress disorders W 6.4% 4.9% % 65+ with schizophrenia & other psychotic disorders W 6.4% 4.9% % 65+ with post-traumatic stress disorders W 6.4% 4.9% % 65+ with schizophrenia & other psychotic disorders W 6.4% 4.9% % 65+ with schizophrenia & other psychotic disorders W 6.5% 5.5% % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with bobacco use disorders 10.0% 6.5% M 65+ with bobacco use disorders 4.78 4.8 M 65+ with bobacco use disorders 11.9% 1.20 M 65+ with bobacco use disorders 11.9% 12.0% M 65+ with brobacco use disorders 11.9% 12.0% M 65+ with stream 11.9% 12.0% M 65+ with brobacco u	BEHAVIORAL HEALTH			
% 65+ with pipolar disorders 3.4% 21.9% % 65+ with pipolar disorders 3.4% 3.1% % 65+ with pipolar disorders W 6.4% 1.4% % 65+ with personality disorders W 6.4% 4.9% % 65+ with personality disorders 1.5% 1.1% % 65+ with personality disorders 1.5% 1.1% % 65+ with personality disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with dubstace use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with dubstace use disorders 11.9% 12.0% % 65+ with dubstace use disorders (drug use +/or alcohol abuse) 28.1% 28.2% % 65+ with dubstace dubstace use disorders 11.9% 12.0% % 65+ with dubstace dubstace use or related dementias 11.9% 12.0% % 65+ with stroke 10.9% 65+ with hubsta	% 60+ with 15+ days poor mental health last month		6.4%	6.9%
% 65+ with bipolar disorders 3.4% 3.1% % 65+ with post-traumatic stress disorder 1.8% 1.4% % 65+ with personality disorders W 6.4% 4.9% % 65+ with personality disorders W 6.4% 4.9% % 65+ with with personality disorders 1.5% 1.1279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with tobacco use disorders 10.0% 10.0% CHRONIC DISEASE % 65+ with diabetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with stroke 10.9% 10.8% % 65+ with troke inchronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with presention 69.1% 70.2% % 65+ with presention 4,7% 4,5% % 65+ with trial fibrillation 15.6% 14.7% % 65+ with osteoperiosis W 16.4% 14.7% % 65+ with osteoporrosis W 20.2% 17.4%	% 65+ with depression		29.4%	28.8%
% 65+ with post-traumatic stress disorder 1.8% 1.4% % 65+ with schizophrenia & other psychotic disorders W 6.4% 4.9% % 65+ with personality disorders 1.5% 1.1% * opioid deaths (all ages) (county) 276 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with tubbacco use disorders 10.0% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with stroke 10.9% 10.8% % 65+ with obstructive pulmonary disease 19.4% 20.5% % 65+ with hypertension 99.1% 70.2% % 65+ with personality is schemic heart attack 4.7% 4.5% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with congestive heart failure 18.0% 14.7% % 65+ with osteoporosis W 16.4% 14.7% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with osteoporosis W 1.9% 2.8%	% 65+ with anxiety disorders		23.4%	21.9%
% 65+ with schizophrenia & other psychotic disorders W 6.4% 4.9% % 65+ with personality disorders 1.5% 1.1% # opioid deaths (all ages) (county) 276 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% 65+ with tobacco use disorders 10.0% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with stroke 10.9% 10.8% 65+ with stroke 10.9% 20.5% 65+ with strokine obstructive pulmonary disease 19.4% 20.5% 65+ with asthma 13.1% 13.0% 65+ with asthma 19.4% 20.5% 65+ with apperture of the periphers of the periphe	% 65+ with bipolar disorders		3.4%	3.1%
% 65+ with personality disorders 1.5% 1.1% # opioid deaths (all ages) (county) 276 1,279 % 65+ with substance use disorders 10.0% 5.5% % 65+ with tobacco use disorders 10.0% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease or related dementias 11.9% 28.2% % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease 10.9% 10.8% % 65+ with orthoric obstructive pulmonary disease 19.4% 20.5% % 65+ with pertension 69.1% 20.5% % 65+ with pertension 69.1% 31.3% % 65+ with schemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with netwemia sand lymphomas 19.6% 45.5%	% 65+ with post-traumatic stress disorder		1.8%	1.4%
# opioid deaths (all ages) (county) % 65+ with substance use disorders (drug use +/or alcohol abuse) % 65+ with tobacco use disorders **CHRONIC DISEASE** % 65+ with Alzheimer's disease or related dementias **6 65+ with diabetes **6 65+ with diabetes **6 65+ with diabetes **6 65+ with chronic obstructive pulmonary disease **6 65+ with chronic obstructive pulmonary disease **6 65+ with hypertension **6 65+ with hypertension **6 65+ with hypertension **6 65+ with cheart disease **8 65+ with congestive heart failure **6 65+ with atrial fibrillation **6 65+ with osteografic atrihitis/rheumatoid arthritis **6 65+ with osteografic atrihitis/rheumatoid arthritis **8 65+ with osteografic atrihitis/rheumatoid arthritis **8 65+ with lung cancer **8 65+ with lung cancer **8 65+ women with breast cancer **8 65+ women with prostate cancer **9 65+ women with prostate cancer **9 65+ women with prostate cancer **9 65+ with HIV/AIDS **8 65+ with hypothyroidism **9 65+ with lung disease **9 11.5% **9 65+ with hypothyroidism **9 65+ with lung disease **9 12.5% **9 65+ with hypothyroidism **9 65+ with prophyroidism **9 65+ with prophyroidism **9 65+ with prophyroidism **9 65+ with lung disease **9 12.5% **9 65+ with hypothyroidism **9 65+ with chronic kidney disease **9 65+ with liver diseases **9 65+ with liver diseases **9 65- with li	% 65+ with schizophrenia & other psychotic disorders	W	6.4%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 5.8% 5.5% % 65+ with tobacco use disorders 10.0% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with disbetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with natrial fibrillation 15.6% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with up prostate cancer W 11.4% 1.7% % 65+ women with breast cancer B 1.1% 1.7% % 65+ with hurlynatic	% 65+ with personality disorders		1.5%	1.1%
% 65+ with tobacco use disorders 10.0% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with Alzheimer's disease or related dementias 11.9% 28.2% % 65+ with diabetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with stroke 69.1% 70.2% % 65+ with hypertension 69.1% 70.2% % 65+ with pripheral vascular disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 20.5% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer W 11.1% 1.6% % 65+ women with prostate c	# opioid deaths (all ages) (county)		276	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with diabetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ with pyertension 4.7% 4.5% 65+ with schemic heart disease W 38.9% 34.3% 65+ with congestive heart failure 18.0% 17.7% 65+ with peripheral vascular disease W 16.4% 14.7% 65+ with peripheral vascular disease W 16.4% 14.7% 65+ with osteoporosis W 20.2% 17.4% 65+ with lug cancer	% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.8%	5.5%
% 65+ with Alzheimer's disease or related dementias 11.9% 12.0% % 65+ with diabetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with leukemias and lymphomas 1.9% 20.2% % 65+ with lung cancer 1.7% 1.6% % 65+ with peripheral vascular disease W 11.4% 9.8% % 65+ with lung cancer W 20.2% 17.4% % 65+ with lung cancer W 11.7% 1.6% % 65+ women with prostate cancer W 11.4% 9	% 65+ with tobacco use disorders		10.0%	10.4%
% 65+ with diabetes 28.1% 28.2% % 65+ with stroke 10.9% 10.8% % 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with lug cancer W 20.2% 17.4% % 65+ with lug cancer 1.9% 2.0% % 65+ with colon cancer 2.8% 2.4% % 65+ with prostate cancer W 1.1% 9.8% % 65+ women with prostate cancer B 1.1% 1.7% % 65+ men with prostate cancer W 40.0% 36.8% % 65+ with hypothyroidism W 40.0% 36.8%	CHRONIC DISEASE			
% 65+ with stroke 10.9% 10.8% % 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with lung cancer W 20.2% 17.4% % 65+ with lung cancer 1.7% 1.6% % 65+ with peripheral vascular disease W 11.4% 2.0% % 65+ with lung cancer 17.4% 49.1% 49.1% % 65+ with lung cancer 17.7% 1.6% % 65+ worth lung cancer W 11.4% 9.8% % 65+ worth loon cancer B 1.1% 1.7% % 65+ worth peripheral vascular disease W 10.0%	% 65+ with Alzheimer's disease or related dementias		11.9%	12.0%
% 65+ with chronic obstructive pulmonary disease 19.4% 20.5% % 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with prostate cancer B 1.1% 1.7% % 65+ men with prostate cancer B 1.1% 1.5% % 65+ with helign prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with chronic kidney dise	% 65+ with diabetes		28.1%	28.2%
% 65+ with asthma 13.1% 13.0% % 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoporosis B 45.5% 49.1% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ men with prostate cancer B 1.1% 1.7% % 65+ men with prostate cancer B 1.1% 1.5% % 65+ with heingn prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with nemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 6.9% <td>% 65+ with stroke</td> <td></td> <td>10.9%</td> <td>10.8%</td>	% 65+ with stroke		10.9%	10.8%
% 65+ with hypertension 69.1% 70.2% % 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with attrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with lung cancer 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer W 11.4% 9.8% % 65+ women with prostate cancer W 11.4% 9.8% % 65+ men with prostate cancer B 1.1% 1.7% % 65+ men with prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with nemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7%	% 65+ with chronic obstructive pulmonary disease		19.4%	20.5%
% 65+ ever had a heart attack 4.7% 4.5% % 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with asthma		13.1%	13.0%
% 65+ with ischemic heart disease W 38.9% 34.3% % 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with prostate cancer B 1.1% 1.7% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with nemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with hypertension		69.1%	70.2%
% 65+ with congestive heart failure 18.0% 17.7% % 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with colon cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with prostate cancer B 1.1% 1.7% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with liver diseases 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ ever had a heart attack		4.7%	4.5%
% 65+ with atrial fibrillation 15.6% 14.4% % 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer B 1.1% 1.7% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with ischemic heart disease	W	38.9%	34.3%
% 65+ with peripheral vascular disease W 16.4% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with congestive heart failure		18.0%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis B 45.5% 49.1% % 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with atrial fibrillation		15.6%	14.4%
% 65+ with osteoporosis W 20.2% 17.4% % 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with peripheral vascular disease	W	16.4%	14.7%
% 65+ with leukemias and lymphomas 1.9% 2.0% % 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with osteoarthritis/rheumatoid arthritis	В	45.5%	49.1%
% 65+ with lung cancer 1.7% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with osteoporosis	W	20.2%	17.4%
% 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with leukemias and lymphomas		1.9%	2.0%
% 65+ women with breast cancer W 11.4% 9.8% % 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with lung cancer		1.7%	1.6%
% 65+ women with endometrial cancer B 1.1% 1.7% % 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with colon cancer		2.8%	2.4%
% 65+ men with prostate cancer 12.2% 11.5% % 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ women with breast cancer	W	11.4%	9.8%
% 65+ with benign prostatic hyperplasia W 40.0% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ women with endometrial cancer	В	1.1%	1.7%
% 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ men with prostate cancer		12.2%	11.5%
% 65+ with hypothyroidism W 22.9% 20.8% % 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with benign prostatic hyperplasia	W	40.0%	36.8%
% 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%	% 65+ with HIV/AIDS		0.04%	0.05%
% 65+ with anemia 38.4% 37.3% % 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%		W	22.9%	
% 65+ with chronic kidney disease 21.7% 22.3% % 65+ with liver diseases 6.7% 6.9%				
% 65+ with liver diseases 6.7% 6.9%				
	· · · · · · · · · · · · · · · · · · ·			
	% 65+ with fibromyalgia, chronic pain and fatigue	W	21.3%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		4.6%	4.0%
% 65+ with epilepsy		2.3%	2.1%
% 65+ with traumatic brain injury		1.0%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma		23.7%	22.9%
% 65+ with cataract	В	58.0%	61.2%
% 65+ with pressure ulcer or chronic ulcer		8.0%	7.1%
% 65+ with 4+ (out of 15) chronic conditions		54.1%	54.4%
% 65+ with 0 chronic conditions	В	12.4%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		14.4%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		14.3%	14.4%
% 65+ with self-reported vision difficulty	В	2.7%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	0.9%
% 65+ with self-reported cognition difficulty	В	3.2%	6.9%
% 65+ with self-reported ambulatory difficulty		16.1%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.3%	3.2%
% 65+ with self-reported self-care difficulty		5.4%	5.6%
% 65+ with self-reported independent living difficulty		10.4%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	4.6%	7.9%
% dually eligible for Medicare and Medicaid		8.1%	7.5%
% 60+ with a regular doctor		95.2%	95.8%
% 60+ who did not see doctor when needed due to cost		5.4%	5.4%
# of primary care providers within 5 miles		105	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		2	74
# of home health agencies		11	49
# of community health centers		1	22
# of adult day health centers		0	21
# of memory cafes		1	12
# of dementia-related support groups		1	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	7.8	6.3
Emergency room visits/1000 persons 65+ years per year		648	572
Part D monthly prescription fills per person per year		49.9	49.1
Home health visits per year		2.8	2.5
Durable medical equipment claims per year	*	1.4	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	280	237
Medicare inpatient hospital readmissions (as % of admissions)		18.0%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year	*	91	76
# skilled nursing home Medicare beds/1000 persons 65+ years		64	33
% 65+ getting Medicaid long term services and supports		3.9%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Yes	Yes
# of senior centers		1	44
Air pollution: annual # of unhealthy days for 65+ (county)		2	N/A
% of grandparents raising grandchildren		0.5%	0.8%
% of grandparents who live with grandchildren		1.6%	2.5%
# of assisted living sites		5	134
% of vacant homes in community		4.5%	16.0%
# of universities and community colleges		3	41
# of public libraries		1	234
# of YMCAs		1	12
% in county with access to broadband (all ages)		99.0%	93.0%
% 60+ who used Internet in last month		75.1%	77.6%
Voter participation rate in 2018 election (age 18+)		62.5%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		190	207
Homicide rate /100,000 persons (county)		1	1
# firearm fatalities (county)		92	586
Property crime rate /100,000 persons		2,271	2,012
% 65+ who own a motor vehicle		90.8%	91.0%
% 60+ who always drive wearing a seatbelt		80.9%	77.1%
# of fatal crashes involving adult age 60+/town		3	151
# of fatal crashes involving adult age 60+/county		31	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		4.3%	5.4%
% 60+ receiving food stamps past year		2.8%	5.7%
% 65+ employed past year		25.3%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		15.9%	18.2%
% households with annual income \$20,000-\$49,999		30.3%	36.5%
% households with annual income > \$50,000		53.8%	45.3%
% 60+ own home		69.3%	79.9%
% 60+ have mortgage on home		31.0%	35.3%
% 65+ households spend >35% of income on housing (renter)		15.7%	8.7%
% 65+ households spend >35% of income on housing (owner)		16.0%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,536	\$25,284	1.01
Single, renter, good health	\$26,712	\$26,400	1.01
Couple, homeowner without mortgage, good health	\$37,092	\$37,128	1.00
Couple, renter, good health	\$38,268	\$38,244	1.00

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are not due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016). ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019