
Belmont (Belknap)

Belmont is a town in Belknap County and home to several historic places including the Belmont Public Library and the Bandstand. There are 1,138 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of schizophrenia and other psychotic disorders, Alzheimer's disease or related dementias, osteoarthritis/rheumatoid arthritis, osteoporosis, leukemia and lymphomas, hypothyroidism, migraine, traumatic brain injury, glaucoma, and cataracts. However, they had higher rates of tobacco use disorders, chronic obstructive pulmonary disease, and hypertension. They are less likely to take the health promotion steps of undergoing colorectal cancer screening or wearing a seatbelt while driving. Community resources to support healthy aging include 2 primary care providers within 5 miles, 5 home health agencies, a public library, and access to broadband.







HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		4.3%	3.6%
% moved from different county in New Hampshire		0.0%	1.0%
% moved from different state		0.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		71.7%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		21.8%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		56.3%	58.6%
% 60+ met CDC guidelines for both types of physical activities		16.7%	19.9%
% 60+ getting recommended hours of sleep		58.8%	66.4%
% 60+ injured in a fall within last 12 months		9.6%	10.4%
% 65+ had hip fracture		3.2%	3.3%
% 60+ with self-reported fair or poor health status		17.8%	16.5%
% 60+ with 15+ physically unhealthy days last month		12.4%	12.5%
% 60+ with physical exam/check-up in past year		90.8%	86.5%
% 60+ met CDC preventive health screening goals		38.0%	40.3%
% 60+ flu shot past year		58.8%	59.3%
% 65+ with pneumonia vaccine		77.1%	77.8%
% 60+ with cholesterol screening		93.2%	95.3%
% 60+ women with a mammogram within last 2 years		76.5%	79.3%
% 60+ with colorectal cancer screening	W	69.3%	77.0%
% 60+ with HIV test		9.2%	13.5%
% 60+ current smokers		11.5%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		31.2%	29.0%
% 60+ with annual dental exam		68.9%	75.7%
# of dentists per 100,000 persons (all ages) (county)		69	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		16.8%	20.2%
% 60+ self-reported obese		27.5%	27.2%
% 65+ clinically diagnosed obese		14.5%	16.7%
% 65+ with high cholesterol		72.9%	72.2%
% 60+ excessive drinking		11.0%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

% 60+ with 15+ days poor mental health last month 6.9% 6.9% 6.9% % 65+ with depression 27.0% 28.8% % 65+ with biolar disorders 2.1.4% 21.9% % 65+ with bost-traumatic stress disorder 1.7% 1.44 % 65+ with post-traumatic stress disorders B 3.7% 4.99 % 65+ with post-traumatic stress disorders B 3.7% 4.99 % 65+ with post-traumatic stress disorders (drug use +/or alcohol abuse) 5.6% 5.55 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.55 % 65+ with babcacc use disorders (drug use +/or alcohol abuse) 5.6% 5.55 % 65+ with abustance use disorders (drug use +/or alcohol abuse) 5.6% 5.55 % 65+ with babcacc use disorders (drug use +/or alcohol abuse) 5.6% 5.55 % 65+ with babcacc use disorders W 12.9% 10.49 CHEONIC DISEASE 12.0% 10.08 % 65+ with habibetes 31.0% 28.6% 20.5% % 65+ with stroke 9.9% 10.89 65 % 65+ with stroke 9.9%	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression 27.0% 28.89 % 65+ with anxiety disorders 21.4% 21.99 % 65+ with post-rtaumatic stress disorder 1.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 3.7% 4.99 % 65+ with schizophrenia & other psychotic disorders B 3.7% 4.99 % 65+ with schizophrenia & other psychotic disorders B 3.7% 4.99 % 65+ with schizophrenia & other psychotic disorders W 12.9% 10.49 CHRONC DISEASE W 12.9% 10.49 CHRONC DISEASE 9.9% 10.8% 20.59 % 65+ with disbetes 31.0% 28.29 20.59 % 65+ with chronic obstructive pulmonary disease W 23.6% 20.59 % 65+ with hypertension W 73.6% 70.29 % 65+ with asthma 12.8% 13.00 34.39 % 65+ with other is disease 34.0% 34.39 34.39 % 65+ with other is disease 34.0% 34.39 34.39 % 65+ with other is disease 34.0% </td <td>BEHAVIORAL HEALTH</td> <td></td> <td></td> <td></td>	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders 21.4% 21.9% % 65+ with bolar disorders 2.3% 3.19 % 65+ with borst-traumatic stress disorder 1.7% 1.49 % 65+ with bersonality disorders B 3.7% 4.99 % 65+ with bersonality disorders 0.8% 1.11 # opioid deaths (all ages) (county) 53 1.27 % 65+ with bersonality disorders W 12.9% 10.49 CHRONC DISEASE W 12.0% 10.49 CHRONIC Distase 31.0% 28.29 % 65+ with Alzheimer's disease or related dementias B 8.7% 12.0% % 65+ with Alzheimer's disease or related dementias B 8.7% 12.0% % 65+ with Alzheimer's disease or related dementias B 8.7% 12.0% % 65+ with Alzheimer's disease 31.0% 28.29 30.0% % 65+ with shitha 12.8% 20.59 36.54 31.0% 28.29 36.64 31.0% 34.39 36.54 31.0% 34.39 36.54 31.0% 34.39 36.54 31.	% 60+ with 15+ days poor mental health last month		6.9%	6.9%
% 65+ with bipolar disorders 2.3% 3.19 % 65+ with post-traumatic stress disorder 1.7% 1.49 % 65+ with personality disorders B 3.7% 4.99 % 65+ with personality disorders B 3.7% 4.99 % 65+ with personality disorders B 3.7% 4.99 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.59 % 65+ with substance use disorders W 12.9% 10.4% CHEONIC DISEASE	% 65+ with depression		27.0%	28.8%
% 65+ with post-traumatic stress disorder 1.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 3.7% 4.99 % 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 53 1.27 % 65+ with babacco use disorders W 12.9% 10.49 CHRONIC DISEASE W 12.9% 10.49 % 65+ with babacco use disorders W 12.9% 10.49 % 65+ with babacco use disorders W 12.9% 10.49 CHRONIC DISEASE 31.0% 28.29 % 65+ with diabetes 31.0% 28.29 % 65+ with astroke 9.9% 10.89 % 65+ with astroke 9.9% 10.89 % 65+ with stroke 9.9% 10.89 % 65+ with asthma 12.8% 13.00% % 65+ with asthma 12.8% 13.09 % 65+ with stroke 9.9% 10.49 % 65+ with asthma 12.8% 13.09 % 65+ with asthma 12.8% 13.09 % 65	% 65+ with anxiety disorders		21.4%	21.9%
% 65+ with schizophrenia & other psychotic disorders B 3.7% 4.99 % 65+ with personality disorders 0.8% 1.19 # oploid deaths (all ages) (county) 53 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.5% % 65+ with tobacco use disorders W 12.9% 10.49 CHRONC DISEASE 31.0% 28.29 % 65+ with diabetes 31.0% 28.29 % 65+ with stroke 9.9% 10.89 % 65+ with obstructive pulmonary disease W 23.6% 20.59 % 65+ with stroke 9.9% 10.89 34.0% 34.39 % 65+ with astma 12.8% 13.00% 26.43 34.0% 34.39 % 65+ with schemic heart disease 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% 34.39 34.0% <td>% 65+ with bipolar disorders</td> <td></td> <td>2.3%</td> <td>3.1%</td>	% 65+ with bipolar disorders		2.3%	3.1%
% 65+ with personality disorders 0.8% 1.19 # opioid deaths (all ages) (county) 53 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.5% % 65+ with babaco use disorders W 12.9% 10.4% CHRONIC DISEASE	% 65+ with post-traumatic stress disorder		1.7%	1.4%
# opioid deaths (all ages) (county) 53 1,27 % 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.59 % 65+ with tobacco use disorders W 12.9% 10.49 CHRONIC DISEASE	% 65+ with schizophrenia & other psychotic disorders	В	3.7%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) 5.6% 5.5% % 65+ with tobacco use disorders W 12.9% 10.49 CHRONIC DISEASE ************************************	% 65+ with personality disorders		0.8%	1.1%
% 65+ with tobacco use disorders W 12.9% 10.49 CHRONIC DISEASE 8.7% 12.09 % 65+ with Alzheimer's disease or related dementias B 8.7% 12.09 % 65+ with diabetes 31.0% 28.29 % 65+ with stroke 9.9% 10.89 % 65+ with stroke 9.9% 10.89 % 65+ with asthma 12.8% 33.09 % 65+ with asthma 12.8% 13.09 % 65+ with opertension W 73.6% 70.29 % 65+ with congestive heart failure 16.7% 17.77 % 65+ with congestive heart failure 16.7% 14.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 14.0% 17.49 % 65+ with osteoarthritis/rheumatoid arthritis B 14.0% 17.49 % 65+ with loug cancer 10.3% 9.89 <td># opioid deaths (all ages) (county)</td> <td></td> <td>53</td> <td>1,279</td>	# opioid deaths (all ages) (county)		53	1,279
CHRONIC DISEASE B 8.7% 12.0% % 65+ with Alzheimer's disease or related dementias B 8.7% 12.0% % 65+ with diabetes 31.0% 28.29 % 65+ with stroke 9.9% 10.8% % 65+ with stroke 9.9% 10.8% % 65+ with asthma 12.8% 13.0% % 65+ with ischemic heart disease 34.0% 34.3% % 65+ with atrial fibrillation 13.4% 14.4% % 65+ with astoporosis B 14.0% 17.4% % 65+ with osteoporosis B 14.0% 17.4% % 65+ with loug cancer 2.1% 1.6% 2.4% % 65+ with colon cancer 2.8% 2.4% 65+ with brenay trostat cancer 10.1%	% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.6%	5.5%
% 65+ with Alzheimer's disease or related dementias B 8.7% 12.0% % 65+ with diabetes 31.0% 28.29 % 65+ with stroke 9.9% 10.89 % 65+ with stroke 9.9% 10.89 % 65+ with chronic obstructive pulmonary disease W 23.6% 20.59 % 65+ with asthma 12.8% 13.09 65 % 65+ with asthma 12.8% 13.09 65 % 65+ with hypertension W 73.6% 70.29 % 65+ with schemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.77 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteopartivitis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoparosis B 11.0% 17.49 % 65+ with loug cancer 2.1% 1.6% 2.4% % 65+ with loug cancer 2.1% 1.6% 2.4% % 65+ with loug cancer 2.8% 2.4% 4.5% % 65+ with nug cancer 10.3% <td< td=""><td>% 65+ with tobacco use disorders</td><td>W</td><td>12.9%</td><td>10.4%</td></td<>	% 65+ with tobacco use disorders	W	12.9%	10.4%
% 65+ with diabetes 31.0% 28.29 % 65+ with stroke 9.9% 10.89 % 65+ with chronic obstructive pulmonary disease W 23.6% 20.59 % 65+ with asthma 12.8% 13.09 65+ % 65+ with hypertension W 73.6% 70.29 % 65+ with hypertension W 73.6% 70.29 % 65+ with congestive heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with colon cancer 2.1% 1.69 % 65+ with lung cancer 2.1% 1.69 % 65+ women with prestat cancer 10.3% 9.88 % 65+ women with prostate cancer 10.1% 11.59 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with holing norostatic hyperplasia 39.2% 36.88 %	CHRONIC DISEASE			
% 65+ with stroke 9.9% 10.89 % 65+ with chronic obstructive pulmonary disease W 23.6% 20.59 % 65+ with asthma 12.8% 13.00 % 65+ with hypertension W 73.6% 70.29 % 65+ ever had a heart attack 4.4% 4.59 % 65+ ever had a heart attack 4.4% 34.39 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ women with prostate cancer 10.3% 9.89 % 65+ women with prostate cancer 10.3% 9.89 % 65+ with heingn prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89	% 65+ with Alzheimer's disease or related dementias	В	8.7%	12.0%
% 65+ with chronic obstructive pulmonary disease W 23.6% 20.59 % 65+ with asthma 12.8% 13.00 % 65+ with hypertension W 73.6% 70.29 % 65+ ever had a heart attack 4.4% 4.59 % 65+ ever had a heart attack 4.4% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ woren with prostate cancer 2.1% 1.69 % 65+ woren with prostate cancer 10.3% 9.89 % 65+ woren with prostate cancer 10.1% 11.59 % 65+ woren with prostate cancer 10.1% 11.59 % 65+ woren with prostate cancer 0.04% 0.058 % 65+ with hulpy protyroidism B 15.7% 20.89	% 65+ with diabetes		31.0%	28.2%
% 65+ with asthma 12.8% 13.09 % 65+ with hypertension W 73.6% 70.29 % 65+ ever had a heart attack 4.4% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoorthritis/rheumatoid arthritis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with colon cancer 2.8% 2.49 % 65+ women with prostate cancer 10.3% 9.89 % 65+ women with nedometrial cancer 10.1% 11.59 % 65+ with heingn prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7%<	% 65+ with stroke		9.9%	10.8%
% 65+ with hypertension W 73.6% 70.29 % 65+ ever had a heart attack 4.4% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with colon cancer 2.1% 1.69 2.09 % 65+ with colon cancer 2.8% 2.49 3.89 % 65+ women with breast cancer 10.3% 9.89 3.92% 36.89 % 65+ women with prostate cancer 10.1% 11.59 3.92% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 3.92% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 3.92% 36.89 % 65+ with hyp	% 65+ with chronic obstructive pulmonary disease	W	23.6%	20.5%
% 65+ ever had a heart attack 4.4% 4.59 % 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with congestive heart failure 13.4% 14.49 % 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with colon cancer 2.1% 1.69 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism 8 15.7% 20.89 % 65+ with hypothyroidism 8 15.7% 20.89 % 65+ with hypothyroidism 8 15.7% 20.89 % 65+ with chronic kidney disease 20.6% 22.39	% 65+ with asthma		12.8%	13.0%
% 65+ with ischemic heart disease 34.0% 34.39 % 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoarthritis/rheumatoid arthritis B 14.0% 17.49 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with colon cancer 2.1% 1.69 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with aremia 35.2% 37.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with hypertension	W	73.6%	70.2%
% 65+ with congestive heart failure 16.7% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteopartsis B 14.0% 17.49 % 65+ with osteopartsis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with lung cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 1.7% 1.79 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with chronic kidney disease 20.6%	% 65+ ever had a heart attack		4.4%	4.5%
% 65+ with atria fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.4% % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with long cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 10.3% 9.89 % 65+ with benign prostatic cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% <td>% 65+ with ischemic heart disease</td> <td></td> <td>34.0%</td> <td>34.3%</td>	% 65+ with ischemic heart disease		34.0%	34.3%
% 65+ with peripheral vascular disease 16.7% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with lung cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with congestive heart failure		16.7%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis B 44.1% 49.19 % 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with lung cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 1.7% 1.79 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with anemia 35.2% 37.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with chronic kidney disease 6.5% 6.99	% 65+ with atrial fibrillation		13.4%	14.4%
% 65+ with osteoporosis B 14.0% 17.49 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with lung cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 10.7% 1.79 % 65+ women with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with chronic kidney disease 6.5% 6.99	% 65+ with peripheral vascular disease		16.7%	14.7%
% 65+ with leukemias and lymphomas B 1.3% 2.09 % 65+ with lung cancer 2.1% 1.69 % 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 1.7% 1.79 % 65+ men with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with nemia 35.2% 37.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with osteoarthritis/rheumatoid arthritis	В	44.1%	49.1%
% 65+ with lung cancer 2.1% 1.6% % 65+ with colon cancer 2.8% 2.4% % 65+ women with breast cancer 10.3% 9.8% % 65+ women with endometrial cancer 10.3% 9.8% % 65+ women with prostate cancer 10.1% 11.7% % 65+ with benign prostatic hyperplasia 39.2% 36.8% % 65+ with hypothyroidism B 15.7% 20.8% % 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with osteoporosis	В	14.0%	17.4%
% 65+ with colon cancer 2.8% 2.49 % 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 1.7% 1.79 % 65+ men with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with leukemias and lymphomas	В	1.3%	2.0%
% 65+ women with breast cancer 10.3% 9.89 % 65+ women with endometrial cancer 1.7% 1.79 % 65+ men with prostate cancer 10.1% 11.59 % 65+ with benign prostatic hyperplasia 39.2% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism B 15.7% 20.89 % 65+ with anemia 35.2% 37.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with lung cancer		2.1%	1.6%
% 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer 10.1% 11.5% % 65+ with benign prostatic hyperplasia 39.2% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 15.7% 20.8% % 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with colon cancer		2.8%	2.4%
% 65+ men with prostate cancer 10.1% 11.5% % 65+ with benign prostatic hyperplasia 39.2% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 15.7% 20.8% % 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ women with breast cancer		10.3%	9.8%
% 65+ with benign prostatic hyperplasia 39.2% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 15.7% 20.8% % 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ women with endometrial cancer		1.7%	1.7%
% 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism B 15.7% 20.8% % 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ men with prostate cancer		10.1%	11.5%
% 65+ with hypothyroidism B 15.7% 20.89 % 65+ with anemia 35.2% 37.39 % 65+ with chronic kidney disease 20.6% 22.39 % 65+ with liver diseases 6.5% 6.99	% 65+ with benign prostatic hyperplasia		39.2%	36.8%
% 65+ with anemia 35.2% 37.3% % 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with HIV/AIDS		0.04%	0.05%
% 65+ with chronic kidney disease 20.6% 22.3% % 65+ with liver diseases 6.5% 6.9%	% 65+ with hypothyroidism	В	15.7%	20.8%
% 65+ with liver diseases 6.5% 6.9%	% 65+ with anemia		35.2%	37.3%
	% 65+ with chronic kidney disease		20.6%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue 16.8% 18.6%	% 65+ with liver diseases		6.5%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue		16.8%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache	В	3.1%	4.0%
% 65+ with epilepsy		1.9%	2.1%
% 65+ with traumatic brain injury	В	0.6%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	В	16.9%	22.9%
% 65+ with cataract	В	54.3%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.2%	7.1%
% 65+ with 4+ (out of 15) chronic conditions		53.0%	54.4%
% 65+ with 0 chronic conditions		10.1%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		20.6%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		13.4%	14.4%
% 65+ with self-reported vision difficulty		3.8%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.6%	0.9%
% 65+ with self-reported cognition difficulty		3.5%	6.9%
% 65+ with self-reported ambulatory difficulty		19.8%	18.8%
% 65+ with clinical diagnosis of mobility impairments		2.7%	3.2%
% 65+ with self-reported self-care difficulty		4.9%	5.6%
% 65+ with self-reported independent living difficulty		7.8%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	10.6%	7.9%
% dually eligible for Medicare and Medicaid		9.0%	7.5%
% 60+ with a regular doctor		93.5%	95.8%
% 60+ who did not see doctor when needed due to cost		6.5%	5.4%
# of primary care providers within 5 miles		2	2,961
# of hospitals within 5 miles		0	26
# of nursing homes within 5 miles		0	74
# of home health agencies		5	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	6.9	6.3
Emergency room visits/1000 persons 65+ years per year		588	572
Part D monthly prescription fills per person per year		50.7	49.1
Home health visits per year		2.0	2.5
Durable medical equipment claims per year	*	2.9	2.0
Inpatient hospital stays/1000 persons 65+ years per year		255	237
Medicare inpatient hospital readmissions (as % of admissions)		15.8%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		91	76
# skilled nursing home Medicare beds/1000 persons 65+ years		0	33
% 65+ getting Medicaid long term services and supports	*	2.1%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		0	44
Air pollution: annual # of unhealthy days for 65+ (county)		0	N/A
% of grandparents raising grandchildren		0.3%	0.8%
% of grandparents who live with grandchildren		2.8%	2.5%
# of assisted living sites		0	134
% of vacant homes in community		24.6%	16.0%
# of universities and community colleges		0	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		95.0%	93.0%
% 60+ who used Internet in last month		70.4%	77.6%
Voter participation rate in 2018 election (age 18+)		46.9%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		256	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		31	586
Property crime rate /100,000 persons		2,880	2,012
% 65+ who own a motor vehicle		98.3%	91.0%
% 60+ who always drive wearing a seatbelt	W	66.1%	77.1%
# of fatal crashes involving adult age 60+/town		0	151
# of fatal crashes involving adult age 60+/county		6	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		5.6%	5.4%
% 60+ receiving food stamps past year		5.6%	5.7%
% 65+ employed past year		26.3%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		20.5%	18.2%
% households with annual income \$20,000-\$49,999		40.5%	36.5%
% households with annual income > \$50,000		39.0%	45.3%
% 60+ own home		89.3%	79.9%
% 60+ have mortgage on home		33.6%	35.3%
% 65+ households spend >35% of income on housing (renter)		0.0%	8.7%
% 65+ households spend >35% of income on housing (owner)		28.4%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$24,000	\$25,284	0.95
Single, renter, good health	\$25,032	\$26,400	0.95
Couple, homeowner without mortgage, good health	\$35,940	\$37,128	0.97
Couple, renter, good health	\$36,972	\$38,244	0.97

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019