

## Berlin (Coös)

Berlin is a city in northern New Hampshire located along the Androscoggin River. There are 1,938 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of bipolar and personality disorders, breast cancer, benign prostatic hyperplasia, migraine, traumatic brain injury, and visual impairment. However, they had higher rates of high cholesterol, Alzheimer's disease, diabetes, stroke, COPD, hypertension, ischemic heart disease, peripheral vascular disease, anemia, chronic kidney disease, fibromyalgia/chronic pain/fatigue, cataracts, and hearing impairment. They are less likely to take the health promotion steps of doing physical activity within the past month or following the CDC guidelines for both muscle-strengthening and aerobic physical activity, meeting the CDC preventive health screening goals, having an annual dental exam, or wearing a seatbelt while driving. Community resources to support healthy aging include a hospital and 2 nursing homes within 5 miles, 2 home health agencies, 3 community health centers, an adult day health center, a senior center, a university or community college, and a public library.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		10,154	1,327,503
Population 60 years or older as % of total population		25.3%	22.7%
Total population 60 years or older		2,565	301,630
Population 65 years or older as % of total population		19.1%	15.8%
Total population 65 years or older		1,938	210,385
% 65-74 years		41.6%	58.5%
% 75-84 years		35.6%	28.6%
% 85 years or older		22.8%	12.9%
Gender (65+ population)			
% female		58.9%	54.7%
Race/Ethnicity (65+ population)			
% White		97.9%	97.7%
% African American		0.4%	0.5%
% Asian		0.0%	0.9%
% Other		1.8%	0.9%
% Hispanic/Latino		0.5%	0.9%
Marital Status (65+ population)			
% married		47.4%	58.5%
% divorced/separated		9.7%	14.0%
% widowed		34.0%	22.9%
% never married		8.9%	4.6%
Education (65+ population)			
% with less than high school education		20.3%	12.3%
% with high school or some college		72.5%	57.1%
% with college degree		7.2%	30.6%
% of 65+ population living alone		39.4%	26.1%
% of 65+ population who speak only English at home		55.9%	91.3%
% of 65+ population who are veterans of military service		26.8%	24.8%
Age-sex adjusted 1-year mortality rate		4.9%	4.1%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		4.9%	3.6%
% moved from different county in New Hampshire		0.6%	1.0%
% moved from different state		0.9%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month	W	64.7%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity	W	17.7%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		61.2%	58.6%
% 60+ met CDC guidelines for both types of physical activities	W	12.7%	19.9%
% 60+ getting recommended hours of sleep		64.2%	66.4%
% 60+ injured in a fall within last 12 months		11.3%	10.4%
% 65+ had hip fracture		3.7%	3.3%
% 60+ with self-reported fair or poor health status		24.8%	16.5%
% 60+ with 15+ physically unhealthy days last month		15.2%	12.5%
% 60+ with physical exam/check-up in past year		90.7%	86.5%
% 60+ met CDC preventive health screening goals	W	26.9%	40.3%
% 60+ flu shot past year		63.0%	59.3%
% 65+ with pneumonia vaccine		80.2%	77.8%
% 60+ with cholesterol screening		92.3%	95.3%
% 60+ women with a mammogram within last 2 years		75.9%	79.3%
% 60+ with colorectal cancer screening		69.1%	77.0%
% 60+ with HIV test		9.9%	13.5%
% 60+ current smokers		11.1%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth	W	43.3%	29.0%
% 60+ with annual dental exam	W	60.2%	75.7%
# of dentists per 100,000 persons (all ages) (county)		59	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		13.9%	20.2%
% 60+ self-reported obese		26.2%	27.2%
% 65+ clinically diagnosed obese		17.0%	16.7%
% 65+ with high cholesterol	W	77.7%	72.2%
% 60+ excessive drinking		7.2%	9.2%
% 65+ with poor supermarket access		39.5%	28.4%

% 65+ with depression         30.6%         28.8%           % 65+ with anxiety disorders         21.2%         21.9%           % 65+ with post-traumatic stress disorder         1.3%         1.4%           % 65+ with schizophrenia & other psychotic disorders         4.5%         4.9%           % 65+ with schizophrenia & other psychotic disorders         8         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.278           % 65+ with schizophrenia & other psychotic disorders         10.4%         10.4%           CHRONC DISEASE         0.6%         1.1%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with asthma         13.8%         13.0%           % 65+ with hypertension         W         14.7%         12.0%           % 65+ with hypertension         W         77.7%         70.2%           % 65+ with sthma         13.8%         13.0%         46.5%           % 65+ with other tailatek         5.4%         4.5%         4.5%           % 65+ with other tailatek         5.4%         4.5%         4.5% </th <th>HEALTHY AGING INDICATORS</th> <th>BETTER / WORSE STATE RATE<sup>1</sup></th> <th>COMMUNITY ESTIMATE</th> <th>STATE ESTIMATE</th>	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression         30.6%         28.8%           % 65+ with anxiety disorders         21.2%         21.9%           % 65+ with post-traumatic stress disorder         1.3%         1.4%           % 65+ with schizophrenia & other psychotic disorders         4.5%         4.9%           % 65+ with schizophrenia & other psychotic disorders         8         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.278           % 65+ with schizophrenia & other psychotic disorders         10.4%         10.4%           CHRONC DISEASE         0.6%         1.1%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with asthma         13.8%         13.0%           % 65+ with hypertension         W         14.7%         12.0%           % 65+ with hypertension         W         77.7%         70.2%           % 65+ with sthma         13.8%         13.0%         46.5%           % 65+ with other tailatek         5.4%         4.5%         4.5%           % 65+ with other tailatek         5.4%         4.5%         4.5% </td <td>BEHAVIORAL HEALTH</td> <td></td> <td></td> <td></td>	BEHAVIORAL HEALTH			
% 65+ with anxiety disorders         21.2%         21.9%           % 65+ with biolar disorders         B         1.7%         3.1%           % 65+ with bersonality disorders         4.5%         4.9%           % 65+ with bersonality disorders         B         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.273           % 65+ with bersonality disorders         B         0.5%         1.1%           % 65+ with bersonality disorders         B         0.5%         1.1%           % 65+ with bersonality disorders         10.0%         10.4%           % 65+ with bersonality disorders         10.0%         10.4%           % 65+ with status         W         14.7%         12.0%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with status         W         14.0%         10.8%           % 65+ with status         13.8%         13.0%         28.2%           % 65+ with status         W         14.0%         10.8%           % 65+ with status         W         14.8%         10.8%           % 65+ with conscious tructive pulmonary disease         W         40.1%         34.3%           % 65+ with status         5.4%         4.5%<	% 60+ with 15+ days poor mental health last month		8.1%	6.9%
% 65+ with bipolar disorders         B         1.7%         3.1%           % 65+ with post-traumatic stress disorder         1.3%         1.4%           % 65+ with personality disorders         4.5%         4.9%           % 65+ with personality disorders         B         0.5%         1.1%           % 65+ with substance use disorders (drug use +/or alcohol abuse)         5.8%         5.5%           % 65+ with substance use disorders         10.4%         10.4%           CHEONIC DISEASE         W         14.7%         12.0%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with diabetes         W         38.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with bronic obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with schernic heart disease         W         40.1%         34.3%           % 65+ with schernic heart disease         W         40.1%         34.3%           % 65+ with s	% 65+ with depression		30.6%	28.8%
% 65+ with post-traumatic stress disorder         1.3%         1.4%           % 65+ with schizophrenia & other psychotic disorders         4.5%         4.9%           % 65+ with personality disorders         B         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.27%           % 65+ with substance use disorders (drug use +/or alcohol abuse)         5.8%         5.5%           % 65+ with substance use disorders         10.4%         10.4%           CHRONIC DISEASE         V         36.0%         28.2%           % 65+ with Alzheimer's disease or related dementias         W         14.0%         10.8%           % 65+ with stroke         W         38.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         24.8%         20.5%           % 65+ with stroke         W         24.8%         20.5%           % 65+ with brenic heart disease         W         24.8%         20.5%           % 65+ with schemic heart disease         W         24.8%         20.5%           % 65+ with congestive heart disease         W         20.1%         36.3%           % 65+ with congestive heart disease         W         20.1%         37.7%	% 65+ with anxiety disorders		21.2%	21.9%
% 65+ with schizophrenia & other psychotic disorders         B         0.5%         1.1%           % 65+ with personality disorders         B         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.279           % 65+ with substance use disorders (drug use +/or alcohol abuse)         5.8%         5.5%           % 65+ with tobacco use disorders         10.4%         10.4%         10.4%           CHRONIC DISEASE         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with orboric obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with astma         13.8%         13.0%         65%           % 65+ with orenic obstructive pulmonary disease         W         40.1%         34.3%           % 65+ with astma         13.8%         13.0%         65%           % 65+ with congestive heart failure         18.8%         17.7%           % 65+ with orgeneration disease         W         40.1%         34.3%           % 65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with osteoarthritis/rheumatoid arthritis         1	% 65+ with bipolar disorders	В	1.7%	3.1%
% 65+ with personality disorders         B         0.5%         1.1%           # opioid deaths (all ages) (county)         37         1.279           % 65+ with substance use disorders         10.4%         10.4%           CHRONIC DISEASE         0.4%         12.0%           % 65+ with Alzheimer's disease or related dementias         W         14.7%         12.0%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with diabetes         W         14.0%         10.8%           % 65+ with asthma         13.8%         13.0%           % 65+ with asthma         13.8%         13.0%           % 65+ with asthma         13.8%         13.0%           % 65+ with congestive heart dialure         18.8%         17.7%           % 65+ with congestive heart dialure         18.8%         17.7%           % 65+ with osteoparthitis/rheumatoid arthritis         47.8%         49.1%           % 65+ with cole cancer         1.6%         2.0%           % 65+ with lucy cancer         1.5%         1.6%           % 65+ with colon cancer         1.5%	% 65+ with post-traumatic stress disorder		1.3%	1.4%
# opioid deaths (all ages) (county)         37         1.279           % 65+ with substance use disorders (drug use +/or alcohol abuse)         5.8%         5.5%           % 65+ with tobacco use disorders         10.4%         10.4%           CHRONIC DISEASE          14.7%         12.0%           % 65+ with Alzheimer's disease or related dementias         W         14.7%         12.0%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         24.8%         20.5%           % 65+ with asthma         13.8%         13.0%           % 65+ with operations         W         77.7%         70.2%           % 65+ with congestive heart disease         W         44.3%         45.9%           % 65+ with congestive heart disease         W         25.5%         14.7%           % 65+ with ober optrovisis         17.9%         17.4%           % 65+ with osteoprovisis         17.9%         17.4%           % 65+ w	% 65+ with schizophrenia & other psychotic disorders		4.5%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse)         5.8%         5.5%           % 65+ with tobacco use disorders         10.4%         10.4%           CHRONIC DISEASE         ************************************	% 65+ with personality disorders	В	0.5%	1.1%
% 65+ with tobacco use disorders         10.4%         10.4%           CHRONIC DISEASE         ************************************	# opioid deaths (all ages) (county)		37	1,279
CHRONIC DISEASE         W         14.7%         12.0%           % 65+ with Alzheimer's disease or related dementias         W         35.0%         28.2%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with asthma         13.8%         13.0%           % 65+ with ischemic heart disease         W         40.1%         34.3%           % 65+ with atrial fibrillation         15.0%         14.4%           % 65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with osteoporosis         17.9%         17.4%           % 65+ with leukemias and lymphomas         1.6%         2.0%           % 65+ with loop cancer         1.5%         1.6%           % 65+ with loop cancer         1.2%         1.7%	% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.8%	5.5%
% 65+ with Alzheimer's disease or related dementias         W         14.7%         12.0%           % 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with chronic obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with asthma         13.8%         13.0%         45.0%           % 65+ with asthma         13.8%         13.0%         45.4%           % 65+ with achemic heart disease         W         40.1%         34.3%           % 65+ with congestive heart failure         18.8%         17.7%           % 65+ with atrial fibrillation         15.0%         14.4%           % 65+ with osteoparthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with osteoparosis         17.9%         17.4%           % 65+ with leukemias and lymphomas         1.6%         2.0%           % 65+ with colon cancer         3.1%         2.4%           % 65+ with loug cancer         1.5%         1.6%           % 65+ with loug cancer         1.2%         1.7%           % 65+ with colon cancer         1.2%         1.7% <t< td=""><td>% 65+ with tobacco use disorders</td><td></td><td>10.4%</td><td>10.4%</td></t<>	% 65+ with tobacco use disorders		10.4%	10.4%
% 65+ with diabetes         W         35.0%         28.2%           % 65+ with stroke         W         14.0%         10.8%           % 65+ with chronic obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with asthma         13.8%         13.0%         65.4%           % 65+ with hypertension         W         77.7%         70.2%           % 65+ with hypertension         W         40.1%         34.3%           % 65+ with congestive heart disease         W         40.1%         34.3%           % 65+ with congestive heart failure         18.8%         17.7%           % 65+ with atrial fibrillation         15.0%         14.4%           % 65+ with peripheral vascular disease         W         25.5%         14.7%           % 65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with leukemias and lymphomas         1.6%         2.0%           % 65+ with long cancer         1.5%         1.6%           % 65+ with long cancer         1.5%         1.6%           % 65+ with long cancer         1.5%         1.6%           % 65+ with networe with prostate cancer         1.2%         1.7%           % 65+ with horedometrial cancer         1.2%         1.7% <td>CHRONIC DISEASE</td> <td></td> <td></td> <td></td>	CHRONIC DISEASE			
% 65+ with stroke         W         14.0%         10.8%           % 65+ with chronic obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with asthma         13.8%         13.0%           % 65+ with hypertension         W         77.7%         70.2%           % 65+ ever had a heart attack         5.4%         4.5%           % 65+ ever had a heart attack         5.4%         4.5%           % 65+ with ischemic heart disease         W         40.1%         34.3%           % 65+ with congestive heart failure         18.8%         17.7%           % 65+ with atrial fibrillation         15.0%         14.4%           % 65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with osteoporosis         17.9%         17.4%           % 65+ with leukemias and lymphomas         1.6%         2.0%           % 65+ women with presat cancer         1.5%         1.6%           % 65+ women with prostate cancer         1.2%         1.7%           % 65+ with hypothyroidism         20.5%         20.5%           % 65+ with hypothyroidism         20.5%         20.8%           % 65+ with hypothyroidism         20.5%         20.8%           % 65+ with hypothyroidism         20	% 65+ with Alzheimer's disease or related dementias	W	14.7%	12.0%
% 65+ with chronic obstructive pulmonary disease         W         24.8%         20.5%           % 65+ with asthma         13.8%         13.0%           % 65+ with hypertension         W         77.7%         70.2%           % 65+ ever had a heart attack         5.4%         4.5%           % 65+ ever had a heart attack         5.4%         4.5%           % 65+ with ischemic heart disease         W         40.1%         34.3%           % 65+ with congestive heart failure         18.8%         17.7%           % 65+ with attrial fibrillation         15.0%         14.4%           % 65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           % 65+ with osteoporosis         17.9%         17.4%           % 65+ with leukemias and lymphomas         1.6%         2.0%           % 65+ woren with perstate cancer         1.5%         1.6%           % 65+ woren with prostate cancer         1.2%         1.7%           % 65+ woren with prostate cancer         10.4%         11.5%           % 65+ with hilping prostate chyperplasia         B         32.3%         36.8%           % 65+ with hypothyroidism         20.5%         20.5%         20.8%           % 65+ with hypothyroidism         20.5%         20.5%         20.8% <td>% 65+ with diabetes</td> <td>W</td> <td>35.0%</td> <td>28.2%</td>	% 65+ with diabetes	W	35.0%	28.2%
% 65+ with asthma       13.8%       13.0%         % 65+ with hypertension       W       77.7%       70.2%         % 65+ ever had a heart attack       5.4%       4.5%         % 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with congestive heart failure       18.8%       17.7%         % 65+ with atrial fibrillation       15.0%       14.4%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoorthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with colon cancer       1.6%       2.0%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with hypothyroidism       20.5%       20.5%       20.8%         % 65+ with hypothyroidism       20.5%       20.5%       20.8%         % 65+ with hypothyroidism       20.5%       20.8%       20.5%       20.8%         % 65+ with chronic kidney disease       W       40.4%       37.3%         % 65+ with hypothyroidism       20.	% 65+ with stroke	W	14.0%	10.8%
% 65+ with hypertension       W       77.7%       70.2%         % 65+ ever had a heart attack       5.4%       4.5%         % 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with congestive heart failure       18.8%       17.7%         % 65+ with atrial fibrillation       15.0%       14.4%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoarthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with prostate cancer       10.4%       11.5%       1.6%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with hypothyroidism       20.5%       20.5%       20.8%         % 65+ with hypothyroidism       20.5%       20.8%       65+ with chronic kidney disease       W       40.4%       37.3%         % 65+ with hilt lever diseases <t< td=""><td>% 65+ with chronic obstructive pulmonary disease</td><td>W</td><td>24.8%</td><td>20.5%</td></t<>	% 65+ with chronic obstructive pulmonary disease	W	24.8%	20.5%
% 65+ ever had a heart attack       5.4%       4.5%         % 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with congestive heart failure       18.8%       17.7%         % 65+ with atrial fibrillation       15.0%       14.4%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoarthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with long cancer       1.5%       1.6%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with chronic kidney disease       W       40.4%       37.3%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with chronic kidney disease       8.0%       6.9%	% 65+ with asthma		13.8%	13.0%
% 65+ with ischemic heart disease       W       40.1%       34.3%         % 65+ with congestive heart failure       18.8%       17.7%         % 65+ with atrial fibrillation       15.0%       14.4%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoarthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with long cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with prostate cancer       10.4%       11.5%       1.6%         % 65+ with HIV/AIDS       0.05%       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%       20.8%         % 65+ with nemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with hypertension	W	77.7%	70.2%
% 65+ with congestive heart failure       18.8%       17.7%         % 65+ with atrial fibrillation       15.0%       14.4%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoparthritis/rheumatoid atthritis       47.8%       49.1%         % 65+ with osteoparthritis/rheumatoid atthritis       47.8%       49.1%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with lung cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       8       7.4%       9.8%         % 65+ women with endometrial cancer       10.4%       11.5%         % 65+ with hypothyroidism       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with hypothyroidism       20.5%       20.8%       20.8%         % 65+ with hypothyroidism       20.5%       20.8%       20.8%         % 65+ with hypothyroidisease       W       26.4%<	% 65+ ever had a heart attack		5.4%	4.5%
%         15.0%         14.4%           %         65+ with atrial fibrillation         15.0%         14.7%           %         65+ with peripheral vascular disease         W         25.5%         14.7%           %         65+ with osteoarthritis/rheumatoid arthritis         47.8%         49.1%           %         65+ with osteoporosis         17.9%         17.4%           %         65+ with leukemias and lymphomas         1.6%         2.0%           %         65+ with lung cancer         1.5%         1.6%           %         65+ with colon cancer         3.1%         2.4%           %         65+ women with breast cancer         8         7.4%         9.8%           %         65+ women with prostate cancer         10.4%         11.5%           %         65+ women with prostate cancer         10.4%         11.5%           %         65+ with benign prostatic hyperplasia         B         32.3%         36.8%           %         65+ with hypothyroidism         20.5%         20.8%         20.5%         20.8%           %         65+ with hypothyroidism         20.5%         20.8%         20.5%         20.8%           %         65+ with hypothyroidism         20.5%         20.8%	% 65+ with ischemic heart disease	W	40.1%	34.3%
% 65+ with peripheral vascular disease       W       25.5%       14.7%         % 65+ with osteoarthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with long cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with hypothypothyperplasia       B       32.3%       36.8%         % 65+ with hypothypoidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%	% 65+ with congestive heart failure		18.8%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis       47.8%       49.1%         % 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with lung cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%       20.5%       20.8%         % 65+ with chronic kidney disease       W       40.4%       37.3%         % 65+ with chronic kidney disease       8.0%       6.9%	% 65+ with atrial fibrillation		15.0%	14.4%
% 65+ with osteoporosis       17.9%       17.4%         % 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with lung cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with namenia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with peripheral vascular disease	W	25.5%	14.7%
% 65+ with leukemias and lymphomas       1.6%       2.0%         % 65+ with lung cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ women with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with nemia       W       40.4%       37.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with osteoarthritis/rheumatoid arthritis		47.8%	49.1%
% 65+ with lung cancer       1.5%       1.6%         % 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ men with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with liver disease       W       26.4%       22.3%	% 65+ with osteoporosis		17.9%	17.4%
% 65+ with colon cancer       3.1%       2.4%         % 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ men with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with leukemias and lymphomas		1.6%	2.0%
% 65+ women with breast cancer       B       7.4%       9.8%         % 65+ women with endometrial cancer       1.2%       1.7%         % 65+ men with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with lung cancer		1.5%	1.6%
% 65+ women with endometrial cancer       1.2%       1.7%         % 65+ men with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with colon cancer		3.1%	2.4%
% 65+ men with prostate cancer       10.4%       11.5%         % 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ women with breast cancer	В	7.4%	9.8%
% 65+ with benign prostatic hyperplasia       B       32.3%       36.8%         % 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ women with endometrial cancer		1.2%	1.7%
% 65+ with HIV/AIDS       0.05%       0.05%         % 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ men with prostate cancer		10.4%	11.5%
% 65+ with hypothyroidism       20.5%       20.8%         % 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with benign prostatic hyperplasia	В	32.3%	36.8%
% 65+ with anemia       W       40.4%       37.3%         % 65+ with chronic kidney disease       W       26.4%       22.3%         % 65+ with liver diseases       8.0%       6.9%	% 65+ with HIV/AIDS		0.05%	0.05%
% 65+ with chronic kidney disease         W         26.4%         22.3%           % 65+ with liver diseases         8.0%         6.9%	% 65+ with hypothyroidism		20.5%	20.8%
% 65+ with liver diseases 8.0% 6.9%	% 65+ with anemia	W	40.4%	37.3%
	% 65+ with chronic kidney disease	W	26.4%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue W 24.3% 18.6%	% 65+ with liver diseases		8.0%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	W	24.3%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache	В	3.0%	4.0%
% 65+ with epilepsy		2.5%	2.1%
% 65+ with traumatic brain injury	В	0.6%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma		22.9%	22.9%
% 65+ with cataract	W	65.8%	61.2%
% 65+ with pressure ulcer or chronic ulcer		6.3%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	W	61.8%	54.4%
% 65+ with 0 chronic conditions	W	7.4%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		17.6%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment	W	17.4%	14.4%
% 65+ with self-reported vision difficulty	В	2.0%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment	В	0.4%	0.9%
% 65+ with self-reported cognition difficulty		5.2%	6.9%
% 65+ with self-reported ambulatory difficulty		20.7%	18.8%
% 65+ with clinical diagnosis of mobility impairments		3.8%	3.2%
% 65+ with self-reported self-care difficulty		6.9%	5.6%
% 65+ with self-reported independent living difficulty		13.1%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees	*	1.2%	7.9%
% dually eligible for Medicare and Medicaid	*	16.9%	7.5%
% 60+ with a regular doctor		93.8%	95.8%
% 60+ who did not see doctor when needed due to cost		8.2%	5.4%
# of primary care providers within 5 miles		28	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		2	74
# of home health agencies		2	49
# of community health centers		3	22
# of adult day health centers		1	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	1.5	6.3
Emergency room visits/1000 persons 65+ years per year	*	781	572
Part D monthly prescription fills per person per year	*	61.4	49.1
Home health visits per year		2.6	2.5
Durable medical equipment claims per year	*	2.8	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	300	237
Medicare inpatient hospital readmissions (as % of admissions)		14.8%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year	*	126	76
# skilled nursing home Medicare beds/1000 persons 65+ years		90	33
% 65+ getting Medicaid long term services and supports	*	8.9%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		1	44
Air pollution: annual # of unhealthy days for 65+ (county)		3	N/A
% of grandparents raising grandchildren		0.5%	0.8%
% of grandparents who live with grandchildren		1.6%	2.5%
# of assisted living sites		0	134
% of vacant homes in community		16.9%	16.0%
# of universities and community colleges		1	41
# of public libraries		1	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		70.0%	93.0%
% 60+ who used Internet in last month		61.3%	77.6%
Voter participation rate in 2018 election (age 18+)		33.8%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		253	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		31	586
Property crime rate /100,000 persons		1,749	2,012
% 65+ who own a motor vehicle		86.3%	91.0%
% 60+ who always drive wearing a seatbelt	W	60.7%	77.1%
# of fatal crashes involving adult age 60+/town		0	151
# of fatal crashes involving adult age 60+/county		11	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		10.4%	5.4%
% 60+ receiving food stamps past year		11.0%	5.7%
% 65+ employed past year		14.1%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		33.9%	18.2%
% households with annual income \$20,000-\$49,999		47.9%	36.5%
% households with annual income > \$50,000		18.2%	45.3%
% 60+ own home		72.3%	79.9%
% 60+ have mortgage on home		16.8%	35.3%
% 65+ households spend >35% of income on housing (renter)		9.5%	8.7%
% 65+ households spend >35% of income on housing (owner)		21.8%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$23,496	\$25,284	0.93
Single, renter, good health	\$23,100	\$26,400	0.88
Couple, homeowner without mortgage, good health	\$35,436	\$37,128	0.95
Couple, renter, good health	\$35,040	\$38,244	0.92

## TECHNICAL NOTES

\*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an \*. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

## Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019