Windham (Rockingham)

Windham is an urban town in Rockingham County that hosts an annual Strawberry Festival featuring food from local businesses. There are 1,902 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of hip fracture, depression, anxiety and bipolar disorders, post-traumatic stress disorder, personality disorders, substance and tobacco use disorders, Alzheimer's disease or related dementias, chronic obstructive pulmonary disease, asthma, peripheral vascular disease, benign prostatic hyperplasia, fibromyalgia/chronic pain/fatigue, epilepsy, and pressure ulcers. However, they had a higher rate of high cholesterol. They are more likely to take the health promotion steps of having regular mammograms and annual dental exams. Community resources to support healthy aging include 70 primary care providers and a nursing home within 5 miles, 13 home health agencies, a senior center, 5 assisted living sites, a public library, and access to broadband.

| POPULATION CHARACTERISTICS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|--|---|-----------------------|-------------------|
| Total population all ages | STATE RATE | 14,238 | 1,327,503 |
| Population 60 years or older as % of total population | | 21.0% | 22.7% |
| Total population 60 years or older | | 2,994 | 301,630 |
| | | 13.4% | |
| Population 65 years or older as % of total population | | | 15.8% |
| Total population 65 years or older | | 1,902 | 210,385 |
| % 65-74 years | | 60.6% | 58.5% |
| % 75-84 years | | 28.1% | 28.6% |
| % 85 years or older | | 11.3% | 12.9% |
| Gender (65+ population) | | | |
| % female | | 48.0% | 54.7% |
| Race/Ethnicity (65+ population) | | | |
| % White | | 97.8% | 97.7% |
| % African American | | 0.0% | 0.5% |
| % Asian | | 1.8% | 0.9% |
| % Other | | 0.4% | 0.9% |
| % Hispanic/Latino | | 0.0% | 0.9% |
| Marital Status (65+ population) | | | |
| % married | | 62.1% | 58.5% |
| % divorced/separated | | 14.3% | 14.0% |
| % widowed | | 22.9% | 22.9% |
| % never married | | 0.7% | 4.6% |
| Education (65+ population) | | | |
| % with less than high school education | | 12.6% | 12.3% |
| % with high school or some college | | 52.1% | 57.1% |
| % with college degree | | 35.3% | 30.6% |
| % of 65+ population living alone | | 18.7% | 26.1% |
| % of 65+ population who speak only English at home | | 94.1% | 91.3% |
| % of 65+ population who are veterans of military service | | 30.7% | 24.8% |
| Age-sex adjusted 1-year mortality rate | | 4.3% | 4.1% |





| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|--|---|-----------------------|-------------------|
| Geographic Migration (65+ population) in the past 12 months | | | |
| % moved within same county | | 1.0% | 3.6% |
| % moved from different county in New Hampshire | | 0.0% | 1.0% |
| % moved from different state | | 3.5% | 1.7% |
| WELLNESS & PREVENTION | | | |
| % 60+ with any physical activity within last month | | 75.1% | 74.5% |
| % 60+ met CDC guidelines for muscle-strengthening activity | | 28.5% | 26.0% |
| % 60+ met CDC guidelines for aerobic physical activity | | 61.7% | 58.6% |
| % 60+ met CDC guidelines for both types of physical activities | | 20.5% | 19.9% |
| % 60+ getting recommended hours of sleep | | 64.8% | 66.4% |
| % 60+ injured in a fall within last 12 months | | 9.5% | 10.4% |
| % 65+ had hip fracture | В | 2.5% | 3.3% |
| % 60+ with self-reported fair or poor health status | | 14.3% | 16.5% |
| % 60+ with 15+ physically unhealthy days last month | | 9.5% | 12.5% |
| % 60+ with physical exam/check-up in past year | | 90.4% | 86.5% |
| % 60+ met CDC preventive health screening goals | | 46.3% | 40.3% |
| % 60+ flu shot past year | | 56.8% | 59.3% |
| % 65+ with pneumonia vaccine | | 74.7% | 77.8% |
| % 60+ with cholesterol screening | | 97.6% | 95.3% |
| % 60+ women with a mammogram within last 2 years | В | 93.4% | 79.3% |
| % 60+ with colorectal cancer screening | | 81.1% | 77.0% |
| % 60+ with HIV test | | 9.9% | 13.5% |
| % 60+ current smokers | | 7.9% | 8.1% |
| Oral Health | | | |
| % 60+ with loss of 6 or more teeth | | 25.4% | 29.0% |
| % 60+ with annual dental exam | В | 82.6% | 75.7% |
| # of dentists per 100,000 persons (all ages) (county) | | 72 | 72 |
| NUTRITION/DIET | | | |
| % 60+ with 5 or more servings of fruit or vegetables per day | | 20.3% | 20.2% |
| % 60+ self-reported obese | | 32.0% | 27.2% |
| % 65+ clinically diagnosed obese | | 17.2% | 16.7% |
| % 65+ with high cholesterol | W | 75.0% | 72.2% |
| % 60+ excessive drinking | | 11.2% | 9.2% |
| % 65+ with poor supermarket access | | 77.1% | 28.4% |

| % 65+ with depression B 24.2% 28.89 % 65+ with anxiety disorders B 19.0% 21.99 % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 0.6% 1.19 # opioid deaths (all ages) (county) 276 1.27 % 65+ with schizophrenia & other psychotic disorders B 7.1% 10.49 CHRONC DISEASE B 7.1% 10.49 Ø 65+ with diabetes 28.4% 28.29 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 9.5% 10.89 % 65+ with hypertension 70.7% 70.29 % 65+ with stroke 3.7% 4.55 % 65+ with chronic obstructive pulmonary disease B 11.0% % 65+ with discherat failure 16.5% 17.7% % 65+ with otherangestrain 3.7% 4.59 | HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|--|--|---|-----------------------|-------------------|
| % 65+ with depression B 24.2% 28.89 % 65+ with anxiety disorders B 19.0% 21.99 % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 0.6% 1.19 # opioid deaths (all ages) (county) 276 1.27 % 65+ with schizophrenia & other psychotic disorders B 7.1% 10.49 CHRONC DISEASE B 7.1% 10.49 CHRONC DISEASE 28.4% 28.27 % 65+ with diabetes 28.4% 28.27 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 3.7% 4.55 % 65+ with stroke 3.7% <t< td=""><td>BEHAVIORAL HEALTH</td><td></td><td></td><td></td></t<> | BEHAVIORAL HEALTH | | | |
| % 65+ with anxiety disorders B 19.0% 21.99 % 65+ with bolar disorders B 2.3% 3.19 % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with bersonality disorders B 0.7% 1.47 % 65+ with bersonality disorders B 0.6% 1.11 # opioid deaths (all ages) (county) 276 1.27 % 65+ with bersonality disorders B 0.6% 1.19 % 65+ with backnec use disorders (drug use +/or alcohol abuse) B 4.1% 5.55 % 65+ with Alzheimer's disease or related dementias B 9.9% 12.09 % 65+ with Alzheimer's disease or related dementias B 9.5% 10.49 CHRONC DISEASE 28.4% 28.29 26.4% 28.29 % 65+ with Alzheimer's disease B 11.0% 12.09 % 65+ with strike 9.5% 10.89 2.5% % 65+ with strike 9.5% 10.89 2.5% % 65+ with strike 3.7% 4.56 % 65+ with strike <td< td=""><td>% 60+ with 15+ days poor mental health last month</td><td></td><td>5.3%</td><td>6.9%</td></td<> | % 60+ with 15+ days poor mental health last month | | 5.3% | 6.9% |
| % 65+ with bipolar disorders B 2.3% 3.19 % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 3.9% 4.99 % 65+ with personality disorders B 0.6% 1.19 % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.55 % 65+ with substance use disorders B 7.1% 10.49 CHRONIC DISEASE | % 65+ with depression | В | 24.2% | 28.8% |
| % 65+ with post-traumatic stress disorder B 0.7% 1.49 % 65+ with schizophrenia & other psychotic disorders B 3.9% 4.99 % 65+ with personality disorders B 0.6% 1.19 # opioid deaths (all ages) (county) 276 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.59 % 65+ with tobacco use disorders B 7.1% 10.49 CHRONIC DISEASE 28.4% 28.29 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 9.5% 10.89 % 65+ with stroke 3.7% 4.59 % 65+ with astma B 11.0% 13.00 % 65+ with chonic beatru disease 32.0% 34.39 % 65+ with astima 11.9% 14.47 % 65+ with origonsith fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with o | % 65+ with anxiety disorders | В | 19.0% | 21.9% |
| % 65+ with schizophrenia & other psychotic disorders B 3.9% 4.99 % 65+ with personality disorders B 0.6% 1.19 # opioid deaths (all ages) (county) 276 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.59 % 65+ with tobacco use disorders B 7.1% 10.49 % 65+ with diabetes 28.4% 28.29 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with obstructive pulmonary disease B 16.5% 20.59 % 65+ with stroke 3.7% 4.59 % 65+ with astma B 11.0% 13.09 % 65+ with orgenic heart disease 32.0% 34.39 % 65+ with orgenic heart failure 16.5% 17.79 % 65+ with orgenic heart failure 16.5% 17.79 % 65+ with orgenic heart failure 16.5% 17.79 % 65+ with osteoarthrifis/heumatoid arthritis 47.9% 49.19 % 65+ with osteoarthrifis/heumatoid arthritis 47.9% | % 65+ with bipolar disorders | В | 2.3% | 3.1% |
| % 65+ with personality disorders B 0.6% 1.19 # opioid deaths (all ages) (county) 276 1.27 % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.59 % 65+ with tobacco use disorders B 7.1% 10.44 CHRONIC DISEASE 28.4% 28.29 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with asthma B 11.0% 13.09 % 65+ with asthma B 11.0% 13.09 % 65+ with operstive heart failure 10.5% 17.79 % 65+ with congestive heart failure 10.5% 17.79 % 65+ with atrial librillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 | % 65+ with post-traumatic stress disorder | В | 0.7% | 1.4% |
| # opioid deaths (all ages) (county) 276 1,27 % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.59 % 65+ with tobacco use disorders B 7.1% 10.49 CHRONIC DISEASE | % 65+ with schizophrenia & other psychotic disorders | В | 3.9% | 4.9% |
| % 65+ with substance use disorders (drug use +/or alcohol abuse) B 4.1% 5.5% % 65+ with tobacco use disorders B 7.1% 10.49 CHRONIC DISEASE | % 65+ with personality disorders | В | 0.6% | 1.1% |
| % 65+ with tobacco use disorders B 7.1% 10.49 CHRONIC DISEASE 28.4% 28.29 % 65+ with Alzheimer's disease or related dementias B 9.9% 12.09 % 65+ with Alzheimer's disease or related dementias B 9.9% 12.09 % 65+ with diabetes 28.4% 28.22 28.29 % 65+ with stroke 9.5% 10.89 20.59 % 65+ with stroke B 11.0% 13.09 % 65+ with hypertension 70.7% 70.22 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.7% % 65+ with congestive heart disease B 11.9% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with colon cancer 2.2% 2.0% % 65+ with lung cancer 1.5% 1.69 % 65+ with lung cancer 1.5% 1.69 % 65+ with lung cancer 1.5% 1.69 < | # opioid deaths (all ages) (county) | | 276 | 1,279 |
| CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.9% 12.09 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with chronic obstructive pulmonary disease B 16.5% 20.59 % 65+ with asthma B 11.0% 13.09 % 65+ with stroke 3.70% 70.29 % 65+ with asthma B 11.0% 13.09 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with tarial fibrillation 13.4% 14.49 % 65+ with tarial fibrillation 13.4% 14.49 % 65+ with osteoporosis 16.9% 17.49 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with colon cancer 2.7% 2.49 % 65+ with colon cancer 1.5% 1.69 % 65+ with prostate cancer 12.3% 11.59 % 65+ with prostate cancer </td <td>% 65+ with substance use disorders (drug use +/or alcohol abuse)</td> <td>В</td> <td>4.1%</td> <td>5.5%</td> | % 65+ with substance use disorders (drug use +/or alcohol abuse) | В | 4.1% | 5.5% |
| % 65+ with Alzheimer's disease or related dementias B 9.9% 12.00 % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with chronic obstructive pulmonary disease B 16.5% 20.59 % 65+ with asthma B 11.0% 13.09 % 65+ with hypertension 70.7% 70.29 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.77 % 65+ with distoenci heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoporosis 16.9% 17.49 % 65+ with osteoporosis 16.9% 17.49 % 65+ with loug cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.4% % 65+ worne with prostate cancer 10.1% 9.88 % 65+ worne with prostate cancer 12.3% 11.5% % 65+ wornen with prostate cancer 12.3% | % 65+ with tobacco use disorders | В | 7.1% | 10.4% |
| % 65+ with diabetes 28.4% 28.29 % 65+ with stroke 9.5% 10.89 % 65+ with chronic obstructive pulmonary disease B 16.5% 20.59 % 65+ with asthma B 11.0% 13.09 % 65+ with hypertension 70.7% 70.29 % 65+ with hypertension 70.7% 70.29 % 65+ with ischemic heart disease 32.0% 34.33 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with losteoprosis 16.9% 17.49 % 65+ with losteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with losteoprosis 16.9% 17.49 % 65+ with losteoprosis 16.9% 17.49 % 65+ with losteoprosis 16.9% 17.49 % 65+ with losteon cancer 2.2% 2.09 % 65+ with losteoarcancer 1.5% 1.69 | CHRONIC DISEASE | | | |
| % 65+ with stroke 9.5% 10.8% % 65+ with chronic obstructive pulmonary disease B 16.5% 20.5% % 65+ with chronic obstructive pulmonary disease B 11.0% 13.0% % 65+ with hypertension 70.7% 70.2% % 65+ ever had a heart attack 3.7% 4.55 % 65+ with ischemic heart disease 32.0% 34.33 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with peripheral vascular disease B 11.9% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with loug cancer 1.5% 1.69% % 65+ women with prostate cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.5% % 65+ women with prostate cancer 12.3% 11.5% % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 22.3% % 65+ with hypothyroidism </td <td>% 65+ with Alzheimer's disease or related dementias</td> <td>В</td> <td>9.9%</td> <td>12.0%</td> | % 65+ with Alzheimer's disease or related dementias | В | 9.9% | 12.0% |
| % 65+ with chronic obstructive pulmonary disease B 16.5% 20.59 % 65+ with asthma B 11.0% 13.00 % 65+ with hypertension 70.7% 70.29 % 65+ ever had a heart attack 3.7% 4.59 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with peripheral vascular disease B 11.9% 14.49 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.5% % 65+ women with prostate cancer 12.3% 11.5% % 65+ women with prostate cancer 12.3% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism </td <td>% 65+ with diabetes</td> <td></td> <td>28.4%</td> <td>28.2%</td> | % 65+ with diabetes | | 28.4% | 28.2% |
| % 65+ with ashma B 11.0% 13.09 % 65+ with hypertension 70.7% 70.29 % 65+ ever had a heart attack 3.7% 4.59 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with colon cancer 2.7% 2.49 % 65+ women with prostate cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.5% % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 36.89 % 65+ with hypothyroidism 22.3% 20.89 36.89 % 65+ with hypothyroidism 22.3% 23.89 36.89 36.8 | % 65+ with stroke | | 9.5% | 10.8% |
| % 65+ with hypertension 70.7% 70.29 % 65+ ever had a heart attack 3.7% 4.59 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.0% % 65+ with colon cancer 1.5% 1.69 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.5% % 65+ with being prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 36.89 36.89 36.89 36.89 36.89 36.89 | % 65+ with chronic obstructive pulmonary disease | В | 16.5% | 20.5% |
| % 65+ ever had a heart attack 3.7% 4.59 % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with congestive heart failure 13.4% 14.49 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with loukemias and lymphomas 2.2% 2.09 % 65+ with loug cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.59 % 65+ women with prostate cancer 12.3% 11.59 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with anemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% </td <td>% 65+ with asthma</td> <td>В</td> <td>11.0%</td> <td>13.0%</td> | % 65+ with asthma | В | 11.0% | 13.0% |
| % 65+ with ischemic heart disease 32.0% 34.39 % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with leukemias and lymphomas 2.2% 2.0% % 65+ with loug cancer 1.5% 1.6% % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.5% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 22.3% 20.89 % 65+ with anemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver d | % 65+ with hypertension | | 70.7% | 70.2% |
| % 65+ with congestive heart failure 16.5% 17.79 % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with long cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ women with prostate cancer 12.3% 11.5% % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 22.3% | % 65+ ever had a heart attack | | 3.7% | 4.5% |
| % 65+ with atrial fibrillation 13.4% 14.49 % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with lung cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ with benign prostatic cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with ischemic heart disease | | 32.0% | 34.3% |
| % 65+ with peripheral vascular disease B 11.9% 14.79 % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with long cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ women with prostate cancer 12.3% 11.59 % 65+ with hypothyroidism B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with anemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.9% % 65+ with chronic kidney disease 6.9% 6.9% | % 65+ with congestive heart failure | | 16.5% | 17.7% |
| % 65+ with osteoarthritis/rheumatoid arthritis 47.9% 49.19 % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with lung cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ women with prostate cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with hypothyroidism 22.3% 20.89 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with atrial fibrillation | | 13.4% | 14.4% |
| % 65+ with osteoporosis 16.9% 17.49 % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with lung cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 10.1% 9.89 % 65+ women with prostate cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with nemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with peripheral vascular disease | В | 11.9% | 14.7% |
| % 65+ with leukemias and lymphomas 2.2% 2.09 % 65+ with lung cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ men with prostate cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with namenia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with osteoarthritis/rheumatoid arthritis | | 47.9% | 49.1% |
| % 65+ with lung cancer 1.5% 1.69 % 65+ with colon cancer 2.7% 2.49 % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ men with prostate cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with osteoporosis | | 16.9% | 17.4% |
| % 65+ with colon cancer 2.7% 2.4% % 65+ women with breast cancer 10.1% 9.8% % 65+ women with endometrial cancer 1.8% 1.7% % 65+ men with prostate cancer 12.3% 11.5% % 65+ with benign prostatic hyperplasia B 32.8% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 22.3% 20.8% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ with leukemias and lymphomas | | 2.2% | 2.0% |
| % 65+ women with breast cancer 10.1% 9.89 % 65+ women with endometrial cancer 1.8% 1.79 % 65+ men with prostate cancer 12.3% 11.59 % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with anemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ with lung cancer | | 1.5% | 1.6% |
| % 65+ women with endometrial cancer 1.8% 1.7% % 65+ men with prostate cancer 12.3% 11.5% % 65+ with benign prostatic hyperplasia B 32.8% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 22.3% 20.8% % 65+ with anemia 37.0% 37.3% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ with colon cancer | | 2.7% | 2.4% |
| % 65+ men with prostate cancer 12.3% 11.5% % 65+ with benign prostatic hyperplasia B 32.8% 36.8% % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 22.3% 20.8% % 65+ with anemia 37.0% 37.3% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ women with breast cancer | | 10.1% | 9.8% |
| % 65+ with benign prostatic hyperplasia B 32.8% 36.89 % 65+ with HIV/AIDS 0.04% 0.059 % 65+ with hypothyroidism 22.3% 20.89 % 65+ with anemia 37.0% 37.39 % 65+ with chronic kidney disease 22.9% 22.39 % 65+ with liver diseases 6.9% 6.9% | % 65+ women with endometrial cancer | | 1.8% | 1.7% |
| % 65+ with HIV/AIDS 0.04% 0.05% % 65+ with hypothyroidism 22.3% 20.8% % 65+ with anemia 37.0% 37.3% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ men with prostate cancer | | 12.3% | 11.5% |
| % 65+ with hypothyroidism 22.3% 20.8% % 65+ with anemia 37.0% 37.3% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ with benign prostatic hyperplasia | В | 32.8% | 36.8% |
| % 65+ with anemia 37.0% 37.3% % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ with HIV/AIDS | | 0.04% | 0.05% |
| % 65+ with chronic kidney disease 22.9% 22.3% % 65+ with liver diseases 6.9% 6.9% | % 65+ with hypothyroidism | | 22.3% | 20.8% |
| % 65+ with liver diseases 6.9% 6.9% | % 65+ with anemia | | 37.0% | 37.3% |
| | % 65+ with chronic kidney disease | | 22.9% | 22.3% |
| % 65+ with fibromyalgia, chronic pain and fatigue B 15.4% 18.6% | % 65+ with liver diseases | | 6.9% | 6.9% |
| | % 65+ with fibromyalgia, chronic pain and fatigue | В | 15.4% | 18.6% |

| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|---|---|-----------------------|-------------------|
| % 65+ with migraine and other chronic headache | | 4.0% | 4.0% |
| % 65+ with epilepsy | В | 1.5% | 2.1% |
| % 65+ with traumatic brain injury | | 1.1% | 1.1% |
| % 65+ with autism spectrum disorders | | N/A | 0.03% |
| % 65+ with glaucoma | | 23.9% | 22.9% |
| % 65+ with cataract | | 61.9% | 61.2% |
| % 65+ with pressure ulcer or chronic ulcer | В | 5.5% | 7.1% |
| % 65+ with 4+ (out of 15) chronic conditions | В | 51.4% | 54.4% |
| % 65+ with 0 chronic conditions | | 9.4% | 10.3% |
| LIVING WITH DISABILITY | | | |
| % 65+ with self-reported hearing difficulty | В | 9.7% | 15.0% |
| % 65+ with clinical diagnosis of deafness or hearing impairment | | 13.7% | 14.4% |
| % 65+ with self-reported vision difficulty | | 4.1% | 5.2% |
| % 65+ with clinical diagnosis of blindness or visual impairment | | 0.7% | 0.9% |
| % 65+ with self-reported cognition difficulty | | 5.7% | 6.9% |
| % 65+ with self-reported ambulatory difficulty | В | 10.3% | 18.8% |
| % 65+ with clinical diagnosis of mobility impairments | | 2.6% | 3.2% |
| % 65+ with self-reported self-care difficulty | | 4.1% | 5.6% |
| % 65+ with self-reported independent living difficulty | В | 5.3% | 11.3% |
| ACCESS TO CARE | | | |
| Medicare (65+ population) | | | |
| % Medicare managed care enrollees | * | 10.2% | 7.9% |
| % dually eligible for Medicare and Medicaid | * | 3.5% | 7.5% |
| % 60+ with a regular doctor | В | 99.2% | 95.8% |
| % 60+ who did not see doctor when needed due to cost | | 11.0% | 5.4% |
| # of primary care providers within 5 miles | | 70 | 2,961 |
| # of hospitals within 5 miles | | 0 | 26 |
| # of nursing homes within 5 miles | | 1 | 74 |
| # of home health agencies | | 13 | 49 |
| # of community health centers | | 0 | 22 |
| # of adult day health centers | | 0 | 21 |
| # of memory cafes | | 0 | 12 |
| | | 0 | 14 |

| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|--|---|-----------------------|-------------------|
| SERVICE UTILIZATION | | | |
| Physician visits per year | * | 7.4 | 6.3 |
| Emergency room visits/1000 persons 65+ years per year | * | 376 | 572 |
| Part D monthly prescription fills per person per year | * | 45.1 | 49.1 |
| Home health visits per year | * | 1.9 | 2.5 |
| Durable medical equipment claims per year | | 1.8 | 2.0 |
| Inpatient hospital stays/1000 persons 65+ years per year | | 209 | 237 |
| Medicare inpatient hospital readmissions (as % of admissions) | | 16.7% | 15.8% |
| # skilled nursing facility stays/1000 persons 65+ years per year | | 59 | 76 |
| # skilled nursing home Medicare beds/1000 persons 65+ years | | 13 | 33 |
| % 65+ getting Medicaid long term services and supports | * | 2.7% | 3.7% |
| COMMUNITY VARIABLES & CIVIC ENGAGEMENT | | | |
| AARP Age-Friendly efforts in community | | Not yet | Yes |
| # of senior centers | | 1 | 44 |
| Air pollution: annual # of unhealthy days for 65+ (county) | | 2 | N/A |
| % of grandparents raising grandchildren | | 1.0% | 0.8% |
| % of grandparents who live with grandchildren | | 3.5% | 2.5% |
| # of assisted living sites | | 5 | 134 |
| % of vacant homes in community | | 4.0% | 16.0% |
| # of universities and community colleges | | 0 | 41 |
| # of public libraries | | 1 | 234 |
| # of YMCAs | | 0 | 12 |
| % in county with access to broadband (all ages) | | 99.0% | 93.0% |
| % 60+ who used Internet in last month | | 80.1% | 77.6% |
| Voter participation rate in 2018 election (age 18+) | | 63.6% | 54.7% |
| SAFETY & TRANSPORTATION | | | |
| Violent crime rate /100,000 persons | | 59 | 207 |
| Homicide rate /100,000 persons (county) | | 1 | 1 |
| # firearm fatalities (county) | | 92 | 586 |
| Property crime rate /100,000 persons | | 716 | 2,012 |
| % 65+ who own a motor vehicle | | 93.1% | 91.0% |
| % 60+ who always drive wearing a seatbelt | | 78.4% | 77.1% |
| # of fatal crashes involving adult age 60+/town | | 2 | 151 |
| # of fatal crashes involving adult age 60+/county | | 31 | 151 |

| HEALTHY AGING INDICATORS | BETTER / WORSE STATE RATE ¹ | COMMUNITY ESTIMATE | STATE ESTIMATE |
|---|---|-----------------------|-------------------------|
| ECONOMIC & HOUSING VARIABLES | | | |
| % 65+ with income below the poverty line past year | | 4.8% | 5.4% |
| % 60+ receiving food stamps past year | | 2.5% | 5.7% |
| % 65+ employed past year | | 30.8% | 24.8% |
| Household income (65+ householder) | | | |
| % households with annual income < \$20,000 | | 9.3% | 18.2% |
| % households with annual income \$20,000-\$49,999 | | 24.1% | 36.5% |
| % households with annual income > \$50,000 | | 66.6% | 45.3% |
| % 60+ own home | | 92.7% | 79.9% |
| % 60+ have mortgage on home | | 48.9% | 35.3% |
| % 65+ households spend >35% of income on housing (renter) | | 4.1% | 8.7% |
| % 65+ households spend >35% of income on housing (owner) | | 26.7% | 21.2% |
| COST OF LIVING | \$ COUNTY ESTIMATE | \$ STATE ESTIMATE | RATIO (COUNTY/STATE) |
| Elder Economic Security Standard Index | | | |
| Single, homeowner without mortgage, good health | \$25,536 | \$25,284 | 1.01 |
| Single, renter, good health | \$26,712 | \$26,400 | 1.01 |
| Couple, homeowner without mortgage, good health | \$37,092 | \$37,128 | 1.00 |
| Couple, renter, good health | \$38,268 | \$38,244 | 1.00 |

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019