

Concord (Merrimack)

Concord is the capital city of New Hampshire, located in Merrimack County and home to the gold-domed State House. There are 6,387 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of ischemic heart disease, leukemias and lymphomas, liver diseases. However, they had higher rates of hip fracture, obesity, high cholesterol, depression, anxiety/bipolar/personality disorders, schizophrenia, substance/tobacco use disorders, Alzheimer's disease, stroke, peripheral vascular disease, arthritis, osteoporosis, breast cancer, benign prostatic hyperplasia, hypothyroidism, chronic kidney disease, fibromyalgia, migraine, epilepsy, traumatic brain injury, glaucoma, cataracts, ulcers, hearing/visual/mobility impairments. They are more likely to take the health promotion step of having a regular doctor. Community resources to support healthy aging include 3 senior centers, 163 primary care providers, a hospital, and 4 nursing homes within 5 miles, 8 home health agencies, a memory cafe, a dementia support group, 3 assisted living sites, 3 universities or community colleges, 2 public libraries, a YMCA, and access to broadband.



POPULATION CHARACTERISTICS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Total population all ages		42,634	1,327,503
Population 60 years or older as % of total population		21.0%	22.7%
Total population 60 years or older		8,970	301,630
Population 65 years or older as % of total population		15.0%	15.8%
Total population 65 years or older		6,387	210,385
% 65-74 years		51.2%	58.5%
% 75-84 years		26.9%	28.6%
% 85 years or older		21.9%	12.9%
Gender (65+ population)			
% female		59.7%	54.7%
Race/Ethnicity (65+ population)			
% White		97.1%	97.7%
% African American		0.4%	0.5%
% Asian		0.9%	0.9%
% Other		1.7%	0.9%
% Hispanic/Latino		0.5%	0.9%
Marital Status (65+ population)			
% married		43.0%	58.5%
% divorced/separated		21.4%	14.0%
% widowed		28.0%	22.9%
% never married		7.6%	4.6%
Education (65+ population)			
% with less than high school education		10.0%	12.3%
% with high school or some college		57.5%	57.1%
% with college degree		32.5%	30.6%
% of 65+ population living alone		38.5%	26.1%
% of 65+ population who speak only English at home		96.4%	91.3%
% of 65+ population who are veterans of military service		25.9%	24.8%
Age-sex adjusted 1-year mortality rate	W	4.8%	4.1%
0 1/44 : 1)			DA 05 4

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		4.7%	3.6%
% moved from different county in New Hampshire		0.3%	1.0%
% moved from different state		2.3%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		74.5%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		29.2%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		61.8%	58.6%
% 60+ met CDC guidelines for both types of physical activities		22.9%	19.9%
% 60+ getting recommended hours of sleep		67.1%	66.4%
% 60+ injured in a fall within last 12 months		14.0%	10.4%
% 65+ had hip fracture	W	4.5%	3.3%
% 60+ with self-reported fair or poor health status		16.0%	16.5%
% 60+ with 15+ physically unhealthy days last month		12.8%	12.5%
% 60+ with physical exam/check-up in past year		88.8%	86.5%
% 60+ met CDC preventive health screening goals		41.4%	40.3%
% 60+ flu shot past year		60.3%	59.3%
% 65+ with pneumonia vaccine		76.9%	77.8%
% 60+ with cholesterol screening		96.9%	95.3%
% 60+ women with a mammogram within last 2 years		76.1%	79.3%
% 60+ with colorectal cancer screening		77.8%	77.0%
% 60+ with HIV test		11.9%	13.5%
% 60+ current smokers		8.0%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		24.2%	29.0%
% 60+ with annual dental exam		78.5%	75.7%
# of dentists per 100,000 persons (all ages) (county)		78	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		19.4%	20.2%
% 60+ self-reported obese		28.9%	27.2%
% 65+ clinically diagnosed obese	W	18.3%	16.7%
% 65+ with high cholesterol	W	74.7%	72.2%
% 60+ excessive drinking		6.6%	9.2%
% 65+ with poor supermarket access		36.4%	28.4%

BEHAVIORAL HEALTH 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.98% 6.98% 6.9% 6.98% 6.98% 6.9% 6.9% 6.88% 6.9% 6.9% 6.9% 6.9% 6.19% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 4.9% 6.9% 6.9% 6.9% 4.9% 6.9% 6.9% 4.9% 6.9% 6.9% 4.9% 6.9% 6.9% 4.9% 6.9% 6.9% 6.9% 4.9% 6.9% 6.9% 4.9% 6.9% 6.9% 6.9% 1.1% 4.9% 6.9% 6.9% 6.1% 1.1% 4.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.9% 6.	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with depression W 35.4% 28.8% % 65+ with anxiety disorders W 28.8% 21.9% % 65+ with bipolar disorders W 5.5% 3.1% % 65+ with post-traumatic stress disorder W 2.1% 4.1% % 65+ with post-traumatic stress disorders W 2.5% 4.9% % 65+ with post-traumatic stress disorders W 2.7% 1.1% % 65+ with schizophrenia & other psychotic disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% % 65+ with bobacco use disorders W 12.2% 10.4% CHRONIC DISEASE W 17.0% 12.0% K 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% K 65+ with disabetes 29.6% 28.2% K 65+ with stroben W 12.3% 10.8% K 65+ with stroben W 12.3% 10.0% K 65+ with hypertension <	BEHAVIORAL HEALTH			
% 65+ with pipolar disorders W 28.8% 21.9% % 65+ with pipolar disorders W 5.5% 3.1% % 65+ with pipolar disorders W 2.1% 1.4% % 65+ with personality disorders W 2.7% 1.1% % 65+ with personality disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1.279 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% % 65+ with stubstance use disorders W 12.0% 5.5% % 65+ with tobacco use disorders W 17.0% 12.0% 65+ with broke cuse disorders W 17.0% 12.0% 65+ with disbetes 29.6% 28.2% 65+ with disbetes W 17.0% 12.0% 65+ with stroke W 12.3% 10.8% 65+ with stroke W 12.3% 10.8% 65+ with conic obstructive pulmonary disease B 3.2.7% 3.4.3% 65+ with hypertension 71.0% 70.2%	% 60+ with 15+ days poor mental health last month		6.9%	6.9%
% 65+ with bipolar disorders W 5.5% 3.1% % 65+ with post-traumatic stress disorder W 2.1% 1.4% % 65+ with personality disorders W 8.5% 4.9% % 65+ with personality disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1,279 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% 65+ with tobacco use disorders W 12.0% 12.0% CHRONIC DISEASE W 17.0% 12.0% 65+ with diabetes 29.6% 28.2% 65+ with stroke W 12.3% 10.8% 65+ with trorolic obstructive pulmonary disease W 12.3% 10.8% 65+ with stroke W 12.3% 10.8% 65+ with thronic obstructive pulmonary disease W 12.3% 10.8% 65+ with attalfabilities T,10% 70.2% 65+ with premission 71.0% 70.2% 65+ with congestive heart failure B 32.7% 34.3%	% 65+ with depression	W	35.4%	28.8%
% 65+ with bost-traumatic stress disorder W 2.1% 1.4% % 65+ with schizophrenia & other psychotic disorders W 8.5% 4.9% % 65+ with personality disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1.279 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% 65+ with tobacco use disorders W 12.2% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% 65+ with stroke W 12.3% 10.8% 65+ with stroke W 12.3% 10.8% 65+ with horronic obstructive pulmonary disease W 12.3% 10.8% 65+ with papertension 71.0% 70.2% 65+ with papertension 71.0% 70.2% 65+ with papertension 8 32.7% 34.3% 65+ with congestive heart disease B 32.7% 34.3% 65+ with bosteoporosis W 16.3% 17.7% <t< td=""><td>% 65+ with anxiety disorders</td><td>W</td><td>28.8%</td><td>21.9%</td></t<>	% 65+ with anxiety disorders	W	28.8%	21.9%
% 65+ with schizophrenia & other psychotic disorders W 8.5% 4.9% % 65+ with personality disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1.279 6 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% 6 65+ with tobacco use disorders W 12.2% 10.4% CHRONIC DISEASE 8 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% 6 65+ with diabetes 29.6% 28.2% 6 65+ with stroke W 12.3% 10.8% 6 65+ with strokic W 12.3% 10.8% 6 65+ with strokic obstructive pulmonary disease W 12.3% 10.5% 6 65+ with asthma 13.6% 13.0% 13.0% 6 65+ with asthma 13.6% 13.0% 13.0% 6 65+ with asthma 13.6% 13.0% 14.5% 6 65+ with asthma 13.6% 13.0% 14.5% 6 5+ with schemic heart disease B 32.7% 34.3%	% 65+ with bipolar disorders	W	5.5%	3.1%
% 65+ with personality disorders W 2.7% 1.1% # opioid deaths (all ages) (county) 118 1,279 % 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% % 65+ with babacco use disorders W 12.2% 10.4% CHRONIC DISEASE W 17.0% 12.0% 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% % 65+ with Alzheimer's disease or related dementias W 12.3% 12.0% % 65+ with Alzheimer's disease or related dementias W 12.3% 12.0% % 65+ with Alzheimer's disease or related dementias W 12.3% 12.0% % 65+ with diabetes 29.6% 28.2% 28.2% % 65+ with obstance in with poper disease W 12.3% 10.8% % 65+ with poper disease and a heart attack 4.5% 4.5% % 65+ with osteoarthritis/rheumatoid arthritis W 16.3% 17.7% % 65+ with osteoarthritis/rheumatoid arthritis W 16.3% 17.4% % 65+ with lung can	% 65+ with post-traumatic stress disorder	W	2.1%	1.4%
# opioid deaths (all ages) (county) % 65+ with substance use disorders (drug use +/or alcohol abuse) % 65+ with tobacco use disorders W 12.2% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias % 65+ with diabetes % 65+ with diabetes % 65+ with chronic obstructive pulmonary disease % 65+ with stroke W 12.3% 20.5% % 65+ with hashma 13.6% 13.0% % 65+ with nashma 13.6% 13.0% % 65+ with nashma 13.6% 13.0% % 65+ with nashma 13.6% 13.0% % 65+ with inchemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with ongestive heart failure 18.3% 14.4% % 65+ with ongestive heart failure 18.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 19.6% 14.4% % 65+ with osteoarthritis/rheumatoid arthritis W 19.6% 10.0% % 65+ with lung cancer 18.3% 1.5% % 65+ women with prostate cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.5% 11.5% % 65+ women with prostate cancer W 23.5% 22.8% % 65+ with HIV/AIDS W 23.5% 20.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with prophyroidism W 23.5% 20.8% % 65+ with lung cancer W 39.4% 36.8% % 65+ with prophyroidism W 23.5% 20.8% % 65+ with chronic kidney disease W 25.1% 20.3% % 65+ with chronic kidney disease	% 65+ with schizophrenia & other psychotic disorders	W	8.5%	4.9%
% 65+ with substance use disorders (drug use +/or alcohol abuse) W 7.6% 5.5% % 65+ with tobacco use disorders W 12.2% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% % 65+ with disbetes 29.6% 28.2% % 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with nonic obstructive pulmonary disease 20.3% 20.5% % 65+ with ashma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 6.6+ with lung cancer 1.6%	% 65+ with personality disorders	W	2.7%	1.1%
% 65+ with tobacco use disorders W 12.2% 10.4% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% % 65+ with diabetes 29.6% 28.2% % 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease W 12.3% 10.8% % 65+ with asthma 31.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ with bypertension 71.0% 70.2% % 65+ with propertines of the eart disease B 32.7% 34.3% % 65+ with compastive heart failure 18.3% 17.7% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with leukemias and lymphomas B	# opioid deaths (all ages) (county)		118	1,279
CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% % 65+ with diabetes 29.6% 28.2% % 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with asthma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ with pypertension 4.5% 4.5% % 65+ with pypertension 4.5% 4.5% % 65+ with pypertension 4.5% 4.5% % 65+ with schemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoporosis W 16.3% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer W 11.2% 9.8% % 65+ women with presat cancer W	% 65+ with substance use disorders (drug use +/or alcohol abuse)	W	7.6%	5.5%
% 65+ with Alzheimer's disease or related dementias W 17.0% 12.0% % 65+ with diabetes 29.6% 28.2% % 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with asthma 13.6% 13.0% % 65+ with pypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer W 19.6% 17.4% % 65+ with lung cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.2% 9.8% % 65+ women with prostate cancer W 39.4% 36.8% % 65+ with ben	% 65+ with tobacco use disorders	W	12.2%	10.4%
% 65+ with diabetes 29.6% 28.2% % 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with asthma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis//rheumatoid arthritis W 52.0% 49.1% % 65+ with lug cancer W 19.6% 17.4% % 65+ with lug cancer B 1.5% 2.0% % 65+ with colon cancer 2.3% 2.4% % 65+ with prostate cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.5% 11.5% % 65+ women with prostate cancer W 39.4% 36.8% % 65+ with hypothyroidism W </td <td>CHRONIC DISEASE</td> <td></td> <td></td> <td></td>	CHRONIC DISEASE			
% 65+ with stroke W 12.3% 10.8% % 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with asthma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with lug cancer W 19.6% 17.4% % 65+ with lug cancer B 1.5% 2.0% % 65+ with lug cancer 1.6% 1.6% 1.6% % 65+ women with preast cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.5% 11.5% % 65+ women with prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with nemia <td>% 65+ with Alzheimer's disease or related dementias</td> <td>W</td> <td>17.0%</td> <td>12.0%</td>	% 65+ with Alzheimer's disease or related dementias	W	17.0%	12.0%
% 65+ with chronic obstructive pulmonary disease 20.3% 20.5% % 65+ with asthma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ women with presat cancer W 11.2% 9.8% % 65+ momen with prostate cancer W 11.5% 11.5% % 65+ men with prostate cancer W 39.4% 36.8% % 65+ mith hypothyroidism W 23.5% 20.8% % 65+ with hypothyroidism W 23.5% 20.8% % 6	% 65+ with diabetes		29.6%	28.2%
% 65+ with asthma 13.6% 13.0% % 65+ with hypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoparthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with nemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases	% 65+ with stroke	W	12.3%	10.8%
% 65+ with hypertension 71.0% 70.2% % 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with lung cancer B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.2% 9.8% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with hyl/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with nemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 20.3%	% 65+ with chronic obstructive pulmonary disease		20.3%	20.5%
% 65+ ever had a heart attack 4.5% 4.5% % 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ men with prostate cancer W 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9% <	% 65+ with asthma		13.6%	13.0%
% 65+ with ischemic heart disease B 32.7% 34.3% % 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ men with prostate cancer W 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with hypertension		71.0%	70.2%
% 65+ with congestive heart failure 18.3% 17.7% % 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with colon cancer 1.6% 1.6% 1.6% % 65+ with colon cancer W 11.2% 9.8% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with prostate cancer W 11.5% 11.5% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ ever had a heart attack		4.5%	4.5%
% 65+ with atrial fibrillation 14.5% 14.4% % 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with ischemic heart disease	В	32.7%	34.3%
% 65+ with peripheral vascular disease W 16.3% 14.7% % 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with congestive heart failure		18.3%	17.7%
% 65+ with osteoarthritis/rheumatoid arthritis W 52.0% 49.1% % 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with atrial fibrillation		14.5%	14.4%
% 65+ with osteoporosis W 19.6% 17.4% % 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with peripheral vascular disease	W	16.3%	14.7%
% 65+ with leukemias and lymphomas B 1.5% 2.0% % 65+ with lung cancer 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with osteoarthritis/rheumatoid arthritis	W	52.0%	49.1%
% 65+ with lung cancer 1.6% 1.6% % 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with osteoporosis	W	19.6%	17.4%
% 65+ with colon cancer 2.3% 2.4% % 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with leukemias and lymphomas	В	1.5%	2.0%
% 65+ women with breast cancer W 11.2% 9.8% % 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with lung cancer		1.6%	1.6%
% 65+ women with endometrial cancer 1.7% 1.7% % 65+ men with prostate cancer 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with colon cancer		2.3%	2.4%
% 65+ men with prostate cancer 11.5% 11.5% % 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ women with breast cancer	W	11.2%	9.8%
% 65+ with benign prostatic hyperplasia W 39.4% 36.8% % 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ women with endometrial cancer		1.7%	1.7%
% 65+ with HIV/AIDS 0.07% 0.05% % 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ men with prostate cancer		11.5%	11.5%
% 65+ with hypothyroidism W 23.5% 20.8% % 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with benign prostatic hyperplasia	W	39.4%	36.8%
% 65+ with anemia 36.0% 37.3% % 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with HIV/AIDS		0.07%	0.05%
% 65+ with chronic kidney disease W 25.1% 22.3% % 65+ with liver diseases B 5.7% 6.9%	% 65+ with hypothyroidism	W	23.5%	20.8%
% 65+ with liver diseases B 5.7% 6.9%	% 65+ with anemia		36.0%	37.3%
% 65+ with liver diseases B 5.7% 6.9%	% 65+ with chronic kidney disease	W	25.1%	22.3%
% 65+ with fibromyalgia, chronic pain and fatigue W 22.1% 18.6%	% 65+ with liver diseases	В	5.7%	6.9%
	% 65+ with fibromyalgia, chronic pain and fatigue	W	22.1%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache	W	4.8%	4.0%
% 65+ with epilepsy	W	3.0%	2.1%
% 65+ with traumatic brain injury	W	2.5%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma	W	28.6%	22.9%
% 65+ with cataract	W	66.0%	61.2%
% 65+ with pressure ulcer or chronic ulcer	W	9.7%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	W	58.3%	54.4%
% 65+ with 0 chronic conditions	W	8.4%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		14.6%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment	W	16.0%	14.4%
% 65+ with self-reported vision difficulty		5.8%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment	W	1.5%	0.9%
% 65+ with self-reported cognition difficulty	W	12.0%	6.9%
% 65+ with self-reported ambulatory difficulty		21.8%	18.8%
% 65+ with clinical diagnosis of mobility impairments	W	4.2%	3.2%
% 65+ with self-reported self-care difficulty		7.3%	5.6%
% 65+ with self-reported independent living difficulty	W	17.0%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		7.9%	7.9%
% dually eligible for Medicare and Medicaid	*	12.4%	7.5%
% 60+ with a regular doctor	В	98.4%	95.8%
% 60+ who did not see doctor when needed due to cost		4.1%	5.4%
# of primary care providers within 5 miles		163	2,961
# of hospitals within 5 miles		1	26
# of nursing homes within 5 miles		4	74
# of home health agencies		8	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		1	12
# of dementia-related support groups		1	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year		6.4	6.3
Emergency room visits/1000 persons 65+ years per year	*	723	572
Part D monthly prescription fills per person per year	*	56.5	49.1
Home health visits per year	*	5.0	2.5
Durable medical equipment claims per year		2.0	2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	279	237
Medicare inpatient hospital readmissions (as % of admissions)		16.0%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year	*	96	76
# skilled nursing home Medicare beds/1000 persons 65+ years		51	33
% 65+ getting Medicaid long term services and supports	*	8.2%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		3	44
Air pollution: annual # of unhealthy days for 65+ (county)		0	N/A
% of grandparents raising grandchildren		0.6%	0.8%
% of grandparents who live with grandchildren		2.5%	2.5%
# of assisted living sites		3	134
% of vacant homes in community		7.3%	16.0%
# of universities and community colleges		3	41
# of public libraries		2	234
# of YMCAs		1	12
% in county with access to broadband (all ages)		92.0%	93.0%
% 60+ who used Internet in last month		78.7%	77.6%
Voter participation rate in 2018 election (age 18+)		52.4%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		228	207
Homicide rate /100,000 persons (county)		N/A	1
# firearm fatalities (county)		51	586
Property crime rate /100,000 persons		2,255	2,012
% 65+ who own a motor vehicle		86.0%	91.0%
% 60+ who always drive wearing a seatbelt		80.4%	77.1%
# of fatal crashes involving adult age 60+/town		3	151
# of fatal crashes involving adult age 60+/county		18	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		6.2%	5.4%
% 60+ receiving food stamps past year		7.8%	5.7%
% 65+ employed past year		20.6%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		18.6%	18.2%
% households with annual income \$20,000-\$49,999		42.5%	36.5%
% households with annual income > \$50,000		38.9%	45.3%
% 60+ own home		59.3%	79.9%
% 60+ have mortgage on home		26.4%	35.3%
% 65+ households spend >35% of income on housing (renter)		22.3%	8.7%
% 65+ households spend >35% of income on housing (owner)		15.5%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$25,860	\$25,284	1.02
Single, renter, good health	\$26,376	\$26,400	1.00
Couple, homeowner without mortgage, good health	\$37,800	\$37,128	1.02
Couple, renter, good health	\$38,316	\$38,244	1.00

TECHNICAL NOTES

*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older people, and when the implication is unclear, we use an *. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town and 5 Manchester neighborhoods, 4 Nashua neighborhoods). For example, the population characteristics were reported for all 244 units. For other data (i.e., highly prevalent chronic disease, health services utilization) we could report for 154 geographic units. For less prevalent conditions we report for 69 geographic units. For the BRFSS data we report for 28 geographic units, and for the lowest prevalence conditions (e.g., HIV) we report for 4 geographic units. The same estimate is reported for all cities/towns within aggregated geographic areas. Maps of the different geographic groupings and the rationale behind the groupings are in the Technical Report.

Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Agriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team.</u> Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019