## **Easton (Grafton)**



Easton is a rural town located in Grafton County situated between Mount Moosilauke and Mount Lafayette. There are 78 residents age 65 or older. Compared to state average rates, older residents fared better on some healthy aging indicators with lower rates of obesity, high cholesterol, anxiety disorders, bipolar and tobacco use disorders, Alzheimer's disease or related dementias, diabetes, chronic obstructive pulmonary disease, asthma, hypertension, ischemic heart disease, congestive heart failure, peripheral vascular disease, osteoarthritis/rheumatoid arthritis, benign prostatic hyperplasia, hypothyroidism, anemia, chronic kidney disease, and fibromyalgia/chronic pain/fatigue. Community resources to support healthy aging include 2 home health agencies and a senior center.

BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE 274	STATE ESTIMATE 1,327,503
	274	1 327 502
		1,527,503
	49.3%	22.7%
	135	301,630
	28.5%	15.8%
	78	210,385
	66.7%	58.5%
	29.5%	28.6%
	3.8%	12.9%
	50.0%	54.7%
	100.0%	97.7%
	0.0%	0.5%
	0.0%	0.9%
	0.0%	0.9%
	0.0%	0.9%
	76.9%	58.5%
	15.4%	14.0%
	2.6%	22.9%
	5.1%	4.6%
	9.0%	12.3%
	37.2%	57.1%
	53.8%	30.6%
	21.8%	26.1%
	91.0%	91.3%
	21.8%	24.8%
	3.4%	4.1%
		28.5% 78 66.7% 29.5% 3.8% 50.0% 50.0% 0.0% 0.0% 0.0% 0.0% 0.0% 0

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
Geographic Migration (65+ population) in the past 12 months			
% moved within same county		0.0%	3.6%
% moved from different county in New Hampshire		0.0%	1.0%
% moved from different state		0.0%	1.7%
WELLNESS & PREVENTION			
% 60+ with any physical activity within last month		73.6%	74.5%
% 60+ met CDC guidelines for muscle-strengthening activity		23.8%	26.0%
% 60+ met CDC guidelines for aerobic physical activity		55.3%	58.6%
% 60+ met CDC guidelines for both types of physical activities		17.3%	19.9%
% 60+ getting recommended hours of sleep		64.8%	66.4%
% 60+ injured in a fall within last 12 months		9.6%	10.4%
% 65+ had hip fracture		3.4%	3.3%
% 60+ with self-reported fair or poor health status		18.0%	16.5%
% 60+ with 15+ physically unhealthy days last month		16.6%	12.5%
% 60+ with physical exam/check-up in past year		83.2%	86.5%
% 60+ met CDC preventive health screening goals		38.2%	40.3%
% 60+ flu shot past year		52.1%	59.3%
% 65+ with pneumonia vaccine		74.9%	77.8%
% 60+ with cholesterol screening		92.9%	95.3%
% 60+ women with a mammogram within last 2 years		77.0%	79.3%
% 60+ with colorectal cancer screening		80.9%	77.0%
% 60+ with HIV test		16.2%	13.5%
% 60+ current smokers		10.6%	8.1%
Oral Health			
% 60+ with loss of 6 or more teeth		33.6%	29.0%
% 60+ with annual dental exam		69.2%	75.7%
# of dentists per 100,000 persons (all ages) (county)		83	72
NUTRITION/DIET			
% 60+ with 5 or more servings of fruit or vegetables per day		15.2%	20.2%
% 60+ self-reported obese		27.6%	27.2%
% 65+ clinically diagnosed obese	В	10.9%	16.7%
% 65+ with high cholesterol	В	62.8%	72.2%
% 60+ excessive drinking		7.6%	9.2%
% 65+ with poor supermarket access		0.0%	28.4%

AVIORAL HEALTH + with 15+ days poor mental health last month + with depression + with anxiety disorders + with bipolar disorders + with post-traumatic stress disorder + with schizophrenia & other psychotic disorders + with personality disorders oid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse) + with tobacco use disorders	B B B B B B	8.0% 25.2% 17.2% 1.6% 1.8% 4.4% 0.9% 45 4.7% 5.8%	6.9% 28.8% 21.9% 3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with depression</li> <li>+ with anxiety disorders</li> <li>+ with bipolar disorders</li> <li>+ with post-traumatic stress disorder</li> <li>+ with schizophrenia &amp; other psychotic disorders</li> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	B	25.2% 17.2% 1.6% 1.8% 4.4% 0.9% 45 4.7% 5.8% 9.3%	28.8% 21.9% 3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with anxiety disorders</li> <li>+ with bipolar disorders</li> <li>+ with post-traumatic stress disorder</li> <li>+ with schizophrenia &amp; other psychotic disorders</li> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	B	17.2% 1.6% 1.8% 4.4% 0.9% 45 4.7% 5.8% 9.3%	21.9% 3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with bipolar disorders</li> <li>+ with post-traumatic stress disorder</li> <li>+ with schizophrenia &amp; other psychotic disorders</li> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	B	1.6%         1.8%         4.4%         0.9%         45         4.7%         5.8%         9.3%	3.1% 1.4% 4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with post-traumatic stress disorder</li> <li>+ with schizophrenia &amp; other psychotic disorders</li> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	B	1.8%         4.4%         0.9%         45         4.7%         5.8%         9.3%	1.4% 4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with schizophrenia &amp; other psychotic disorders</li> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	В	4.4% 0.9% 45 4.7% 5.8% 9.3%	4.9% 1.1% 1,279 5.5% 10.4%
<ul> <li>+ with personality disorders</li> <li>oid deaths (all ages) (county)</li> <li>+ with substance use disorders (drug use +/or alcohol abuse)</li> </ul>	В	0.9% 45 4.7% 5.8% 9.3%	1.1% 1,279 5.5% 10.4%
oid deaths (all ages) (county) + with substance use disorders (drug use +/or alcohol abuse)	В	45 4.7% 5.8% 9.3%	1,279 5.5% 10.4%
+ with substance use disorders (drug use +/or alcohol abuse)	В	4.7% 5.8% 9.3%	5.5% 10.4%
	В	5.8% 9.3%	10.4%
+ with tobacco use disorders	В	9.3%	
			12.00/
ONIC DISEASE			10 00/
+ with Alzheimer's disease or related dementias	В		12.0%
+ with diabetes		18.0%	28.2%
+ with stroke		8.3%	10.8%
+ with chronic obstructive pulmonary disease	В	9.2%	20.5%
+ with asthma	В	10.2%	13.0%
+ with hypertension	В	61.0%	70.2%
+ ever had a heart attack		3.0%	4.5%
+ with ischemic heart disease	В	24.4%	34.3%
+ with congestive heart failure	В	13.7%	17.7%
+ with atrial fibrillation		12.1%	14.4%
+ with peripheral vascular disease	В	10.4%	14.7%
+ with osteoarthritis/rheumatoid arthritis	В	43.0%	49.1%
+ with osteoporosis		17.0%	17.4%
+ with leukemias and lymphomas		1.7%	2.0%
+ with lung cancer		1.4%	1.6%
+ with colon cancer		2.6%	2.4%
+ women with breast cancer		9.5%	9.8%
+ women with endometrial cancer		2.0%	1.7%
+ men with prostate cancer		12.6%	11.5%
+ with benign prostatic hyperplasia	В	29.4%	36.8%
+ with HIV/AIDS		0.05%	0.05%
+ with hypothyroidism	В	15.6%	20.8%
+ with anemia	В	31.0%	37.3%
+ with chronic kidney disease	В	11.7%	22.3%
+ with liver diseases		5.0%	6.9%
+ with fibromyalgia, chronic pain and fatigue	В	13.5%	18.6%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
% 65+ with migraine and other chronic headache		3.8%	4.0%
% 65+ with epilepsy		1.8%	2.1%
% 65+ with traumatic brain injury		0.9%	1.1%
% 65+ with autism spectrum disorders		N/A	0.03%
% 65+ with glaucoma		25.2%	22.9%
% 65+ with cataract		65.1%	61.2%
% 65+ with pressure ulcer or chronic ulcer		5.4%	7.1%
% 65+ with 4+ (out of 15) chronic conditions	В	39.2%	54.4%
% 65+ with 0 chronic conditions	В	13.7%	10.3%
LIVING WITH DISABILITY			
% 65+ with self-reported hearing difficulty		10.3%	15.0%
% 65+ with clinical diagnosis of deafness or hearing impairment		17.5%	14.4%
% 65+ with self-reported vision difficulty		9.0%	5.2%
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	0.9%
% 65+ with self-reported cognition difficulty		11.5%	6.9%
% 65+ with self-reported ambulatory difficulty		15.4%	18.8%
% 65+ with clinical diagnosis of mobility impairments		2.7%	3.2%
% 65+ with self-reported self-care difficulty		5.1%	5.6%
% 65+ with self-reported independent living difficulty		10.3%	11.3%
ACCESS TO CARE			
Medicare (65+ population)			
% Medicare managed care enrollees		6.6%	7.9%
% dually eligible for Medicare and Medicaid	*	4.6%	7.5%
% 60+ with a regular doctor		96.4%	95.8%
% 60+ who did not see doctor when needed due to cost		9.7%	5.4%
# of primary care providers within 5 miles		0	2,961
# of hospitals within 5 miles		0	26
# of nursing homes within 5 miles		0	74
# of home health agencies		2	49
# of community health centers		0	22
# of adult day health centers		0	21
# of memory cafes		0	12
# of dementia-related support groups		0	14

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
SERVICE UTILIZATION			
Physician visits per year	*	4.4	6.3
Emergency room visits/1000 persons 65+ years per year		480	572
Part D monthly prescription fills per person per year	*	37.8	49.1
Home health visits per year	*	1.2	2.5
Durable medical equipment claims per year	*	1.3	2.0
Inpatient hospital stays/1000 persons 65+ years per year		184	237
Medicare inpatient hospital readmissions (as % of admissions)		18.8%	15.8%
# skilled nursing facility stays/1000 persons 65+ years per year		71	76
# skilled nursing home Medicare beds/1000 persons 65+ years		122	33
% 65+ getting Medicaid long term services and supports		2.6%	3.7%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT			
AARP Age-Friendly efforts in community		Not yet	Yes
# of senior centers		1	44
Air pollution: annual # of unhealthy days for 65+ (county)		0	N/A
% of grandparents raising grandchildren		2.3%	0.8%
% of grandparents who live with grandchildren		2.3%	2.5%
# of assisted living sites		0	134
% of vacant homes in community		36.6%	16.0%
# of universities and community colleges		0	41
# of public libraries		0	234
# of YMCAs		0	12
% in county with access to broadband (all ages)		89.0%	93.0%
% 60+ who used Internet in last month		72.9%	77.6%
Voter participation rate in 2018 election (age 18+)		80.0%	54.7%
SAFETY & TRANSPORTATION			
Violent crime rate /100,000 persons		N/A	207
Homicide rate /100,000 persons (county)		2	1
# firearm fatalities (county)		40	586
Property crime rate /100,000 persons		N/A	2,012
% 65+ who own a motor vehicle		86.8%	91.0%
% 60+ who always drive wearing a seatbelt		73.0%	77.1%
# of fatal crashes involving adult age 60+/town		1	151
# of fatal crashes involving adult age 60+/county		17	151

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	STATE ESTIMATE
ECONOMIC & HOUSING VARIABLES			
% 65+ with income below the poverty line past year		0.0%	5.4%
% 60+ receiving food stamps past year		7.0%	5.7%
% 65+ employed past year		23.1%	24.8%
Household income (65+ householder)			
% households with annual income < \$20,000		7.5%	18.2%
% households with annual income \$20,000-\$49,999		49.1%	36.5%
% households with annual income > \$50,000		43.4%	45.3%
% 60+ own home		98.8%	79.9%
% 60+ have mortgage on home		26.2%	35.3%
% 65+ households spend >35% of income on housing (renter)		0.0%	8.7%
% 65+ households spend >35% of income on housing (owner)		22.6%	21.2%
COST OF LIVING	\$ COUNTY ESTIMATE	\$ STATE ESTIMATE	RATIO (COUNTY/STATE)
Elder Economic Security Standard Index			
Single, homeowner without mortgage, good health	\$23,496	\$25,284	0.93
Single, renter, good health	\$26,592	\$26,400	1.01
Couple, homeowner without mortgage, good health	\$35,436	\$37,128	0.95
Couple, renter, good health	\$38,532	\$38,244	1.01

## TECHNICAL NOTES

\*See our technical report (online at https://healthyagingdatareports.org/) for more information on data sources, measures, methodology, and margin of errors. For most indicators the reported community and state values are both estimates derived from sample data. Thus, it is possible that some of the differences between state and community estimates may be due to chance associated with population sampling. We use the terms "better" and "worse" to highlight differences between community and state estimates that we are confident are <u>not</u> due to chance. "Better" is used where a higher/lower value has positive implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications for the health of older residents. "Worse" is used where a higher/lower score has negative implications to report the health of older people, and when the implication is unclear, we use an \*. We balance two goals. First, we aim to report data at very local levels because we believe change is often locally driven. Second, we vowed to protect the privacy of the people providing the information reported. Thus, given the constraints of the data analyzed we used a hierarchical approach to reporting. When possible we report estimates for 244 geographic units (i.e., every NH city/town an

## Data Sources:

- Population Characteristics: The U.S. Census Bureau (American Community Survey 2012-2016).
- Wellness & Prevention: The Behavioral Risk Factor Surveillance System (2011-2016).
- Nutrition/Diet: BRFSS (2011-2016), CMS (2014-2015), and the U.S. Department of Ágriculture Food Atlas (2017).
- Behavioral Health: BRFSS (2011-2016), CMS (2014-2015), CDC Wonder website (2014-2016).
- Chronic Disease: The Master Beneficiary Summary File ABCD/Other from CMS (2014-2015).
- Disability: CMS (2014-2015) for the clinical measures, and ACS (2012-2016) for the self-reported disability.
- Access to Care: BRFSS (2011-2016), CMS (2015), Medicare.gov (June-July 2018), the NH Division of Public Health Services (2018), National Adult Day Services Association (2018), memorycaredirectory.com (2018), and the Alzheimer's Association (July 2018).
- Service Utilization: CMS (2015), and Medicare Nursing Home Compare (December 2018).
- Community & Civic Engagement: AARP (2018 update; https://www.aarp.org/livable-communities/network-age-friendly-communities/info-2014/member-list.html), the Aging & Disability Resource Center, U.S. Environmental Protection Agency Air Compare (2016), assistedlivingfacilities.org, ACS (2012-2016), the NH Department of Business and Economic Affairs Division of Travel and Tourism Development (August 2018), NH YMCA (July 2018), the Federal Communications Commission (2016), BRFSS (2011-2016), and the NH Secretary of State.
- Safety & Transportation: U.S. Department of Justice Federal Bureau of Investigation (August 2017), the County Health Rankings (2018), BRFSS (2011-2016), ACS (2012-2016), and the National Highway Traffic Safety Administration (2011-2015).
- Economic & Housing, Cost of Living: ACS (2012-2016) and the Center for Social and Demographic Research on Aging at the University of Massachusetts Boston (August 2017).

<u>Healthy Aging Data Report Team</u>. Many people contributed to this research. The 2018 research team: Beth Dugan PhD, Frank Porell PhD, Nina Silverstein PhD, Chae Man Lee PhD, ShuangShuang Wang PhD, Bon Kim, Natalie Pitheckoff, Haowei Wang, Sae Hwang Han, Richard Chunga, & Shiva Prisad from the Gerontology Institute in the McCormack Graduate School of Policy and Global Studies at the University of Massachusetts Boston. The Tufts Health Plan Foundation supported the research and provided important guidance. Our Advisory Committees helped to make the Data Report best address the needs of New Hampshire. We thank JSI for their continued partnership. Questions? Beth.dugan@umb.edu Update: 05/2019