

Hampton (Rockingham)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		4.0%	[3.35% - 4.73%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month	B	84.5%	[79.94% - 89.10%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity	B	37.2%	[30.96% - 43.49%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		60.6%	[54.08% - 67.04%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities	B	29.6%	[23.79% - 35.46%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		65.1%	[58.24% - 72.04%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		9.0%	[5.48% - 12.51%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		3.4%	[2.80% - 4.00%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		12.8%	[8.39% - 17.25%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		9.6%	[5.84% - 13.37%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		90.9%	[87.11% - 94.66%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		45.0%	[37.21% - 52.78%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		59.3%	[52.85% - 65.85%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		79.5%	[73.64% - 85.41%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		96.9%	[95.01% - 98.89%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		80.6%	[73.46% - 87.66%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		82.4%	[77.48% - 87.40%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		12.6%	[7.46% - 17.73%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		6.4%	[3.41% - 9.46%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth	B	20.4%	[14.88% - 25.87%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam	B	85.1%	[79.58% - 90.59%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		26.9%	[21.13% - 32.69%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		21.1%	[15.57% - 26.53%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese	B	14.4%	[12.98% - 15.75%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	W	74.7%	[73.02% - 76.31%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		9.5%	[5.46% - 13.52%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		5.8%	[2.67% - 8.98%]	6.9%	[6.29% - 7.61%]
% 65+ with depression	B	26.7%	[25.03% - 28.28%]	28.8%	[28.59% - 29.01%]
% 65+ with anxiety disorders		22.7%	[21.04% - 24.31%]	21.9%	[21.74% - 22.15%]
% 65+ with bipolar disorders		2.7%	[2.11% - 3.29%]	3.1%	[2.97% - 3.14%]
% 65+ with post-traumatic stress disorder		1.5%	[0.98% - 1.93%]	1.4%	[1.35% - 1.48%]
% 65+ with schizophrenia & other psychotic disorders		4.5%	[3.78% - 5.26%]	4.9%	[4.81% - 5.02%]
% 65+ with personality disorders		0.8%	[0.47% - 1.10%]	1.1%	[1.00% - 1.11%]
% 65+ with substance use disorders (drug use +/- alcohol abuse)		5.9%	[4.98% - 6.85%]	5.5%	[5.41% - 5.64%]
% 65+ with tobacco use disorders		10.2%	[8.97% - 11.41%]	10.4%	[10.23% - 10.54%]
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	8.9%	[7.84% - 9.87%]	12.0%	[11.89% - 12.19%]
% 65+ with diabetes	B	26.2%	[24.49% - 27.92%]	28.2%	[27.97% - 28.41%]
% 65+ with stroke		10.0%	[8.91% - 11.07%]	10.8%	[10.64% - 10.92%]
% 65+ with chronic obstructive pulmonary disease		19.6%	[18.18% - 21.06%]	20.5%	[20.36% - 20.73%]
% 65+ with asthma	B	10.6%	[9.43% - 11.68%]	13.0%	[12.89% - 13.21%]
% 65+ with hypertension		70.4%	[68.67% - 72.12%]	70.2%	[70.02% - 70.46%]
% 65+ ever had a heart attack		4.8%	[4.06% - 5.60%]	4.5%	[4.37% - 4.56%]
% 65+ with ischemic heart disease		36.3%	[34.43% - 38.18%]	34.3%	[34.12% - 34.58%]
% 65+ with congestive heart failure		17.2%	[15.74% - 18.59%]	17.7%	[17.57% - 17.93%]
% 65+ with atrial fibrillation		15.3%	[14.00% - 16.60%]	14.4%	[14.28% - 14.60%]
% 65+ with peripheral vascular disease		15.8%	[14.40% - 17.11%]	14.7%	[14.51% - 14.84%]
% 65+ with osteoarthritis/rheumatoid arthritis		48.9%	[46.93% - 50.89%]	49.1%	[48.82% - 49.32%]
% 65+ with osteoporosis		17.3%	[15.96% - 18.67%]	17.4%	[17.22% - 17.57%]
% 65+ with leukemias and lymphomas		2.0%	[1.50% - 2.50%]	2.0%	[1.91% - 2.05%]
% 65+ with lung cancer		1.6%	[1.18% - 2.05%]	1.6%	[1.54% - 1.66%]
% 65+ with colon cancer		2.4%	[1.90% - 2.93%]	2.4%	[2.37% - 2.51%]
% 65+ women with breast cancer		10.9%	[9.41% - 12.48%]	9.8%	[9.59% - 9.96%]
% 65+ women with endometrial cancer		1.8%	[1.21% - 2.43%]	1.7%	[1.60% - 1.76%]
% 65+ men with prostate cancer		12.8%	[10.98% - 14.62%]	11.5%	[11.24% - 11.68%]
% 65+ with benign prostatic hyperplasia		37.1%	[34.41% - 39.71%]	36.8%	[36.47% - 37.14%]
% 65+ with HIV/AIDS		0.04%	[0.02% - 0.06%]	0.05%	[0.04% - 0.06%]
% 65+ with hypothyroidism		20.5%	[19.04% - 21.98%]	20.8%	[20.62% - 20.99%]
% 65+ with anemia		38.9%	[37.12% - 40.69%]	37.3%	[37.12% - 37.57%]
% 65+ with chronic kidney disease		22.6%	[21.02% - 24.21%]	22.3%	[22.14% - 22.54%]
% 65+ with liver diseases	W	8.2%	[7.16% - 9.32%]	6.9%	[6.75% - 6.99%]
% 65+ with fibromyalgia, chronic pain and fatigue		18.8%	[17.29% - 20.33%]	18.6%	[18.42% - 18.80%]
% 65+ with migraine and other chronic headache		4.4%	[3.68% - 5.19%]	4.0%	[3.95% - 4.15%]
% 65+ with epilepsy		2.0%	[1.55% - 2.54%]	2.1%	[2.00% - 2.14%]
% 65+ with traumatic brain injury		1.2%	[0.83% - 1.61%]	1.1%	[1.06% - 1.16%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma	B	20.7%	[19.21% - 22.13%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	W	63.5%	[61.69% - 65.33%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		7.6%	[6.65% - 8.63%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions		53.9%	[52.03% - 55.74%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		9.6%	[8.41% - 10.70%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		15.2%	[13.85% - 16.56%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	[0.51% - 1.10%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		2.8%	[2.21% - 3.38%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees		8.0%	[7.16% - 8.93%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	5.1%	[4.34% - 5.76%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		97.4%	[95.22% - 99.64%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		2.6%	[0.32% - 4.83%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	7.8	[7.5 - 8.1]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year		568	[511 - 624]	572	[566 - 578]
Part D monthly prescription fills per person per year	*	45.0	[43.5 - 46.5]	49.1	[48.9 - 49.3]
Home health visits per year		3.1	[2.5 - 3.6]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year	*	1.6	[1.4 - 1.8]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		243	[217 - 269]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		14.0%	[11.01% - 16.95%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		73	[60 - 86]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	2.4%	[1.87% - 2.85%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month	*	86.3%	[82.13% - 90.46%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		83.7%	[78.05% - 89.38%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu