

Henniker (Merrimack)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		3.5%	[1.94% - 5.06%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		74.2%	[67.37% - 80.97%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		23.1%	[17.41% - 28.70%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		61.5%	[54.68% - 68.29%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		17.7%	[12.66% - 22.78%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		66.8%	[59.69% - 73.83%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		7.1%	[3.38% - 10.80%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		3.2%	[2.62% - 3.73%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		15.4%	[9.64% - 21.19%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		10.7%	[6.12% - 15.36%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		84.8%	[79.83% - 89.87%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		35.3%	[26.27% - 44.24%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		58.9%	[51.65% - 66.10%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		73.7%	[66.00% - 81.35%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		92.1%	[88.33% - 95.93%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		84.1%	[76.93% - 91.33%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		78.7%	[72.47% - 84.88%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		17.4%	[11.44% - 23.33%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		8.3%	[3.58% - 12.94%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		24.0%	[17.45% - 30.62%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		77.3%	[71.00% - 83.64%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	[17.23% - 28.22%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		24.3%	[17.69% - 30.94%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		15.7%	[11.98% - 19.39%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	B	64.5%	[59.82% - 69.18%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		6.5%	[2.94% - 10.08%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		7.8% [3.52% - 12.02%]	6.9% [6.29% - 7.61%]		
% 65+ with depression	B	23.4% [19.44% - 27.34%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders	B	17.4% [13.65% - 21.20%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders		2.5% [1.90% - 3.05%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder		1.1% [0.71% - 1.46%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders	B	3.7% [3.08% - 4.34%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders	B	0.5% [0.24% - 0.76%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		5.2% [2.92% - 7.47%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		9.0% [5.98% - 12.10%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	6.2% [4.06% - 8.27%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes	B	21.4% [17.30% - 25.54%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		8.2% [5.76% - 10.66%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease	B	11.5% [8.71% - 14.35%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma	B	9.3% [6.63% - 11.88%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension	B	59.2% [54.48% - 63.98%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack		3.6% [1.94% - 5.23%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease	B	26.9% [22.47% - 31.34%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure	B	10.8% [7.97% - 13.65%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		13.0% [9.93% - 16.00%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease	B	10.3% [7.37% - 13.15%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis	B	39.9% [34.99% - 44.85%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis	B	13.5% [10.40% - 16.53%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		1.5% [1.06% - 1.94%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.3% [0.90% - 1.64%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.0% [1.54% - 2.44%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		9.2% [5.49% - 12.92%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.9% [1.25% - 2.49%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		10.5% [6.41% - 14.56%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia	B	26.6% [20.74% - 32.36%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.07% [0.04% - 0.10%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism		18.3% [14.69% - 21.89%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia	B	28.4% [24.21% - 32.53%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease	B	17.3% [13.63% - 21.03%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases	B	3.5% [1.82% - 5.15%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		15.0% [11.44% - 18.50%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		4.3% [3.57% - 5.01%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy	B	1.1% [0.73% - 1.46%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury		1.4% [1.00% - 1.76%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		25.3%	[21.29% - 29.35%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	B	55.7%	[50.85% - 60.45%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		5.6%	[3.37% - 7.77%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	B	42.7%	[38.04% - 47.36%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions	B	14.0%	[10.44% - 17.49%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		13.9%	[10.59% - 17.27%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	[0.44% - 1.02%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments	B	2.5%	[1.97% - 3.03%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees		9.9%	[7.51% - 12.20%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	4.7%	[3.02% - 6.35%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		95.7%	[92.56% - 98.74%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		4.4%	[1.17% - 7.72%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year		5.9	[5.4 - 6.4]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year		523	[418 - 629]	572	[566 - 578]
Part D monthly prescription fills per person per year		45.1	[41.3 - 48.9]	49.1	[48.9 - 49.3]
Home health visits per year		3.0	[1.8 - 4.3]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		1.9	[1.3 - 2.5]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year	*	167	[108 - 226]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		27.5%	[14.39% - 40.61%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year	*	34	[10 - 58]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	1.6%	[0.62% - 2.61%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		83.0%	[77.62% - 88.43%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		75.0%	[68.20% - 81.83%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu