

Hooksett (Merrimack)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		4.1%	[3.24% - 5.04%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		72.8%	[67.02% - 78.59%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		21.3%	[16.48% - 26.20%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		58.6%	[52.52% - 64.77%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		16.9%	[12.35% - 21.41%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		66.9%	[60.24% - 73.48%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		7.6%	[4.56% - 10.69%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture	B	2.4%	[1.94% - 2.92%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		14.5%	[9.91% - 19.16%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		14.3%	[9.64% - 19.00%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		88.6%	[85.02% - 92.10%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		38.5%	[30.38% - 46.59%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		55.0%	[48.49% - 61.59%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		72.9%	[65.88% - 79.92%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		95.4%	[92.85% - 97.99%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		85.5%	[80.12% - 90.90%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		78.1%	[72.35% - 83.90%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		14.4%	[9.70% - 19.01%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		6.2%	[3.69% - 8.61%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth	B	22.2%	[16.75% - 27.74%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		80.9%	[75.38% - 86.43%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		24.1%	[18.86% - 29.44%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		24.2%	[18.77% - 29.65%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		17.4%	[15.40% - 19.47%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	W	78.1%	[76.04% - 80.12%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		9.4%	[5.83% - 13.02%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		7.0% [3.60% - 10.32%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		29.2% [27.03% - 31.35%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders		24.1% [21.87% - 26.33%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders		3.3% [2.58% - 3.95%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder	B	0.9% [0.48% - 1.23%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders	B	4.0% [3.28% - 4.62%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders		1.1% [0.69% - 1.42%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		5.1% [3.91% - 6.23%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		9.5% [7.94% - 11.05%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	9.9% [8.44% - 11.32%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes		29.7% [27.27% - 32.10%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		11.4% [9.90% - 12.85%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease		20.2% [18.28% - 22.05%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		14.0% [12.40% - 15.68%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension	W	73.9% [71.71% - 76.04%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack		3.6% [2.70% - 4.41%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease		33.0% [30.55% - 35.38%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure	B	15.4% [13.63% - 17.21%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		13.4% [11.81% - 15.02%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease		13.7% [11.99% - 15.38%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis	W	52.6% [49.90% - 55.21%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis		18.6% [16.85% - 20.44%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		2.1% [1.56% - 2.62%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.4% [1.04% - 1.82%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer	B	1.8% [1.31% - 2.19%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		9.9% [8.00% - 11.73%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.6% [0.99% - 2.14%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		11.8% [9.52% - 14.13%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia		38.2% [34.70% - 41.72%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.07% [0.04% - 0.10%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism	W	25.3% [23.23% - 27.35%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia		36.9% [34.61% - 39.18%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease		23.7% [21.56% - 25.88%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		8.1% [6.66% - 9.51%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		19.5% [17.48% - 21.54%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		4.1% [3.35% - 4.85%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy	B	1.1% [0.76% - 1.53%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury		0.9% [0.58% - 1.26%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma	W	26.3%	[24.19% - 28.35%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		60.6%	[58.23% - 63.01%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		7.1%	[5.82% - 8.34%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	W	58.0%	[55.62% - 60.42%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions	W	8.3%	[6.87% - 9.67%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		14.6%	[12.84% - 16.38%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	[0.48% - 1.06%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments	B	2.2%	[1.70% - 2.74%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees		8.6%	[7.47% - 9.81%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	4.3%	[3.45% - 5.14%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		96.4%	[94.49% - 98.34%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		4.5%	[1.62% - 7.48%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	7.2	[6.8 - 7.5]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year	*	485	[428 - 542]	572	[566 - 578]
Part D monthly prescription fills per person per year	*	51.7	[49.7 - 53.7]	49.1	[48.9 - 49.3]
Home health visits per year		2.5	[1.9 - 3.0]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		2.0	[1.8 - 2.3]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		241	[207 - 275]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		19.4%	[14.67% - 24.09%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		69	[53 - 85]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	2.1%	[1.49% - 2.68%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		81.9%	[76.86% - 86.95%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		80.3%	[74.88% - 85.73%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu