## Jackson (Carroll)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE		MARGIN OF ERROR <sup>2</sup>	STATE ESTIMATE	MARGIN OF ERROR <sup>2</sup>
POPULATION CHARACTERISTICS						
Age-sex adjusted 1-year mortality rate	В	2.7%	[ 1.6	9% - 3.70% ]	4.1%	[ 4.02% - 4.19%
WELLNESS & PREVENTION						
% 60+ with any physical activity within last month	В	80.6%	[ 75.6	88% - 85.59% ]	74.5%	[ 73.36% - 75.59%
% 60+ met CDC guidelines for muscle-strengthening activity		28.5%	[ 21.3	37% - 35.54% ]	26.0%	[ 24.93% - 27.14%
% 60+ met CDC guidelines for aerobic physical activity		65.4%	[ 57.9	93% - 72.77% ]	58.6%	[ 57.33% - 59.82%
% 60+ met CDC guidelines for both types of physical activities		21.0%	[ 14.5	50% - 27.51% ]	19.9%	[ 18.92% - 20.93%
% 60+ getting recommended hours of sleep		68.4%	[ 62.6	62% - 74.25% ]	66.4%	[ 65.16% - 67.60%
% 60+ injured in a fall within last 12 months		11.1%	[ 6.9	95% - 15.22% ]	10.4%	[ 9.65% - 11.16%
% 65+ had hip fracture	В	2.7%	[ 2.7	4% - 3.19% ]	3.3%	[ 3.19% - 3.34%
% 60+ with self-reported fair or poor health status		14.9%	[ 10.6	6% - 19.18% ]	16.5%	[ 15.54% - 17.40%
% 60+ with 15+ physically unhealthy days last month		12.6%	[ 8.7	79% - 16.50% ]	12.5%	[ 11.69% - 13.37%
% 60+ with physical exam/check-up in past year		82.7%	[ 77.2	25% - 88.19% ]	86.5%	[ 85.66% - 87.39%
% 60+ met CDC preventive health screening goals		39.7%	[ 32.4	1% - 47.03% ]	40.3%	[ 38.77% - 41.81%
% 60+ flu shot past year		58.8%	[ 51.8	37% - 65.65% ]	59.3%	[ 58.06% - 60.63%
% 65+ with pneumonia vaccine		72.5%	[ 65.′	4% - 79.91% ]	77.8%	[ 76.56% - 79.12%
% 60+ with cholesterol screening		92.7%	[ 88.6	6% - 96.64% ]	95.3%	[ 94.78% - 95.82%
% 60+ women with a mammogram within last 2 years		78.4%	[ 71.5	52% - 85.33% ]	79.3%	[ 77.96% - 80.69%
% 60+ with colorectal cancer screening		73.0%	[ 66.3	37% - 79.71% ]	77.0%	[ 75.92% - 78.16%
% 60+ with HIV test		11.4%	[ 6.9	96% - 15.84% ]	13.5%	[ 12.50% - 14.41%
% 60+ current smokers		6.6%	[ 3.6	3% - 9.51% ]	8.1%	[ 7.38% - 8.79%
Oral Health						
% 60+ with loss of 6 or more teeth		29.4%	[ 23.2	27% - 35.48% ]	29.0%	[ 27.79% - 30.12%
% 60+ with annual dental exam		76.4%	[ 70.9	95% - 81.81% ]	75.7%	[ 74.65% - 76.84%
NUTRITION/DIET						
% 60+ with 5 or more servings of fruit or vegetables per day		17.8%	[ 12.0	6% - 23.57% ]	20.2%	[ 19.23% - 21.21%
% 60+ self-reported obese		25.9%	[ 19.2	27% - 32.58% ]	27.2%	[ 26.03% - 28.35%
% 65+ clinically diagnosed obese	В	10.2%	[ 8.2	7% - 12.27% ]	16.7%	[ 16.56% - 16.94%
% 65+ with high cholesterol		69.4%	[ 66.3	33% - 72.39% ]	72.2%	[ 72.01% - 72.44%
% 60+ excessive drinking		12.3%	[ 8.0	3% - 16.50% ]	9.2%	[ 8.45% - 9.94%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE	MARGIN OF STATE ERROR <sup>2</sup> ESTIMATE	MARGIN OF ERROR <sup>2</sup>
BEHAVIORAL HEALTH				
% 60+ with 15+ days poor mental health last month	В	3.8%	1.61% - 5.94% ] 6.9% [ 6.2	29% - 7.61%]
% 65+ with depression	В	23.4%	20.73% - 26.10% ] 28.8% [ 28.8	59% - 29.01%]
% 65+ with anxiety disorders	В	14.1%	11.82% - 16.41% ] 21.9% [ 21.7	74% - 22.15% ]
% 65+ with bipolar disorders	В	1.8%	1.28% - 2.25% ] 3.1% [ 2.9	97% - 3.14% ]
% 65+ with post-traumatic stress disorder		1.4%	0.95% - 1.81% ] 1.4% [ 1.3	35% - 1.48% ]
% 65+ with schizophrenia & other psychotic disorders	В	3.9%	3.26% - 4.63% ] 4.9% [ 4.8	31% - 5.02% ]
% 65+ with personality disorders	В	0.7%	0.37% - 0.99% ] 1.1% [ 1.0	00% - 1.11% ]
% 65+ with substance use disorders (drug use +/or alcohol abuse)		4.9%	3.44% - 6.39% ] 5.5% [ 5.4	41% - 5.64% ]
% 65+ with tobacco use disorders	В	7.6%	5.78% - 9.34% ] 10.4% [ 10.2	23% - 10.54% ]
CHRONIC DISEASE				
% 65+ with Alzheimer's disease or related dementias	В	7.4%	5.77% - 9.04% ] 12.0% [ 11.8	39% - 12.19% ]
% 65+ with diabetes	В	19.3%		- 97% - 28.41% ]
% 65+ with stroke	В	7.3%		64% - 10.92% ]
% 65+ with chronic obstructive pulmonary disease		18.8%		36% - 20.73% ]
% 65+ with asthma		11.1%	9.16% - 13.12% ] 13.0% [ 12.8	39% - 13.21% ]
% 65+ with hypertension	В	65.7%	62.55% - 68.76% ] 70.2% [ 70.0	02% - 70.46% ]
% 65+ ever had a heart attack		3.8%	2.66% - 5.01% ] 4.5% [ 4.3	37% - 4.56% ]
% 65+ with ischemic heart disease	В	27.3%	24.29% - 30.22% ] 34.3% [ 34.7	12% - 34.58%]
% 65+ with congestive heart failure	В	14.0%	11.74% - 16.17% ] 17.7% [ 17.5	57% - 17.93%]
% 65+ with atrial fibrillation		15.8%	13.50% - 18.07% ] 14.4% [ 14.2	28% - 14.60%]
% 65+ with peripheral vascular disease	В	11.2%	9.21% - 13.25% ] 14.7% [ 14.5	51% - 14.84%]
% 65+ with osteoarthritis/rheumatoid arthritis	W	53.5%	50.05% - 56.97% ] 49.1% [ 48.8	32% - 49.32% ]
% 65+ with osteoporosis	В	12.0%	9.99% - 14.02% ] 17.4% [ 17.2	22% - 17.57% ]
% 65+ with leukemias and lymphomas		1.5%	1.11% - 1.98% ] 2.0% [ 1.9	91% - 2.05% ]
% 65+ with lung cancer	В	1.1%	0.74% - 1.43% ] 1.6% [ 1.5	54% - 1.66% ]
% 65+ with colon cancer		2.7%	2.13% - 3.22% ] 2.4% [ 2.3	37% - 2.51% ]
% 65+ women with breast cancer		9.8%	7.14% - 12.46% ] 9.8% [ 9.8	59% - 9.96% ]
% 65+ women with endometrial cancer		1.9%	1.25% - 2.52% ] 1.7% [ 1.6	60% - 1.76% ]
% 65+ men with prostate cancer		12.3%	9.40% - 15.24% ] 11.5% [ 11.2	24% - 11.68% ]
% 65+ with benign prostatic hyperplasia		38.6%		47% - 37.14%]
% 65+ with HIV/AIDS		0.04%	0.02% - 0.06% ] 0.05% [ 0.0	04% - 0.06% ]
% 65+ with hypothyroidism	В	16.9%	14.53% - 19.27% ] 20.8% [ 20.6	62% - 20.99% ]
% 65+ with anemia	В	31.6%	28.64% - 34.53% ] 37.3% [ 37.	12% - 37.57% ]
% 65+ with chronic kidney disease	В	13.2%		14% - 22.54% ]
% 65+ with liver diseases	В	3.5%		75% - 6.99% ]
% 65+ with fibromyalgia, chronic pain and fatigue	В	15.2%		42% - 18.80% ]
% 65+ with migraine and other chronic headache		3.6%		95% - 4.15% ]
% 65+ with epilepsy	В	1.3%		00% - 2.14% ]
% 65+ with traumatic brain injury		1.3%	0.90% - 1.67% ] 1.1% [ 1.0	06% - 1.16% ]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE <sup>1</sup>	COMMUNITY ESTIMATE		MARGIN OF ERROR <sup>2</sup>	STATE ESTIMATE	MARGIN OF ERROR <sup>2</sup>
% 65+ with autism spectrum disorders				N/A	0.03%	[ 0.02% - 0.04%
% 65+ with glaucoma	W	33.1%	[	30.13% - 36.10% ]	22.9%	[ 22.74% - 23.13%]
% 65+ with cataract		57.9%	[	54.72% - 61.16% ]	61.2%	[ 61.00% - 61.47%
% 65+ with pressure ulcer or chronic ulcer		6.1%	[	4.58% - 7.65% ]	7.1%	[ 6.99% - 7.24%
% 65+ with 4+ (out of 15) chronic conditions	В	48.5%	[	45.23% - 51.67% ]	54.4%	[ 54.19% - 54.66%
% 65+ with 0 chronic conditions		10.7%	[	8.62% - 12.80% ]	10.3%	[ 10.12% - 10.42%
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment	В	10.5%	[	8.59% - 12.48% ]	14.4%	[ 14.23% - 14.56%
% 65+ with clinical diagnosis of blindness or visual impairment	В	0.5%	[	0.24% - 0.71% ]	0.9%	[ 0.81% - 0.90%
% 65+ with clinical diagnosis of mobility impairments		2.8%	[	2.19% - 3.31% ]	3.2%	[ 3.14% - 3.31%
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees		8.0%	[	6.46% - 9.56% ]	7.9%	[ 7.8% - 8.02%
% dually eligible for Medicare and Medicaid	*	3.5%	[	2.49% - 4.59% ]	7.5%	[ 7.40% - 7.62%
% 60+ with a regular doctor		94.2%	[	90.66% - 97.67% ]	95.8%	[ 95.34% - 96.33%
% 60+ who did not see doctor when needed due to cost		6.0%	[	3.00% - 8.93% ]	5.4%	[ 4.75% - 5.97%
SERVICE UTILIZATION						
Physician visits per year	*	3.5	[	3.3 - 3.8 ]	6.3	[ 6.2 - 6.3
Emergency room visits/1000 persons 65+ years per year		522	[	445 - 599 ]	572	[ 566 - 578
Part D monthly prescription fills per person per year	*	41.1	[	38.6 - 43.6 ]	49.1	[ 48.9 - 49.3
Home health visits per year		2.1	[	1.4 - 2.7 ]	2.5	[ 2.4 - 2.5
Durable medical equipment claims per year		1.7	[	1.3 - 2.0 ]	2.0	[ 2.0 - 2.0
Inpatient hospital stays/1000 persons 65+ years per year		202	[	162 - 241 ]	237	[ 234 - 241
Medicare inpatient hospital readmissions (as % of admissions)		15.8%	[	10.28% - 21.36% ]	15.8%	[ 15.41% - 16.22%
# skilled nursing facility stays/1000 persons 65+ years per year		73	[	48 - 98 ]	76	[ 74 - 77
% 65+ getting Medicaid long term services and supports	*	0.8%	[	0.32% - 1.36% ]	3.7%	[ 3.59% - 3.74%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month		80.0%	[	74.84% - 85.09% ]	77.6%	[ 76.53% - 78.64%
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt		75.0%	[	69.20% - 80.87% ]	77.1%	[ 76.02% - 78.18%
-			-	-		

Notes. <sup>1</sup> Community-state differences where the margin of error do not overlap each other are noted B, W, or \*.
<sup>2</sup> All community and state margins of error are 95% confidence intervals.
<sup>3</sup> The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu