

Litchfield (Hillsborough)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		4.8%	[3.50% - 6.15%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		73.1%	[67.04% - 79.07%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		27.7%	[21.54% - 33.79%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		51.4%	[44.66% - 58.09%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		19.6%	[14.20% - 25.02%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		64.5%	[57.61% - 71.35%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		12.0%	[7.30% - 16.66%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		2.7%	[2.23% - 3.19%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		13.5%	[8.99% - 18.00%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		9.6%	[5.82% - 13.34%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		88.7%	[84.64% - 92.74%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		40.5%	[31.99% - 48.97%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		59.4%	[52.51% - 66.21%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		76.3%	[68.86% - 83.71%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		96.4%	[94.00% - 98.71%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		83.3%	[77.32% - 89.28%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		79.8%	[74.67% - 85.00%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		18.5%	[12.02% - 25.02%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		5.9%	[3.19% - 8.64%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		23.9%	[18.20% - 29.59%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		76.5%	[70.73% - 82.25%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		19.6%	[14.26% - 25.04%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese	W	35.2%	[28.39% - 41.94%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		18.0%	[15.35% - 20.67%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	W	77.2%	[74.41% - 79.97%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		8.9%	[4.68% - 13.17%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		5.4% [2.56% - 8.26%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		27.9% [25.08% - 30.76%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders		23.3% [20.39% - 26.23%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders		3.4% [2.81% - 4.01%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder		1.5% [1.08% - 2.01%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders		5.4% [4.69% - 6.13%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders	B	0.4% [0.23% - 0.67%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		5.2% [3.65% - 6.72%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		8.7% [6.74% - 10.62%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias		13.3% [11.20% - 15.49%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes		28.5% [25.42% - 31.65%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		10.6% [8.67% - 12.49%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease		19.4% [16.91% - 21.85%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		14.3% [12.11% - 16.55%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension		70.9% [67.91% - 73.96%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack		3.5% [2.37% - 4.57%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease		33.0% [29.82% - 36.20%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure		17.4% [14.89% - 19.97%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		13.6% [11.44% - 15.69%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease		16.4% [14.02% - 18.85%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis		48.0% [44.47% - 51.44%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis		18.2% [15.85% - 20.61%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		1.9% [1.45% - 2.38%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.7% [1.33% - 2.15%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.3% [1.81% - 2.74%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		9.1% [6.65% - 11.46%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		2.0% [1.42% - 2.64%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		13.1% [9.89% - 16.34%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia		35.6% [30.99% - 40.16%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.04% [0.02% - 0.07%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism		20.2% [17.69% - 22.75%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia		37.6% [34.55% - 40.73%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease	W	26.4% [23.44% - 29.41%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		7.4% [5.56% - 9.21%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		17.5% [14.92% - 20.03%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		4.4% [3.68% - 5.09%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy		2.0% [1.49% - 2.44%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury	B	0.6% [0.38% - 0.91%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		22.2%	[19.63% - 24.86%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		59.1%	[55.90% - 62.37%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer	B	5.4%	[3.98% - 6.79%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions		55.9%	[52.69% - 59.18%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		9.2%	[7.20% - 11.17%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		13.7%	[11.45% - 15.90%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.6%	[0.38% - 0.86%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.3%	[2.69% - 3.93%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees		8.9%	[7.31% - 10.42%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid		6.3%	[4.97% - 7.63%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		94.4%	[91.18% - 97.56%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost	B	1.8%	[0.52% - 3.00%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	7.2	[6.8 - 7.7]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year	*	445	[382 - 508]	572	[566 - 578]
Part D monthly prescription fills per person per year	*	52.3	[49.4 - 55.3]	49.1	[48.9 - 49.3]
Home health visits per year	*	1.8	[1.2 - 2.4]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		1.9	[1.6 - 2.2]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		204	[166 - 242]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		14.7%	[9.47% - 19.97%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		64	[44 - 84]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	5.1%	[3.86% - 6.25%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month	*	85.1%	[81.04% - 89.13%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		79.9%	[74.58% - 85.16%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu