

Barrington (Strafford)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		3.2%	[2.05% - 4.44%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		77.2%	[71.88% - 82.42%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		25.4%	[20.70% - 30.08%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		61.9%	[56.78% - 67.11%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		21.6%	[17.30% - 25.99%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		69.3%	[63.77% - 74.86%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		12.2%	[8.20% - 16.12%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		2.8%	[2.29% - 3.32%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		13.4%	[9.77% - 17.03%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		9.8%	[6.72% - 12.96%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		90.0%	[86.69% - 93.30%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		44.3%	[37.40% - 51.26%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		61.6%	[55.70% - 67.48%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		79.8%	[74.25% - 85.27%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening	B	98.6%	[97.60% - 99.62%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		78.2%	[71.82% - 84.65%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening	B	83.6%	[79.65% - 87.51%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		10.9%	[6.80% - 14.92%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		6.4%	[3.67% - 9.06%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		26.9%	[21.43% - 32.45%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		80.0%	[74.86% - 85.24%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	[18.16% - 27.28%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		24.4%	[18.97% - 29.90%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		19.1%	[16.45% - 21.82%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol		73.2%	[70.32% - 76.05%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		10.6%	[7.04% - 14.23%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		6.9% [4.15% - 9.70%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		29.5% [26.61% - 32.35%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders		22.8% [19.96% - 25.63%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders	B	2.3% [1.78% - 2.85%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder		1.4% [0.97% - 1.86%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders	B	3.4% [2.78% - 3.98%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders	B	0.3% [0.12% - 0.51%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)	B	4.0% [2.66% - 5.37%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		9.8% [7.80% - 11.84%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias		10.1% [8.22% - 12.02%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes		28.0% [24.94% - 31.06%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		9.2% [7.46% - 10.97%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease		19.3% [16.90% - 21.79%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		11.3% [9.35% - 13.30%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension		67.9% [64.87% - 70.90%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack	B	3.2% [2.11% - 4.24%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease		31.8% [28.67% - 34.96%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure	B	14.8% [12.47% - 17.14%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		12.4% [10.35% - 14.43%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease	B	12.4% [10.30% - 14.49%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis		46.6% [43.16% - 50.02%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis	B	12.7% [10.64% - 14.73%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		1.8% [1.31% - 2.20%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.5% [1.08% - 1.86%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.2% [1.75% - 2.69%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		7.8% [5.44% - 10.13%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.4% [0.87% - 1.93%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		8.8% [6.28% - 11.33%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia		34.8% [30.50% - 39.05%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.04% [0.02% - 0.06%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism	B	16.8% [14.48% - 19.17%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia	B	28.9% [26.07% - 31.71%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease		21.3% [18.58% - 24.00%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		6.7% [5.05% - 8.39%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		17.6% [15.08% - 20.18%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		3.5% [2.87% - 4.19%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy		2.1% [1.55% - 2.59%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury		1.5% [1.11% - 1.90%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma	B	18.0%	[15.61% - 20.41%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	B	53.1%	[49.91% - 56.29%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		6.5%	[4.81% - 8.09%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	B	50.9%	[47.74% - 54.11%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		12.2%	[10.09% - 14.40%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		12.9%	[10.64% - 15.07%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	[0.48% - 1.05%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		2.8%	[2.20% - 3.31%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees	*	4.5%	[3.31% - 5.65%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid		6.1%	[4.79% - 7.50%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor	B	98.2%	[97.04% - 99.35%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		4.2%	[1.81% - 6.62%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year		6.6	[6.1 - 7.0]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year		505	[437 - 572]	572	[566 - 578]
Part D monthly prescription fills per person per year		48.8	[45.9 - 51.8]	49.1	[48.9 - 49.3]
Home health visits per year		2.2	[1.5 - 2.8]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		1.7	[1.4 - 2.0]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		225	[187 - 263]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		11.7%	[6.97% - 16.45%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year	*	46	[30 - 62]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports		3.1%	[2.10% - 4.04%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		80.1%	[75.21% - 85.02%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt	B	83.1%	[78.77% - 87.48%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu