New Hampton (Belknap)



POPULATION CHARACTERISTICS Age-sex adjusted 1-year mortality rate VELLNESS & PREVENTION 6 60+ with any physical activity within last month 6 60+ met CDC guidelines for muscle-strengthening activity 6 60+ met CDC guidelines for aerobic physical activity	3.7%	[
VELLNESS & PREVENTION 6 60+ with any physical activity within last month 6 60+ met CDC guidelines for muscle-strengthening activity		[
6 60+ with any physical activity within last month 6 60+ met CDC guidelines for muscle-strengthening activity	71.7%		2.77% - 4.57%]	4.1%	[4.02% - 4.19%
6 60+ met CDC guidelines for muscle-strengthening activity	71.7%					
		[66.07% - 77.43%]	74.5%	[73.36% - 75.59%
6 60+ met CDC guidelines for aerobic physical activity	21.8%	[16.28% - 27.37%]	26.0%	[24.93% - 27.14%
	56.3%	[49.68% - 63.00%]	58.6%	[57.33% - 59.82%
6 60+ met CDC guidelines for both types of physical activities	16.7%	[11.81% - 21.52%]	19.9%	[18.92% - 20.93%
60+ getting recommended hours of sleep	58.8%	[52.38% - 65.29%]	66.4%	[65.16% - 67.60%
6 60+ injured in a fall within last 12 months	9.6%	[5.91% - 13.28%]	10.4%	[9.65% - 11.16%
65+ had hip fracture	3.0%	[2.41% - 3.57%]	3.3%	[3.19% - 3.34%
60+ with self-reported fair or poor health status	17.8%	[12.95% - 22.68%]	16.5%	[15.54% - 17.40%
60+ with 15+ physically unhealthy days last month	12.4%	[7.88% - 17.01%]	12.5%	[11.69% - 13.37%
60+ with physical exam/check-up in past year	90.8%	[87.20% - 94.31%]	86.5%	[85.66% - 87.39%
60+ met CDC preventive health screening goals	38.0%	[30.68% - 45.23%]	40.3%	[38.77% - 41.81%
60+ flu shot past year	58.8%	[52.42% - 65.27%]	59.3%	[58.06% - 60.63%
65+ with pneumonia vaccine	77.1%	[71.00% - 83.26%]	77.8%	[76.56% - 79.12%
60+ with cholesterol screening	93.2%	[89.86% - 96.51%]	95.3%	[94.78% - 95.82%
60+ women with a mammogram within last 2 years	76.5%	[69.05% - 84.05%]	79.3%	[77.96% - 80.69%
60+ with colorectal cancer screening W	69.3%	[63.26% - 75.44%]	77.0%	[75.92% - 78.16%
60+ with HIV test	9.2%	[5.34% - 13.06%]	13.5%	[12.50% - 14.41%
60+ current smokers	11.5%	[7.07% - 15.90%]	8.1%	[7.38% - 8.79%
Oral Health						
% 60+ with loss of 6 or more teeth	31.2%	[25.44% - 36.98%]	29.0%	[27.79% - 30.12%
% 60+ with annual dental exam	68.9%	[62.99% - 74.71%]	75.7%	[74.65% - 76.84%
IUTRITION/DIET						
6 60+ with 5 or more servings of fruit or vegetables per day	16.8%	[11.90% - 21.75%]	20.2%	[19.23% - 21.21%
60+ self-reported obese	27.5%	[21.58% - 33.39%]	27.2%	[26.03% - 28.35%
65+ clinically diagnosed obese B	12.3%	[10.54% - 13.99%]	16.7%	[16.56% - 16.94%
65+ with high cholesterol	70.6%	[68.28% - 72.84%]	72.2%	[72.01% - 72.44%
60+ excessive drinking	11.0%	[6.81% - 15.22%]	9.2%	[8.45% - 9.94%

New Hampton (Belknap) Page 1

% 65+ with depression B 24.5% (2 2.41% 2.657%) 28.8% (2 8.59% 2.201% 2.01% 3.65% 3.1% 1.21% 2.215% 3.21% 2.174% 2.215% 3.65% with anxiety disorders B 18.1% 16.14% 2.008% 2.15% 3.1% 2.174% 2.215% 3.1% 2.174% 2.215% 3.65% with bipolar disorders B 2.0% 1.48% 2.53% 3.1% 2.174% 2.15% 3.1% 3.1% 2.174% 2.215% 3.65% with bipolar disorders B 2.0% 2.27% 1.60% 1.60% 1.4% 1.35% 1.48% 3.62% 3.47% 4.9% 4.81% 5.02% 6.54 with schizophrenia & other psychotic disorders B 2.9% 2.27% 3.47% 4.9% 4.9% 1.10% 1.10% 1.10% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.02% 1.05% 1.00% 1.02% 1.05% 1.00% 1.02% 1.05% 1.00% 1.02% 1.05% 1.00% 1.0	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 655+ with depression B 24.5% (2.41% - 26.57%) 28.8% (2.85% - 2.901% % 65+ with parkly disorders B 11.1% 10.14% - 20.08% 21.9% (2.17.4% - 22.15% 665+ with parkly disorders B 2.0% (1.48% - 2.53%) 3.1% (2.17.4% - 22.15% % 65+ with parkly bipolar disorders B 2.0% (2.27% - 1.60%) 1.40% (1.35% - 1.48% % 65+ with parkly bipolar disorders B 2.9% (2.27% - 1.60%) 1.4% (1.35% - 1.48% % 65+ with schizophrenia & other psychotic disorders B 0.6% (3.35% - 0.99%) 1.1% (1.10% - 1.11% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% 7.9% 6.46% - 9.36%) 1.0.4 1.0.0% 1.11% % 65+ with liabetes B 7.9% 6.46% - 9.36%) 1.0.4 1.0.0% 1.1.1% % 65+ with Alzheimer's disease or related dementias B 9.7% 6.24% 1.1.07% 1.2.0% 1.1.1% % 65+ with Alzheimer's disease or related dementias B 9.7% 1.8.25% 1.1.0	BEHAVIORAL HEALTH						
% 65+ with anxiety disorders B 18.1% [16.14% - 20.08%] 21.9% 2 1.74% - 22.15% % 65+ with bipolar disorders B 2.0% [1.48% - 2.53%] 3.1% [2.97% - 3.14% % 65+ with post-traumatic stress disorder 1.2% [0.72% - 1.60%] 1.4% [1.35% - 1.48% % 65+ with post-traumatic stress disorders B 2.9% [2.27% - 3.47%] 4.9% [4.81% - 5.02%] % 65+ with post-traumatic stress disorders B 0.6% [0.35% - 0.95%] 1.1% [1.00% - 1.11% 5.64% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% [3.67% - 5.91%] 5.5% [5.41% - 5.64% % 65+ with lobacco use disorders B 7.9% [6.46% - 9.36%] 1.0% [1.12% - 1.15% CHRONIC DISEASE C 8 9.7% [8.24% - 11.07%] 1.2.0% [1.18% - 1.219% % 65+ with Alzheimer's disease or related dementias B 9.7% [8.24% - 11.07%] 1.2.0% [1.18% - 1.20%] 1.1.20% 1.1.20% 1.1.20% 1.1.20% 1.1.20% 1.1.20% 1.1.20% 1.1	% 60+ with 15+ days poor mental health last month		6.9%	[3.47% - 10.23%]	6.9%	[6.29% - 7.61%]
% 65+ with bipolar disorders B 2.0% [1.48% 2.53% 3.1% [2.97% 3.14% % 65+ with post-traumatic stress disorder 1.2% [0.72% 3.0% 1.4% [1.35% 1.48% % 65+ with schizophrenia & other psychotic disorders B 2.9% [0.37% -0.95% 1.1% 1.00% 1.1% 1.00% 1.03% -0.95% 1.1% 1.00% 1.1% 1.00% 1.03% -0.95% 1.04% 1.00% 1.13% 6.64 with schizophrenia & other psychotic disorders B 0.6% 1.03% 0.95% 1.04% 1.00% 1.13% 1.00% 1.13% 6.64 with schizophrenia & other psychotic disorders B 0.9% 6.46% 9.95% 1.04% 1.04% 1.05% <td>% 65+ with depression</td> <td>В</td> <td>24.5%</td> <td>[</td> <td>22.41% - 26.57%]</td> <td>28.8%</td> <td>[28.59% - 29.01%]</td>	% 65+ with depression	В	24.5%	[22.41% - 26.57%]	28.8%	[28.59% - 29.01%]
% 65+ with post-traumatic stress disorder 1.2% [0.72% - 1.60%] 1.4% [1.36% - 1.48%] % 65+ with schizophrenia & other psychotic disorders B 2.9% [2.27% - 3.47%] 4.9% [4.81% - 5.02%] % 65+ with schizophrenia & other psychotic disorders B 0.6% [0.05% - 0.95%] 1.1% [1.00% - 1.11%] % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% [3.67% - 5.91%] 5.5% [5.41% - 5.64%] % 65+ with tobacco use disorders (drug use +/or alcohol abuse) 8 7.9% [6.48% - 3.86%] 10.4% [1.02% - 1.02% - 1.05%] CHRONIC DISEASE 8 9.7% [8.24% - 11.07%] 12.0% [11.89% - 12.19%] % 65+ with diabetes B 24.3% [2.213% - 26.52%] 28.2% [2.797% - 28.41%] % 65+ with diabetes B 15.7% [13.96% 17.39%] 10.0% [10.64% - 10.92%] % 65+ with diabetes B 16.7% [13.96% 17.39%] 10.0% [10.64% - 10.92%] % 65+ with diabetes B 16.7% [13.96% 17.39%] 10.0% [10.64% - 10.92%] % 65+ with directribility distribution of the distribution of the distribution of	% 65+ with anxiety disorders	В	18.1%	[16.14% - 20.08%]	21.9%	[21.74% - 22.15%]
% 65+ with schizophrenia & other psychotic disorders B 2.9% [2.27% 3.47% 4.9% [4.9% 6.48% 5.02% % 65+ with personality disorders B 0.6% [0.35% 0.95% 1.1% 1.1% 1.00% 1.11% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% [3.67% 5.91% 5.5% 5.5% 5.41% 5.564% % 65+ with substance use disorders B 7.9% [6.46% 9.36% 10.4% 10.23% 1.02% 1.054% CHRONIC DISEASE 8 7.9% [6.46% 10.7% 12.0% 12.0% 1.18% 1.2.19% 6.51 1.18% 1.1.19% 12.0% 1.1.89% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.64% 1.0.23% 1.0.24%	% 65+ with bipolar disorders	В	2.0%	[1.48% - 2.53%]	3.1%	[2.97% - 3.14%]
% 65+ with personality disorders B 0.6% 0.35% 0.95% 1.1% 1.00% 1.11% % 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% 1.3.67% 5.91% 5.5% 5.41% 5.64% % 65+ with tobacco use disorders B 7.9% 6.46% 9.36% 10.4% 10.23% 10.54% CHRONIC DISEASE % 65+ with diabetes B 9.7% 8.243% 22.13% 26.52% 22.2% 27.97% 2.841% 65+ with diabetes B 24.3% 22.13% 26.52% 10.6% 10.64% 10.20% 11.89% 10.84% 10.25% 20.26% 20.73% 2.841% 665+ with dispose B 24.3% 22.13% 21.00% 21.19% 61.24% 11.00% 10.0% 11.89% 12.98% 12.19% 665+ with dispose 10.0% 18.56% 11.36% 10.0% 12.19% 655+ with dispose 8 10.5% 18.36% 17.33% 20.5% 20.36% 20.37% 48.41% 49.24%	% 65+ with post-traumatic stress disorder		1.2%	[0.72% - 1.60%]	1.4%	[1.35% - 1.48%]
% 65+ with substance use disorders (drug use +/or alcohol abuse) 4.8% [3.67% - 5.91%] 5.5% [5.41% - 5.64%] % 65+ with tobacco use disorders B 7.9% [6.46% - 9.36%] 10.4% [10.23% - 10.54%] CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.7% [8.24% - 11.07%] 12.0% [11.89% - 12.19%] % 65+ with diabetes B 24.3% [22.13% - 26.52%] 28.2% [27.97% - 28.41%] % 65+ with stroke 10.0% [8.55% - 17.43%] 20.5% [20.36% - 20.73%] 10.92% % 65+ with chronic obstructive pulmonary disease B 15.7% [13.96% - 17.43%] 20.5% [20.36% - 20.73%] % 65+ with chronic obstructive pulmonary disease B 10.8% [9.33% - 12.30%] 13.0% [12.89% - 13.24%] % 65+ with chronic obstructive pulmonary disease B 10.8% [9.38% - 16.60%] 7.02% [70.02% - 70.46%] % 65+ with chronic deart disease B 30.9% [28.59% - 33.27%] 34.3% [12.89% - 13.24%] % 65+ with congestive heart failure B 13.9% [11.74% - 15.66%] 17.7% [17.52% - 17.57%]	% 65+ with schizophrenia & other psychotic disorders	В	2.9%	[2.27% - 3.47%]	4.9%	[4.81% - 5.02%]
% 65+ with tobacco use disorders B 7.9% [6.46% - 9.36%] 10.4% [10.23% - 10.54% CHRONIC DISEASE % 65+ with Alzheimer's disease or related dementias B 9.7% [8.24% - 11.07%] 12.0% [11.89% - 12.19% % 65+ with diabetes B 24.3% [22.13% - 26.52%] 28.2% [27.97% - 28.41% % 65+ with stroke 10.0% [8.55% - 11.36%] 10.68% [10.64% - 10.92% % 65+ with stroke 10.0% [8.55% - 11.36%] 10.68% [10.64% - 10.92% % 65+ with stronic obstructive pulmonary disease B 10.5% [13.96% - 17.3%] 12.0% [12.69% - 20.36% - 20.73% % 65+ with stronic obstructive pulmonary disease B 10.6% [3.39% - 12.30%] 13.0% [12.69% - 20.73% 46.54 with stronic obstructive pulmonary disease B 10.6% [3.89% - 18.60%] 17.0% [10.29% - 10.24% - 20.73% 46.54 with stronic batterial disease B 10.6% [3.89% - 18.60%] 12.0% [10.29%] 12.0% [10.29% - 10.46% 46.54 with stronic batterial disease B 10.2% [28.59% - 33.27%] 34.5% [17.57%] 12.00% [17.59%] 45.5% [17.57%] 46.54 with congestive heart failure B 13.4% [11.71%]	% 65+ with personality disorders	В	0.6%	[0.35% - 0.95%]	1.1%	[1.00% - 1.11%]
## CFRONIC DISEASE 65+ with Alzheimer's disease or related dementias B 9.7% [8.24% 11.07%] 12.0% [11.89% 12.19% 65+ with diabetes B 24.3% [22.13% 26.52%] 28.2% [27.97% 28.41% 65+ with stroke 10.0% [8.55% 11.36%] 10.89% 10.64% 10.92% 65+ with stroke 10.0% [8.55% 11.36%] 10.89% 10.64% 10.92% 65+ with stroke B 15.7% [13.96% 17.43%] 20.5% [20.36% 20.36% 20.73% 65+ with hypertension B 66.2% [63.89% 68.60%] 70.2% 70.02% 70.46% 65+ with hypertension B 66.2% [63.89% 68.60%] 70.2% 70.02% 70.46% 65+ with ischemic heart disease B 30.9% [28.59% 33.27%] 34.3% [34.12% 234.58% 65+ with orangestive heart dislure B 13.4% [11.71% 15.06%] 17.7% [17.57% 17.93% 65+ with strial fibrillation 13.0% [11.41% 13.60%] 14.4% [14.28% 14.60% 65+ with strial disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.51% 14.89% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.28% 14.60% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.28% 14.60% 65+ with benepheral vascular disease B 12.5% [10.89% 14.16%] 14.7% [14.28% 14.60% 1	% 65+ with substance use disorders (drug use +/or alcohol abuse)		4.8%	[3.67% - 5.91%]	5.5%	[5.41% - 5.64%]
% 65+ with Alzheimer's disease or related dementias B 9.7% [8.24% - 11.07%] 12.0% [1.89% - 12.19% % 65+ with diabetes B 24.3% [2.213% - 26.52%] 28.2% [2.797% - 28.41% % 65+ with dribote 10.0% [8.55% - 11.36%] 10.8% [1.064% - 10.92% % 65+ with chronic obstructive pulmonary disease B 15.7% [1.369% - 17.43%] 20.5% [2.036% - 20.73% % 65+ with hypertension B 66.2% [6.389% - 88.60%] 10.02% [7.02% - 70.46% % 65+ with bypertension B 66.2% [6.389% - 83.27%] 34.3% [1.377% - 14.69% % 65+ with stand theart attack 4.1% [3.15% - 5.02%] 4.5% [4.57% - 17.93%] 2.02% 70.46% % 65+ with stand librillation B 30.9% [2.859% - 33.27%] 34.3% [3.175% - 17.93%] 4.5% 4.56% % 65+ with berigheral vascular disease B 30.9% [2.859% - 33.27%] 34.3% [3.15% - 17.59%] 1.47% [1.460% 4.56% % 65+ with berigheral vascular disease B	% 65+ with tobacco use disorders	В	7.9%	[6.46% - 9.36%]	10.4%	[10.23% - 10.54%]
% 65+ with diabetes B 24.3% 22.13% 2 6.52% 28.2% 2.797% 2.844% % 65+ with stroke 10.0% 8.55% 11.36% 1 0.8% 1 0.92% 2.03% 2.073% % 65+ with other onic obstructive pulmonary disease B 15.7% 1 3.96% 1 7.43% 2.05% 2.036% 2.037% % 65+ with hypertension B 66.2% 63.89% 68.60% 7.02% 7.02% 7.046% % 65+ with stylement heart disease B 30.9% 2.559% 33.27% 34.3% [4.57% - 1.73% 4.56% % 65+ with schemic heart disease B 30.9% 2.659% 33.27% 34.3% [3.15% - 1.75% - 1.73% - 4.56% % 65+ with schemic heart disease B 30.9% 2.559% 33.27% 34.3% [3.128% - 3.458% % 65+ with congestive heart failure B 13.3% 1 1.11% - 1.600% 1 1.77% 1 7.75% - 1.757% - 1.793% % 65+ with atrial fibrillation 13.0% 1 1.11% - 1.600% 1 4.79% 1 4.51% 1 4.80% - 4.5193%	CHRONIC DISEASE						
% 65+ with diabetes B 24.3% [22.13% 26.52% 28.2% 27.97% 28.41% 6 28.41% % 65+ with stroke 10.0% 8.55% 11.36% 11.36% 10.8% 10.64% 10.92% 6 10.64% 10.92% % 65+ with chronic obstructive pulmonary disease B 15.7% 13.96% 17.43% 20.5% 20.36% 20.73% % 65+ with asthma B 10.8% 62.2% 63.89% 68.60% 70.2% 70.2% 70.46% 65.4 with strokemic heart attack 4.1% 3.15% 50.2% 4.5% 14.37% 4.56% % 65+ with stromenic heart disease B 30.9% 28.59% 33.27% 34.3% 61.12% 14.57% 17.57% 17.93% % 65+ with stromenic heart disease B 30.9% 28.59% 33.27% 34.3% 17.7% 17.57% 17.93% % 65+ with congestive heart failure B 13.4% 11.11% 15.06% 17.7% 17.57% 17.57% 17.93% % 65+ with pripheral vascular disease B 12.5% 10.89% 14.16% 14.7% 14.51% 14.80% % 65+ with pripheral vascular disease B 12.5% 10.99% 13.09% 14.16% 14.7% 14.51% 14.80% % 65+ with osteoporosis 15.6% 13.29% 17.35% 17.4% 17.57% 17.57% 65+ with leukemias and lymphomas 15.6% 13.29% 17.35% 17.4% 17.5% 17.57% 15.5% % 65+ with lung cancer 12.2% 10.7% 10.79% 10.99% 2.0% 1.61% 1.66% 1.66% 1.66% 1.65% 1.61% 1.60% 1.66% 1.66% 1.66% 1.66% 1.66% 1.66% 1.66% 1.	% 65+ with Alzheimer's disease or related dementias	В	9.7%	-	8.24% - 11.07%]	12.0%	[11.89% - 12.19%]
% 65+ with stroke 10.0% [8.55% - 11.36%] 10.8% [10.64%] - 10.92% % 65+ with chronic obstructive pulmonary disease B 15.7% [13.96%] - 17.43%] 20.5% [20.36%] - 20.73% % 65+ with asthma B 10.8% [9.33%] - 12.30%] 13.0% [1 2.89%] - 32.27% % 65+ with sthma B 66.2% [63.89%] - 68.60%] 70.2% [7 0.02%] - 70.46% % 65+ with sthman B 66.2% [63.89%] - 68.60%] 70.2% [7 0.02%] - 70.46% % 65+ with sthman cheart disease B 30.9% [28.59%] - 33.27%] 34.3% [1 4.75%] - 41.60% % 65+ with congestive heart failure B 13.4% [11.71%] - 15.66%] 17.7% [17.57%] - 70.93% % 65+ with still sibrillation 13.0% [11.41%] - 14.60%] 14.4% [1 4.60%] 44.50% - 41.60% 45.71% - 10.60% - 14.7% [1 4.57%] - 14.60% - 65+ with osteoarthritis/rheumatoid arthritis 48.39 [45.71%] - 50.92%] 49.1% - 14.7% - 14.76%	% 65+ with diabetes	В		i 1		28.2%	[27.97% - 28.41%]
% 65+ with asthma B 10.8% [9.33% - 12.30%] 13.0% [12.89% - 13.21% % 65+ with hypertension B 66.2% [63.89% - 68.60%] 70.2% [70.02% - 70.46% % 65+ with hypertension B 66.2% [63.89% - 68.60%] 70.2% [70.02% - 70.46% % 65+ with ad a heart attack 4.1% [3.15% - 5.02%] 4.5% [4.37% - 4.56% % 65+ with congestive heart failure B 30.9% [28.59% - 33.27%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure B 33.4% [11.71% - 15.06%] 17.7% [17.57% - 17.33% % 65+ with atrial fibrillation 13.0% [11.11% - 14.60%] 14.4% [14.28% - 14.60% % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84% % 65+ with peripheral vascular disease B 12.5% [10.89% - 17.35%] 14.1% [14.87% - 14.84% % 65+ with peripheral vascular disease B 12.5% [10.89% - 17.35%] 14.1% [14.27% - 14.28% % 65+ with peripheral vascular disease 15.6% [1.392% - 17.35%] 14.1% [14.27% -	% 65+ with stroke		10.0%	<u>.</u>]		10.8%	[10.64% - 10.92%]
% 65+ with asthma B 10.8% [9.33% - 12.30%] 13.0% [12.89% - 13.21% % 65+ with hypertension B 66.2% [63.89% - 68.60%] 70.2% [70.02% - 70.46% % 65+ with hypertension B 66.2% [63.89% - 68.60%] 70.2% [70.02% - 70.46% % 65+ with ad a heart attack 4.1% [3.15% - 5.02%] 4.5% [4.37% - 4.56% % 65+ with congestive heart failure B 30.9% [28.59% - 33.27%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure B 33.4% [11.71% - 15.06%] 17.7% [17.57% - 17.33% % 65+ with atrial fibrillation 13.0% [11.11% - 14.60%] 14.4% [14.28% - 14.60% % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84% % 65+ with peripheral vascular disease B 12.5% [10.89% - 17.35%] 14.1% [14.87% - 14.84% % 65+ with peripheral vascular disease B 12.5% [10.89% - 17.35%] 14.1% [14.27% - 14.28% % 65+ with peripheral vascular disease 15.6% [1.392% - 17.35%] 14.1% [14.27% -	% 65+ with chronic obstructive pulmonary disease	В	15.7%		13.96% - 17.43%]	20.5%	[20.36% - 20.73%]
% 65+ ever had a heart attack 4.1% [3.15% - 5.02%] 4.5% [4.37% - 4.56% % 65+ with ischemic heart disease B 30.9% [28.59% - 33.27%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure B 13.4% [11.71% - 15.06%] 17.7% [17.57% - 17.93% % 65+ with congestive heart failure B 13.4% [11.71% - 15.06%] 14.4% [14.28% - 14.60% % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84% % 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32% % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with long cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with prostatic cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with prostate cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ with benign prostatic hyperplasia	% 65+ with asthma	В	10.8%		9.33% - 12.30%]	13.0%	[12.89% - 13.21%]
% 65+ with ischemic heart disease B 30.9% [28.59% - 33.27%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure B 13.4% [11.71% - 15.06%] 17.7% [17.57% - 17.93% % 65+ with atrial fibrillation 13.0% [11.41% - 14.60%] 14.4% [14.28% - 14.60% % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84% % 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32% % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with prostate cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ with hendometrial cancer 12.8% [10.45% - 15.17%] 11.5% [1.24% - 1.68% % 65+ with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [1.60% - 1.76% % 65+ with henign prostatic hyperplasia	% 65+ with hypertension	В	66.2%	[63.89% - 68.60%]	70.2%	[70.02% - 70.46%]
% 65+ with congestive heart failure B 13.4% [11.71% - 15.06%] 17.7% [17.57% - 17.93% % 65+ with atrial fibrillation 13.0% [11.41% - 14.60%] 14.4% [14.28% - 14.60%] % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84%] % 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32%] % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57%] % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05%] % 65+ with colon cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66%] % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96%] % 65+ women with prostate cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76%] % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14%] % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99%] % 65+ with chronic kidney disease	% 65+ ever had a heart attack		4.1%	 [3.15% - 5.02%]	4.5%	[4.37% - 4.56%]
% 65+ with atrial fibrillation 13.0% [11.41% - 14.60%] 14.4% [14.28% - 14.60%] % 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84%] % 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32%] % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57%] % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05%] % 65+ with colon cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66%] % 65+ women with breast cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51%] % 65+ women with prostate cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96%] % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14%] % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99%] % 65+ with chronic kidney disease B 15.0% [14.73% - 14.65%] 10.9% [2.14% - 22.54%] % 65+ with fibromyalgia, chronic pain an	% 65+ with ischemic heart disease	В	30.9%	[28.59% - 33.27%]	34.3%	[34.12% - 34.58%]
% 65+ with peripheral vascular disease B 12.5% [10.89% - 14.16%] 14.7% [14.51% - 14.84% % 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32% % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with lung cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with preast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ women with prostate cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with fibromyalgia, chronic pain and fatigue <th< td=""><td>% 65+ with congestive heart failure</td><td>В</td><td>13.4%</td><td>[</td><td>11.71% - 15.06%]</td><td>17.7%</td><td>[17.57% - 17.93%]</td></th<>	% 65+ with congestive heart failure	В	13.4%	[11.71% - 15.06%]	17.7%	[17.57% - 17.93%]
% 65+ with osteoarthritis/rheumatoid arthritis 48.3% [45.71% - 50.92%] 49.1% [48.82% - 49.32% % 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with lung cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 2.51% % 65+ women with prostate cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with fibromyalgia, chronic pain and fatigue B 15.5% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic head	% 65+ with atrial fibrillation		13.0%	[11.41% - 14.60%]	14.4%	[14.28% - 14.60%]
% 65+ with osteoporosis 15.6% [13.92% - 17.35%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with lung cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with nemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with liver diseases B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9%	% 65+ with peripheral vascular disease	В	12.5%	[10.89% - 14.16%]	14.7%	[14.51% - 14.84%]
% 65+ with leukemias and lymphomas 1.5% [1.07% - 1.99%] 2.0% [1.91% - 2.05% % 65+ with lung cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ men with prostate cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04	% 65+ with osteoarthritis/rheumatoid arthritis		48.3%	[45.71% - 50.92%]	49.1%	[48.82% - 49.32%]
% 65+ with lung cancer 1.2% [0.81% - 1.56%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ women with endometrial cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with briornic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with migraine and other chronic pain and fatigue B 15.8% [14.01% - 1	% 65+ with osteoporosis		15.6%	[13.92% - 17.35%]	17.4%	[17.22% - 17.57%]
% 65+ with colon cancer 2.0% [1.53% - 2.49%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ women with endometrial cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.0%	% 65+ with leukemias and lymphomas		1.5%	[1.07% - 1.99%]	2.0%	[1.91% - 2.05%]
% 65+ women with breast cancer 8.5% [6.72% - 10.37%] 9.8% [9.59% - 9.96% % 65+ women with endometrial cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with migraine and other chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with lung cancer		1.2%	[0.81% - 1.56%]	1.6%	[1.54% - 1.66%]
% 65+ women with endometrial cancer 1.2% [0.70% - 1.74%] 1.7% [1.60% - 1.76% % 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with colon cancer		2.0%	[1.53% - 2.49%]	2.4%	[2.37% - 2.51%]
% 65+ men with prostate cancer 12.8% [10.45% - 15.17%] 11.5% [11.24% - 11.68% % 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ women with breast cancer		8.5%	[6.72% - 10.37%]	9.8%	[9.59% - 9.96%]
% 65+ with benign prostatic hyperplasia 38.0% [34.52% - 41.51%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ women with endometrial cancer		1.2%	[0.70% - 1.74%]	1.7%	[1.60% - 1.76%]
% 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ men with prostate cancer		12.8%	[10.45% - 15.17%]	11.5%	[11.24% - 11.68%]
% 65+ with hypothyroidism B 16.5% [14.73% - 18.31%] 20.8% [20.62% - 20.99% % 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with benign prostatic hyperplasia		38.0%	[34.52% - 41.51%]	36.8%	[36.47% - 37.14%]
% 65+ with anemia 35.6% [33.29% - 37.92%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with HIV/AIDS		0.04%	[0.02% - 0.06%]	0.05%	[0.04% - 0.06%]
% 65+ with chronic kidney disease B 15.0% [13.23% - 16.82%] 22.3% [22.14% - 22.54% % 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with hypothyroidism	В	16.5%	[14.73% - 18.31%]	20.8%	[20.62% - 20.99%]
% 65+ with liver diseases 6.2% [4.96% - 7.43%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with anemia		35.6%	[33.29% - 37.92%]	37.3%	[37.12% - 37.57%]
% 65+ with fibromyalgia, chronic pain and fatigue B 15.8% [14.01% - 17.66%] 18.6% [18.42% - 18.80%] % 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15%] % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%]	% 65+ with chronic kidney disease	В	15.0%	[13.23% - 16.82%]	22.3%	[22.14% - 22.54%]
% 65+ with migraine and other chronic headache 3.9% [3.17% - 4.65%] 4.0% [3.95% - 4.15% % 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with liver diseases		6.2%	[4.96% - 7.43%]	6.9%	[6.75% - 6.99%]
% 65+ with epilepsy 1.6% [1.13% - 2.04%] 2.1% [2.00% - 2.14%	% 65+ with fibromyalgia, chronic pain and fatigue	В	15.8%	[14.01% - 17.66%]	18.6%	[18.42% - 18.80%]
	% 65+ with migraine and other chronic headache		3.9%	[3.17% - 4.65%]	4.0%	[3.95% - 4.15%]
0/ OF 1914 1111	% 65+ with epilepsy		1.6%	[1.13% - 2.04%]	2.1%	[2.00% - 2.14%]
% 65+ with traumatic brain injury 1.2% [0.82% - 1.63%] 1.1% [1.06% - 1.16%	% 65+ with traumatic brain injury		1.2%	[0.82% - 1.63%]	1.1%	[1.06% - 1.16%]

New Hampton (Belknap) Page 2

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders				N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		21.5%	[19.57% - 23.52%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		62.0%	[59.56% - 64.40%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		6.8%	[5.61% - 8.05%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	В	47.7%	[45.21% - 50.10%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		10.1%	[8.58% - 11.69%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment		14.5%	[12.78% - 16.26%]	14.4%	[14.23% - 14.56%
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	[0.45% - 1.06%]	0.9%	[0.81% - 0.90%
% 65+ with clinical diagnosis of mobility impairments	В	2.2%	[1.66% - 2.71%]	3.2%	[3.14% - 3.31%
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees		8.2%	[7.03% - 9.41%]	7.9%	[7.8% - 8.02%
% dually eligible for Medicare and Medicaid	*	4.8%	[3.91% - 5.78%]	7.5%	[7.40% - 7.62%
% 60+ with a regular doctor		93.5%	[90.51% - 96.54%]	95.8%	[95.34% - 96.33%
% 60+ who did not see doctor when needed due to cost		6.5%	[2.85% - 10.24%]	5.4%	[4.75% - 5.97%
SERVICE UTILIZATION						
Physician visits per year		6.5	[6.2 - 6.8]	6.3	[6.2 - 6.3
Emergency room visits/1000 persons 65+ years per year	*	487	[435 - 539]	572	[566 - 578
Part D monthly prescription fills per person per year	*	42.5	[40.4 - 44.6]	49.1	[48.9 - 49.3
Home health visits per year		2.0	[1.5 - 2.5]	2.5	[2.4 - 2.5
Durable medical equipment claims per year		1.8	[1.5 - 2.1]	2.0	[2.0 - 2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	191	[162 - 221]	237	[234 - 241
Medicare inpatient hospital readmissions (as % of admissions)		17.7%	[12.97% - 22.51%]	15.8%	[15.41% - 16.22%
# skilled nursing facility stays/1000 persons 65+ years per year		68	[52 - 84]	76	[74 - 77
% 65+ getting Medicaid long term services and supports	*	2.2%	[1.57% - 2.84%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month	*	70.4%	[64.65% - 76.23%]	77.6%	[76.53% - 78.64%
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt	W	66.1%	[60.08% - 72.19%]	77.1%	[76.02% - 78.18%

Questions or suggestions? Beth.Dugan@umb.edu

New Hampton (Belknap) Page 3

Notes.

Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

All community and state margins of error are 95% confidence intervals.

The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last montth, and did not see doctor when needed due to cost.