

Northumberland (Coös)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		3.1%	[1.51% - 4.65%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		67.2%	[59.93% - 74.47%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity	W	16.7%	[10.53% - 22.92%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity	W	48.1%	[39.96% - 56.20%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities	W	10.2%	[5.54% - 14.78%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		63.1%	[55.93% - 70.26%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		11.8%	[6.91% - 16.79%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		2.9%	[2.13% - 3.60%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		23.6%	[16.61% - 30.59%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		12.6%	[7.38% - 17.89%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		91.0%	[86.36% - 95.55%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		38.5%	[29.69% - 47.31%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		58.7%	[51.09% - 66.33%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		82.1%	[75.87% - 88.38%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		93.1%	[89.30% - 96.81%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		77.1%	[69.44% - 84.77%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		77.9%	[71.98% - 83.82%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		13.4%	[8.26% - 18.61%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		11.2%	[6.05% - 16.39%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth	W	41.9%	[34.89% - 48.96%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam	W	52.3%	[45.21% - 59.38%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		17.0%	[10.67% - 23.25%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		34.2%	[26.42% - 41.98%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		18.3%	[14.51% - 22.12%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	W	77.6%	[73.66% - 81.49%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		11.4%	[6.22% - 16.48%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		7.6% [3.33% - 11.81%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		30.2% [26.10% - 34.40%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders		22.4% [18.53% - 26.37%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders	B	2.2% [1.48% - 2.91%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder		2.1% [1.40% - 2.84%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders	B	3.6% [2.73% - 4.45%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders		1.2% [0.67% - 1.75%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		4.2% [2.39% - 6.10%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders	W	14.3% [10.93% - 17.72%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	8.1% [5.67% - 10.56%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes	W	38.6% [33.82% - 43.32%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		10.6% [7.86% - 13.31%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease	W	25.4% [21.48% - 29.34%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		14.2% [11.10% - 17.35%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension	W	76.5% [72.58% - 80.51%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack	W	7.7% [5.35% - 10.11%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease	W	40.6% [35.90% - 45.39%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure		21.4% [17.53% - 25.21%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		14.7% [11.50% - 17.84%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease	W	18.8% [15.15% - 22.49%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis		47.8% [42.94% - 52.66%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis		15.4% [12.22% - 18.54%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		2.2% [1.46% - 2.88%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.6% [1.01% - 2.13%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.7% [1.92% - 3.38%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		8.7% [5.21% - 12.19%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.2% [0.48% - 1.85%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		9.8% [5.92% - 13.67%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia	B	29.9% [23.80% - 35.98%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.05% [0.02% - 0.08%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism	B	17.1% [13.73% - 20.48%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia		37.8% [33.44% - 42.23%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease	W	27.8% [23.56% - 32.10%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		5.7% [3.56% - 7.92%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		18.4% [14.70% - 22.17%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		3.9% [2.95% - 4.83%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy		1.9% [1.30% - 2.56%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury	B	0.6% [0.28% - 1.01%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		20.3%	[16.70% - 23.91%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		60.6%	[56.01% - 65.12%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		8.5%	[5.81% - 11.10%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	W	59.9%	[55.35% - 64.42%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		8.6%	[5.85% - 11.33%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		14.6%	[11.28% - 17.85%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.6%	[0.23% - 0.90%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.0%	[2.21% - 3.84%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees	*	1.3%	[0.35% - 2.30%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	12.1%	[9.32% - 14.88%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		95.0%	[91.91% - 98.00%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		8.4%	[4.04% - 12.84%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	2.6	[2.2 - 2.9]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year	*	928	[759 - 1097]	572	[566 - 578]
Part D monthly prescription fills per person per year		54.3	[49.2 - 59.4]	49.1	[48.9 - 49.3]
Home health visits per year		3.5	[2.3 - 4.7]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year	*	3.2	[2.4 - 3.9]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year	*	348	[269 - 427]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		19.0%	[12.35% - 25.74%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year	*	174	[119 - 230]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports		4.2%	[2.46% - 5.86%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month	*	60.9%	[53.38% - 68.51%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt	W	67.2%	[60.45% - 73.88%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu