

Rollinsford (Strafford)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		4.5%	[2.39% - 6.67%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		77.2%	[71.88% - 82.42%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		25.4%	[20.70% - 30.08%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		61.9%	[56.78% - 67.11%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		21.6%	[17.30% - 25.99%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		69.3%	[63.77% - 74.86%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		12.2%	[8.20% - 16.12%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture		2.7%	[1.97% - 3.44%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		13.4%	[9.77% - 17.03%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		9.8%	[6.72% - 12.96%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		90.0%	[86.69% - 93.30%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		44.3%	[37.40% - 51.26%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		61.6%	[55.70% - 67.48%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		79.8%	[74.25% - 85.27%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening	B	98.6%	[97.60% - 99.62%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years		78.2%	[71.82% - 84.65%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening	B	83.6%	[79.65% - 87.51%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		10.9%	[6.80% - 14.92%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		6.4%	[3.67% - 9.06%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		26.9%	[21.43% - 32.45%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		80.0%	[74.86% - 85.24%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	[18.16% - 27.28%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		24.4%	[18.97% - 29.90%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		20.8%	[15.98% - 25.71%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol		76.4%	[71.72% - 81.17%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		10.6%	[7.04% - 14.23%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		6.9% [4.15% - 9.70%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		32.0% [27.03% - 37.07%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders		25.7% [20.62% - 30.79%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders		3.8% [2.79% - 4.88%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder	W	2.7% [1.79% - 3.68%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders		6.1% [4.95% - 7.28%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders		1.0% [0.44% - 1.56%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		6.8% [3.74% - 9.84%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		10.5% [6.91% - 14.08%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias		12.7% [9.17% - 16.24%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes		29.7% [24.22% - 35.17%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		11.4% [8.14% - 14.56%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease		22.5% [18.06% - 26.96%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		12.6% [9.08% - 16.19%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension		70.3% [65.18% - 75.33%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack		4.6% [2.42% - 6.76%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease		38.0% [32.35% - 43.72%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure		17.1% [12.68% - 21.53%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		15.5% [11.72% - 19.37%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease		15.0% [10.98% - 18.93%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis		50.1% [44.11% - 56.04%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis		15.9% [12.07% - 19.70%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		2.0% [1.29% - 2.78%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		2.1% [1.43% - 2.79%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.7% [1.91% - 3.41%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		10.6% [6.25% - 14.92%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.7% [0.84% - 2.47%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		9.6% [4.85% - 14.30%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia		40.3% [32.10% - 48.42%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.04% [0.02% - 0.06%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism		20.6% [16.38% - 24.89%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia		37.3% [32.09% - 42.46%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease		26.0% [21.02% - 31.01%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		7.8% [4.81% - 10.70%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue	W	24.8% [19.83% - 29.76%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache		4.4% [3.32% - 5.50%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy		2.7% [1.87% - 3.55%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury		1.6% [0.98% - 2.26%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		27.6%	[22.82% - 32.32%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		56.1%	[50.62% - 61.55%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		8.2%	[5.12% - 11.31%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions		55.2%	[49.74% - 60.64%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		11.8%	[8.17% - 15.52%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		18.6%	[14.39% - 22.81%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		1.2%	[0.64% - 1.78%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.4%	[2.46% - 4.34%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees	*	4.7%	[2.62% - 6.73%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	3.4%	[1.67% - 5.22%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor	B	98.2%	[97.04% - 99.35%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		4.2%	[1.81% - 6.62%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	7.8	[6.9 - 8.6]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year		635	[428 - 842]	572	[566 - 578]
Part D monthly prescription fills per person per year		52.8	[47.3 - 58.4]	49.1	[48.9 - 49.3]
Home health visits per year		2.1	[1.1 - 3.0]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		2.6	[1.8 - 3.3]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		203	[137 - 268]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		19.2%	[9.51% - 28.85%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		48	[17 - 79]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	1.7%	[0.46% - 2.99%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		80.1%	[75.21% - 85.02%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt	B	83.1%	[78.77% - 87.48%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu