

Temple (Hillsborough)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		4.1%	[2.61% - 5.57%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		78.4%	[72.00% - 84.76%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		27.7%	[20.83% - 34.59%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		67.0%	[59.63% - 74.39%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		22.4%	[16.22% - 28.56%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		63.9%	[56.31% - 71.42%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		10.7%	[6.68% - 14.72%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture	W	4.2%	[3.36% - 5.04%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		13.9%	[9.11% - 18.62%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		10.6%	[5.81% - 15.37%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		84.3%	[78.65% - 89.91%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		39.4%	[30.09% - 48.66%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		55.8%	[48.03% - 63.53%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		73.5%	[64.28% - 82.80%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		95.6%	[93.13% - 98.13%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years	W	67.9%	[58.48% - 77.24%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		73.9%	[66.86% - 80.86%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		11.8%	[7.22% - 16.33%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		8.1%	[3.69% - 12.52%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		32.7%	[24.82% - 40.49%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		74.5%	[67.87% - 81.05%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		19.3%	[13.65% - 24.92%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		24.9%	[18.25% - 31.59%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese		16.2%	[13.01% - 19.41%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol	B	66.4%	[62.53% - 70.21%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		5.3%	[2.07% - 8.58%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		5.0% [1.57% - 8.36%]	6.9% [6.29% - 7.61%]		
% 65+ with depression		26.0% [22.52% - 29.42%]	28.8% [28.59% - 29.01%]		
% 65+ with anxiety disorders	B	16.9% [13.79% - 20.02%]	21.9% [21.74% - 22.15%]		
% 65+ with bipolar disorders		2.7% [1.99% - 3.49%]	3.1% [2.97% - 3.14%]		
% 65+ with post-traumatic stress disorder		1.1% [0.55% - 1.58%]	1.4% [1.35% - 1.48%]		
% 65+ with schizophrenia & other psychotic disorders		4.0% [3.11% - 4.81%]	4.9% [4.81% - 5.02%]		
% 65+ with personality disorders		1.0% [0.52% - 1.51%]	1.1% [1.00% - 1.11%]		
% 65+ with substance use disorders (drug use +/- alcohol abuse)		4.9% [2.97% - 6.82%]	5.5% [5.41% - 5.64%]		
% 65+ with tobacco use disorders		9.2% [6.75% - 11.62%]	10.4% [10.23% - 10.54%]		
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	8.5% [6.44% - 10.59%]	12.0% [11.89% - 12.19%]		
% 65+ with diabetes	B	21.5% [18.00% - 25.00%]	28.2% [27.97% - 28.41%]		
% 65+ with stroke		8.6% [6.49% - 10.64%]	10.8% [10.64% - 10.92%]		
% 65+ with chronic obstructive pulmonary disease		18.7% [15.66% - 21.73%]	20.5% [20.36% - 20.73%]		
% 65+ with asthma		11.1% [8.61% - 13.55%]	13.0% [12.89% - 13.21%]		
% 65+ with hypertension	B	57.7% [53.72% - 61.73%]	70.2% [70.02% - 70.46%]		
% 65+ ever had a heart attack		4.7% [3.09% - 6.23%]	4.5% [4.37% - 4.56%]		
% 65+ with ischemic heart disease	B	27.6% [23.94% - 31.33%]	34.3% [34.12% - 34.58%]		
% 65+ with congestive heart failure	B	14.7% [11.92% - 17.54%]	17.7% [17.57% - 17.93%]		
% 65+ with atrial fibrillation		12.3% [9.83% - 14.82%]	14.4% [14.28% - 14.60%]		
% 65+ with peripheral vascular disease	B	11.0% [8.56% - 13.41%]	14.7% [14.51% - 14.84%]		
% 65+ with osteoarthritis/rheumatoid arthritis		50.3% [46.03% - 54.63%]	49.1% [48.82% - 49.32%]		
% 65+ with osteoporosis		16.3% [13.48% - 19.09%]	17.4% [17.22% - 17.57%]		
% 65+ with leukemias and lymphomas		1.8% [1.12% - 2.42%]	2.0% [1.91% - 2.05%]		
% 65+ with lung cancer		1.3% [0.79% - 1.79%]	1.6% [1.54% - 1.66%]		
% 65+ with colon cancer		2.0% [1.41% - 2.60%]	2.4% [2.37% - 2.51%]		
% 65+ women with breast cancer		8.8% [5.74% - 11.86%]	9.8% [9.59% - 9.96%]		
% 65+ women with endometrial cancer		1.3% [0.62% - 1.88%]	1.7% [1.60% - 1.76%]		
% 65+ men with prostate cancer		8.7% [5.49% - 11.87%]	11.5% [11.24% - 11.68%]		
% 65+ with benign prostatic hyperplasia		34.2% [28.74% - 39.69%]	36.8% [36.47% - 37.14%]		
% 65+ with HIV/AIDS		0.04% [0.02% - 0.07%]	0.05% [0.04% - 0.06%]		
% 65+ with hypothyroidism	B	16.4% [13.51% - 19.24%]	20.8% [20.62% - 20.99%]		
% 65+ with anemia		34.7% [30.97% - 38.46%]	37.3% [37.12% - 37.57%]		
% 65+ with chronic kidney disease		19.2% [15.99% - 22.38%]	22.3% [22.14% - 22.54%]		
% 65+ with liver diseases		6.4% [4.37% - 8.49%]	6.9% [6.75% - 6.99%]		
% 65+ with fibromyalgia, chronic pain and fatigue		18.0% [14.80% - 21.21%]	18.6% [18.42% - 18.80%]		
% 65+ with migraine and other chronic headache	W	5.4% [4.38% - 6.50%]	4.0% [3.95% - 4.15%]		
% 65+ with epilepsy		2.2% [1.47% - 2.86%]	2.1% [2.00% - 2.14%]		
% 65+ with traumatic brain injury		1.0% [0.55% - 1.38%]	1.1% [1.06% - 1.16%]		

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma	W	28.4%	[24.87% - 31.90%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	W	67.7%	[63.82% - 71.57%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		6.5%	[4.56% - 8.50%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	B	45.3%	[41.29% - 49.22%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		12.8%	[9.98% - 15.66%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		16.4%	[13.46% - 19.39%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.8%	[0.41% - 1.18%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.0%	[2.26% - 3.76%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees	*	5.5%	[3.94% - 7.12%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	4.4%	[2.97% - 5.82%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		95.7%	[92.91% - 98.41%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		5.1%	[1.42% - 8.82%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	4.9	[4.5 - 5.3]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year		530	[424 - 635]	572	[566 - 578]
Part D monthly prescription fills per person per year	*	42.6	[39.1 - 46.1]	49.1	[48.9 - 49.3]
Home health visits per year		2.3	[1.2 - 3.5]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		1.8	[1.4 - 2.3]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		200	[154 - 245]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		11.7%	[5.42% - 18.02%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		63	[40 - 86]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	1.6%	[0.75% - 2.51%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		81.8%	[76.00% - 87.54%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		78.8%	[72.95% - 84.59%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu