Walpole (Cheshire)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²		STATE ESTIMATE	MARGIN OF ERROR ²	
POPULATION CHARACTERISTICS							
Age-sex adjusted 1-year mortality rate	В	2.8%	[1.66% - 3.98%] 4.1%	[4.02% - 4.199	
WELLNESS & PREVENTION							
% 60+ with any physical activity within last month		77.1%	[71.88% - 82.37%] 74.5%	[73.36% - 75.599	
% 60+ met CDC guidelines for muscle-strengthening activity		25.6%	[20.21% - 30.99%] 26.0%	[24.93% - 27.149	
% 60+ met CDC guidelines for aerobic physical activity		62.7%	[56.76% - 68.67%] 58.6%	[57.33% - 59.829	
% 60+ met CDC guidelines for both types of physical activities		21.0%	[16.05% - 26.00%] 19.9%	[18.92% - 20.939	
% 60+ getting recommended hours of sleep		65.4%	[59.50% - 71.24%] 66.4%	[65.16% - 67.60	
% 60+ injured in a fall within last 12 months		8.0%	[4.71% - 11.25%] 10.4%	[9.65% - 11.169	
% 65+ had hip fracture	В	2.5%	[1.89% - 3.07%] 3.3%	[3.19% - 3.349	
% 60+ with self-reported fair or poor health status		18.1%	[13.14% - 23.05%] 16.5%	[15.54% - 17.409	
% 60+ with 15+ physically unhealthy days last month		13.1%	[8.54% - 17.68%] 12.5%	[11.69% - 13.379	
% 60+ with physical exam/check-up in past year	W	79.2%	[74.09% - 84.34%] 86.5%	[85.66% - 87.39	
% 60+ met CDC preventive health screening goals		39.6%	[32.26% - 46.93%] 40.3%	[38.77% - 41.819	
% 60+ flu shot past year		58.1%	[51.87% - 64.30%] 59.3%	[58.06% - 60.639	
% 65+ with pneumonia vaccine		74.2%	[67.79% - 80.57%] 77.8%	[76.56% - 79.129	
% 60+ with cholesterol screening		95.0%	[92.51% - 97.48%] 95.3%	[94.78% - 95.829	
% 60+ women with a mammogram within last 2 years		75.0%	[68.54% - 81.41%] 79.3%	[77.96% - 80.699	
% 60+ with colorectal cancer screening		71.4%	[65.66% - 77.21%] 77.0%	[75.92% - 78.169	
% 60+ with HIV test		16.7%	[11.75% - 21.63%] 13.5%	[12.50% - 14.419	
% 60+ current smokers		7.2%	[4.18% - 10.20%] 8.1%	[7.38% - 8.79	
Oral Health							
% 60+ with loss of 6 or more teeth		29.6%	[24.15% - 35.05%] 29.0%	[27.79% - 30.129	
% 60+ with annual dental exam	W	66.1%	[60.24% - 71.93%] 75.7%	[74.65% - 76.849	
NUTRITION/DIET							
% 60+ with 5 or more servings of fruit or vegetables per day		22.2%	[17.09% - 27.23%] 20.2%	[19.23% - 21.219	
% 60+ self-reported obese		31.4%	[25.24% - 37.53%] 27.2%	[26.03% - 28.359	
% 65+ clinically diagnosed obese	В	13.6%	[10.75% - 16.49%] 16.7%	[16.56% - 16.949	
% 65+ with high cholesterol	В	62.9%]	59.12% - 66.76%] 72.2%	[72.01% - 72.449	
% 60+ excessive drinking		10.1%	[6.62% - 13.64%] 9.2%	[8.45% - 9.949	

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		RGIN OF RROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH						
% 60+ with 15+ days poor mental health last month		9.9%	[5.909	% - 13.95%]	6.9%	[6.29% - 7.61%]
% 65+ with depression		26.2%	[22.839	% - 29.58%]	28.8%	[28.59% - 29.01%]
% 65+ with anxiety disorders	В	16.1%	[13.179	6 - 19.03%]	21.9%	[21.74% - 22.15%]
% 65+ with bipolar disorders	В	2.2%	[1.549	% - 2.79%]	3.1%	[2.97% - 3.14%]
% 65+ with post-traumatic stress disorder	В	0.8%	[0.389	% - 1.16%]	1.4%	[1.35% - 1.48%]
% 65+ with schizophrenia & other psychotic disorders	В	3.8%	[3.069	% - 4.61%]	4.9%	[4.81% - 5.02%]
% 65+ with personality disorders		0.7%	[0.349	6 - 1.06%]	1.1%	[1.00% - 1.11%]
% 65+ with substance use disorders (drug use +/or alcohol abuse)	В	3.8%	[2.369	6 - 5.30%]	5.5%	[5.41% - 5.64%]
% 65+ with tobacco use disorders		8.3%	[6.029	6 - 10.49%]	10.4%	[10.23% - 10.54%]
CHRONIC DISEASE						
% 65+ with Alzheimer's disease or related dementias	W	20.1%	[17.039	% - 23.14%]	12.0%	[11.89% - 12.19%]
% 65+ with diabetes	В	22.7%	[19.239	- 6 - 26.08%]	28.2%	[27.97% - 28.41%]
% 65+ with stroke		10.7%	[8.389	6 - 13.03%]	10.8%	[10.64% - 10.92%]
% 65+ with chronic obstructive pulmonary disease	В	15.1%	[12.449	6 - 17.79%]	20.5%	[20.36% - 20.73%]
% 65+ with asthma	В	9.5%	[7.299	% - 11.78%]	13.0%	[12.89% - 13.21%]
% 65+ with hypertension	В	61.2%	[57.399	% - 65.09%]	70.2%	[70.02% - 70.46%]
% 65+ ever had a heart attack		3.2%	[1.909	% - 4.50%]	4.5%	[4.37% - 4.56%]
% 65+ with ischemic heart disease	В	29.0%	[25.289	6 - 32.65%]	34.3%	[34.12% - 34.58%]
% 65+ with congestive heart failure		17.2%	[14.299	% - 20.13%]	17.7%	[17.57% - 17.93%]
% 65+ with atrial fibrillation		13.2%	[10.629	% - 15.70%]	14.4%	[14.28% - 14.60%]
% 65+ with peripheral vascular disease	В	10.8%	[8.469	% - 13.20%]	14.7%	[14.51% - 14.84%]
% 65+ with osteoarthritis/rheumatoid arthritis		48.4%	[44.199	6 - 52.53%]	49.1%	[48.82% - 49.32%]
% 65+ with osteoporosis	В	13.9%	[11.319	6 - 16.53%]	17.4%	[17.22% - 17.57%]
% 65+ with leukemias and lymphomas		1.6%	[1.079	% - 2.16%]	2.0%	[1.91% - 2.05%]
% 65+ with lung cancer		1.3%	[0.809	% - 1.71%]	1.6%	[1.54% - 1.66%]
% 65+ with colon cancer	В	1.8%	[1.279	% - 2.31%]	2.4%	[2.37% - 2.51%]
% 65+ women with breast cancer	W	13.7%	•	6 - 17.31%]	9.8%	[9.59% - 9.96%]
% 65+ women with endometrial cancer		1.9%	[1.129		1.7%	[1.60% - 1.76%]
% 65+ men with prostate cancer		8.3%	•	6 - 11.33%]	11.5%	[11.24% - 11.68%]
% 65+ with benign prostatic hyperplasia		39.3%	-	6 - 44.78%]	36.8%	[36.47% - 37.14%]
% 65+ with HIV/AIDS		0.04%	[0.029	-	0.05%	[0.04% - 0.06%]
% 65+ with hypothyroidism	В	17.1%	-	% - 19.96% <u>]</u>	20.8%	[20.62% - 20.99%]
% 65+ with anemia		36.5%	-	6 - 40.16%]	37.3%	[37.12% - 37.57%]
% 65+ with chronic kidney disease		19.2%	-	6 - 22.31%]	22.3%	[22.14% - 22.54%]
% 65+ with liver diseases		5.3%	[3.549	-	6.9%	[6.75% - 6.99%]
% 65+ with fibromyalgia, chronic pain and fatigue	В	15.3%	-	6 - 18.24%]	18.6%	[18.42% - 18.80%]
% 65+ with migraine and other chronic headache		3.3%	[2.529	-	4.0%	[3.95% - 4.15%]
% 65+ with epilepsy		1.5%	[1.049		2.1%	[2.00% - 2.14%]
% 65+ with traumatic brain injury		0.8%	[0.449	% - 1.15%]	1.1%	[1.06% - 1.16%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders				N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma	В	17.8%	[14.85% - 20.67%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	W	65.8%	[61.99% - 69.60%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer	В	4.9%]	3.22% - 6.50%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	В	47.3%	[43.44% - 51.18%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions	В	13.5%	[10.62% - 16.32%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment		13.4%	[10.82% - 15.95%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	[0.39% - 1.04%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.7%	[2.88% - 4.43%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees	*	5.2%	[3.61% - 6.69%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	4.9%	[3.40% - 6.40%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		94.6%	[91.78% - 97.51%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		3.3%	[1.16% - 5.46%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION						
Physician visits per year	*	5.5	[5.1 - 6.0]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year	*	463	[379 - 547]	572	[566 - 578]
Part D monthly prescription fills per person per year	*	43.4	[40.1 - 46.7]	49.1	[48.9 - 49.3]
Home health visits per year		2.2	[1.5 - 2.9]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year	*	1.3	[0.9 - 1.6]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year	*	159	[121 - 197]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		13.2%	[7.58% - 18.83%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year	*	46	[28 - 64]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	1.9%	[0.94% - 2.83%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month		77.1%	[72.03% - 82.23%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt		75.0%	[69.58% - 80.44%]	77.1%	[76.02% - 78.18%]
· ·			-	-		

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.
² All community and state margins of error are 95% confidence intervals.
³ The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu