Warner (Merrimack)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS						
Age-sex adjusted 1-year mortality rate		3.3%	[2.00% - 4.57%]	4.1%	[4.02% - 4.19%
WELLNESS & PREVENTION						
% 60+ with any physical activity within last month		74.2%	[67.37% - 80.97%]	74.5%	[73.36% - 75.59%
% 60+ met CDC guidelines for muscle-strengthening activity		23.1%	[17.41% - 28.70%]	26.0%	[24.93% - 27.14%
% 60+ met CDC guidelines for aerobic physical activity		61.5%	[54.68% - 68.29%]	58.6%	[57.33% - 59.82%
% 60+ met CDC guidelines for both types of physical activities		17.7%	[12.66% - 22.78%]	19.9%	[18.92% - 20.93%
% 60+ getting recommended hours of sleep		66.8%	[59.69% - 73.83%]	66.4%	[65.16% - 67.60%
% 60+ injured in a fall within last 12 months		7.1%	[3.38% - 10.80%]	10.4%	[9.65% - 11.16%
% 65+ had hip fracture		2.6%	[2.01% - 3.25%]	3.3%	[3.19% - 3.34%
% 60+ with self-reported fair or poor health status		15.4%	[9.64% - 21.19%]	16.5%	[15.54% - 17.40%
% 60+ with 15+ physically unhealthy days last month		10.7%	[6.12% - 15.36%]	12.5%	[11.69% - 13.37%
% 60+ with physical exam/check-up in past year		84.8%	[79.83% - 89.87%]	86.5%	[85.66% - 87.39%
% 60+ met CDC preventive health screening goals		35.3%	[26.27% - 44.24%]	40.3%	[38.77% - 41.81%
% 60+ flu shot past year		58.9%	[51.65% - 66.10%]	59.3%	[58.06% - 60.63%
% 65+ with pneumonia vaccine		73.7%	[66.00% - 81.35%]	77.8%	[76.56% - 79.12%
% 60+ with cholesterol screening		92.1%	[88.33% - 95.93%]	95.3%	[94.78% - 95.82%
% 60+ women with a mammogram within last 2 years		84.1%	[76.93% - 91.33%]	79.3%	[77.96% - 80.69%
% 60+ with colorectal cancer screening		78.7%	[72.47% - 84.88%]	77.0%	[75.92% - 78.16%
% 60+ with HIV test		17.4%	[11.44% - 23.33%]	13.5%	[12.50% - 14.41%
% 60+ current smokers		8.3%	[3.58% - 12.94%]	8.1%	[7.38% - 8.79%
Oral Health						
% 60+ with loss of 6 or more teeth		24.0%	[17.45% - 30.62%]	29.0%	[27.79% - 30.12%
% 60+ with annual dental exam		77.3%	[71.00% - 83.64%]	75.7%	[74.65% - 76.84%
NUTRITION/DIET						
% 60+ with 5 or more servings of fruit or vegetables per day		22.7%	[17.23% - 28.22%]	20.2%	[19.23% - 21.21%
% 60+ self-reported obese		24.3%	[17.69% - 30.94%]	27.2%	[26.03% - 28.35%
% 65+ clinically diagnosed obese		16.9%	[13.92% - 19.78%]	16.7%	[16.56% - 16.94%
% 65+ with high cholesterol		70.1%	[66.61% - 73.57%]	72.2%	[72.01% - 72.44%
% 60+ excessive drinking		6.5%	[2.94% - 10.08%]	9.2%	[8.45% - 9.94%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH						
% 60+ with 15+ days poor mental health last month		7.8%	[3.52% - 12.02%]	6.9%	[6.29% - 7.61%]
% 65+ with depression		26.6%	[23.35% - 29.85%]	28.8%	[28.59% - 29.01%]
% 65+ with anxiety disorders	В	18.6%	[15.59% - 21.67%]	21.9%	[21.74% - 22.15%]
% 65+ with bipolar disorders		2.4%	[1.73% - 3.03%]	3.1%	[2.97% - 3.14%]
% 65+ with post-traumatic stress disorder		1.4%	[0.83% - 1.88%]	1.4%	[1.35% - 1.48%]
% 65+ with schizophrenia & other psychotic disorders	В	2.9%	[2.24% - 3.61%]	4.9%	[4.81% - 5.02%]
% 65+ with personality disorders		1.0%	[0.53% - 1.42%]	1.1%	[1.00% - 1.11%]
% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.7%	[3.84% - 7.51%]	5.5%	[5.41% - 5.64%]
% 65+ with tobacco use disorders		10.0%	[7.64% - 12.29%]	10.4%	[10.23% - 10.54%]
CHRONIC DISEASE						
% 65+ with Alzheimer's disease or related dementias	В	9.7%	ſ	7.55% - 11.79%]	12.0%	[11.89% - 12.19%]
% 65+ with diabetes	B	22.6%		19.33% - 25.79%]	28.2%	[27.97% - 28.41%]
% 65+ with stroke	В	8.5%	[6.52% - 10.48%]	10.8%	[10.64% - 10.92%]
% 65+ with chronic obstructive pulmonary disease		18.0%	[15.23% - 20.85%]	20.5%	[20.36% - 20.73%]
% 65+ with asthma		10.8%	[8.56% - 13.13%]	13.0%	[12.89% - 13.21%
% 65+ with hypertension	В	65.4%	[61.83% - 69.04%]	70.2%	[70.02% - 70.46%]
% 65+ ever had a heart attack		4.0%	[2.65% - 5.34%]	4.5%	[4.37% - 4.56%]
% 65+ with ischemic heart disease	В	28.2%	[24.83% - 31.64%]	34.3%	[34.12% - 34.58%]
% 65+ with congestive heart failure	В	11.1%	[8.86% - 13.38%]	17.7%	[17.57% - 17.93%
% 65+ with atrial fibrillation		12.9%	[10.43% - 15.29%]	14.4%	[14.28% - 14.60%
% 65+ with peripheral vascular disease	В	9.8%	[7.65% - 12.02%]	14.7%	[14.51% - 14.84%
% 65+ with osteoarthritis/rheumatoid arthritis		46.4%	[42.49% - 50.36%]	49.1%	[48.82% - 49.32%
% 65+ with osteoporosis		15.5%	[12.94% - 18.12%]	17.4%	[17.22% - 17.57%
% 65+ with leukemias and lymphomas		1.9%	[1.34% - 2.42%]	2.0%	[1.91% - 2.05%
% 65+ with lung cancer	В	1.1%	[0.70% - 1.52%]	1.6%	[1.54% - 1.66%
% 65+ with colon cancer		2.0%	[1.43% - 2.53%]	2.4%	[2.37% - 2.51%
% 65+ women with breast cancer		9.0%	[6.09% - 11.97%]	9.8%	[9.59% - 9.96%
% 65+ women with endometrial cancer		1.2%	[0.57% - 1.74%]	1.7%	[1.60% - 1.76%]
% 65+ men with prostate cancer		11.7%	[8.44% - 14.88%]	11.5%	[11.24% - 11.68%
% 65+ with benign prostatic hyperplasia		36.9%	[31.77% - 41.93%]	36.8%	[36.47% - 37.14%
% 65+ with HIV/AIDS		0.07%	[0.04% - 0.10%]	0.05%	[0.04% - 0.06%
% 65+ with hypothyroidism		19.3%	-	16.41% - 22.15%]	20.8%	[20.62% - 20.99%]
% 65+ with anemia	В	29.8%	-	26.44% - 33.10%]	37.3%	[37.12% - 37.57%]
% 65+ with chronic kidney disease	B	17.8%]	14.94% - 20.71%]	22.3%	[22.14% - 22.54%
% 65+ with liver diseases	В	4.3%	[2.76% - 5.84%]	6.9%	[6.75% - 6.99%
% 65+ with fibromyalgia, chronic pain and fatigue		16.7%	-	13.83% - 19.48%]	18.6%	[18.42% - 18.80%
% 65+ with migraine and other chronic headache		3.6%	[2.77% - 4.39%]	4.0%	[3.95% - 4.15%
% 65+ with epilepsy		1.6%	[1.12% - 2.15%]	2.1%	[2.00% - 2.14%
% 65+ with traumatic brain injury		0.8%	l	0.41% - 1.09%]	1.1%	[1.06% - 1.16%

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders				N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		24.3%	[21.21% - 27.48%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		59.8%	[56.05% - 63.49%]	61.2%	[61.00% - 61.47%
% 65+ with pressure ulcer or chronic ulcer	В	4.2%	[2.72% - 5.74%]	7.1%	[6.99% - 7.24%
% 65+ with 4+ (out of 15) chronic conditions	В	48.2%]	44.52% - 51.97%]	54.4%	[54.19% - 54.66%
% 65+ with 0 chronic conditions		9.9%	[7.55% - 12.26%]	10.3%	[10.12% - 10.42%
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment	В	11.1%	[8.73% - 13.40%]	14.4%	[14.23% - 14.56%
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	[0.39% - 1.02%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		2.7%	[2.04% - 3.36%]	3.2%	[3.14% - 3.31%
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees		7.8%	[6.04% - 9.48%]	7.9%	[7.8% - 8.02%
% dually eligible for Medicare and Medicaid		6.1%	[4.60% - 7.69%]	7.5%	[7.40% - 7.62%
% 60+ with a regular doctor		95.7%	[92.56% - 98.74%]	95.8%	[95.34% - 96.33%
% 60+ who did not see doctor when needed due to cost		4.4%	[1.17% - 7.72%]	5.4%	[4.75% - 5.97%
SERVICE UTILIZATION						
Physician visits per year	*	5.4	[4.9 - 5.8]	6.3	[6.2 - 6.3
Emergency room visits/1000 persons 65+ years per year		522	[366 - 678]	572	[566 - 578
Part D monthly prescription fills per person per year	*	43.9	[40.7 - 47.1]	49.1	[48.9 - 49.3
Home health visits per year		2.9	[1.9 - 3.9]	2.5	[2.4 - 2.5
Durable medical equipment claims per year		1.7	[1.3 - 2.1]	2.0	[2.0 - 2.0
Inpatient hospital stays/1000 persons 65+ years per year		192	[140 - 243]	237	[234 - 241
Medicare inpatient hospital readmissions (as % of admissions)		19.1%	[10.72% - 27.57%]	15.8%	[15.41% - 16.22%
# skilled nursing facility stays/1000 persons 65+ years per year	*	45	[25 - 64]	76	[74 - 77
% 65+ getting Medicaid long term services and supports	*	2.3%	[1.31% - 3.22%]	3.7%	[3.59% - 3.74%
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month		83.0%	[77.62% - 88.43%]	77.6%	[76.53% - 78.64%
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt		75.0%	[68.20% - 81.83%]	77.1%	[76.02% - 78.18%
				-		

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.
² All community and state margins of error are 95% confidence intervals.
³ The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu