Bethlehem (Grafton)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²	
POPULATION CHARACTERISTICS							
Age-sex adjusted 1-year mortality rate		5.4%	[3.06% - 7.82%]	4.1%	[4.02% - 4.19%	
WELLNESS & PREVENTION							
% 60+ with any physical activity within last month		73.6%	[65.84% - 81.31%]	74.5%	[73.36% - 75.59%	
% 60+ met CDC guidelines for muscle-strengthening activity		23.8%	[16.42% - 31.24%]	26.0%	[24.93% - 27.14%	
% 60+ met CDC guidelines for aerobic physical activity		55.3%	[46.49% - 64.01%]	58.6%	[57.33% - 59.82%	
% 60+ met CDC guidelines for both types of physical activities		17.3%	[10.93% - 23.67%]	19.9%	[18.92% - 20.93%	
% 60+ getting recommended hours of sleep		64.8%	[56.58% - 73.06%]	66.4%	[65.16% - 67.60%	
% 60+ injured in a fall within last 12 months		9.6%	[4.98% - 14.31%]	10.4%	[9.65% - 11.16%	
% 65+ had hip fracture		3.4%	[2.66% - 4.21%]	3.3%	[3.19% - 3.34%	
% 60+ with self-reported fair or poor health status		18.0%	[11.61% - 24.32%]	16.5%	[15.54% - 17.40%	
% 60+ with 15+ physically unhealthy days last month		16.6%	[10.00% - 23.16%]	12.5%	[11.69% - 13.37%	
% 60+ with physical exam/check-up in past year		83.2%	[76.56% - 89.85%]	86.5%	[85.66% - 87.39%	
% 60+ met CDC preventive health screening goals		38.2%	[28.68% - 47.64%]	40.3%	[38.77% - 41.81%	
% 60+ flu shot past year		52.1%	[43.69% - 60.52%]	59.3%	[58.06% - 60.63%	
% 65+ with pneumonia vaccine		74.9%	[67.21% - 82.59%]	77.8%	[76.56% - 79.12%	
% 60+ with cholesterol screening		92.9%	[88.51% - 97.35%]	95.3%	[94.78% - 95.82%	
% 60+ women with a mammogram within last 2 years		77.0%	[67.08% - 86.85%]	79.3%	[77.96% - 80.69%	
% 60+ with colorectal cancer screening		80.9%	[75.27% - 86.56%]	77.0%	[75.92% - 78.16%	
% 60+ with HIV test		16.2%	[9.42% - 23.00%]	13.5%	[12.50% - 14.41%	
% 60+ current smokers		10.6%	[5.99% - 15.25%]	8.1%	[7.38% - 8.79%	
Oral Health							
% 60+ with loss of 6 or more teeth		33.6%	[26.22% - 40.88%]	29.0%	[27.79% - 30.12%	
% 60+ with annual dental exam		69.2%	[62.16% - 76.20%]	75.7%	[74.65% - 76.84%	
NUTRITION/DIET							
% 60+ with 5 or more servings of fruit or vegetables per day		15.2%	[9.62% - 20.72%]	20.2%	[19.23% - 21.21%	
% 60+ self-reported obese		27.6%	[19.69% - 35.47%]	27.2%	[26.03% - 28.35%	
% 65+ clinically diagnosed obese	В	12.2%	[8.61% - 15.82%]	16.7%	[16.56% - 16.94%	
% 65+ with high cholesterol	В	59.9%	[54.46% - 65.36%]	72.2%	[72.01% - 72.44%	
% 60+ excessive drinking		7.6%	[3.22% - 12.07%]	9.2%	[8.45% - 9.94%	

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH						
% 60+ with 15+ days poor mental health last month		8.0%	[3.0	9% - 12.88%]	6.9%	[6.29% - 7.61%]
% 65+ with depression	В	22.3%	[17.7	8% - 26.83%]	28.8%	[28.59% - 29.01%]
% 65+ with anxiety disorders	В	16.1%	[11.9	4% - 20.28%]	21.9%	[21.74% - 22.15%]
% 65+ with bipolar disorders	В	1.6%	[1.0	3% - 2.26%]	3.1%	[2.97% - 3.14%]
% 65+ with post-traumatic stress disorder		1.8%	[1.1	6% - 2.54%]	1.4%	[1.35% - 1.48%]
% 65+ with schizophrenia & other psychotic disorders		4.4%	[3.4	5% - 5.38%]	4.9%	[4.81% - 5.02%]
% 65+ with personality disorders		0.9%	[0.4	4% - 1.34%]	1.1%	[1.00% - 1.11%]
% 65+ with substance use disorders (drug use +/or alcohol abuse)		6.6%	[3.6	9% - 9.60%]	5.5%	[5.41% - 5.64%]
% 65+ with tobacco use disorders		9.1%	[5.9	1% - 12.33%]	10.4%	[10.23% - 10.54%]
CHRONIC DISEASE						
% 65+ with Alzheimer's disease or related dementias		9.8%	[6.6	5% - 12.95%]	12.0%	[11.89% - 12.19%]
% 65+ with diabetes	В	19.4%	[15.0	5% - 23.83%]	28.2%	[27.97% - 28.41%]
% 65+ with stroke	В	7.2%	[4.5	9% - 9.85%]	10.8%	[10.64% - 10.92%]
% 65+ with chronic obstructive pulmonary disease	В	15.9%	[12.0	0% - 19.82%]	20.5%	[20.36% - 20.73%]
% 65+ with asthma		12.3%	[8.8	0% - 15.80%]	13.0%	[12.89% - 13.21%]
% 65+ with hypertension	В	62.2%	[56.7	9% - 67.62%]	70.2%	[70.02% - 70.46%]
% 65+ ever had a heart attack		4.4%	[2.2	3% - 6.57%]	4.5%	[4.37% - 4.56%]
% 65+ with ischemic heart disease		29.9%	[24.7	8% - 34.92%]	34.3%	[34.12% - 34.58%]
% 65+ with congestive heart failure		18.2%	[13.9	4% - 22.37%]	17.7%	[17.57% - 17.93%]
% 65+ with atrial fibrillation	В	10.7%	[7.4	6% - 13.87%]	14.4%	[14.28% - 14.60%]
% 65+ with peripheral vascular disease	В	10.8%	[7.4	5% - 14.12%]	14.7%	[14.51% - 14.84%]
% 65+ with osteoarthritis/rheumatoid arthritis	В	38.4%	[33.0	0% - 43.79%]	49.1%	[48.82% - 49.32%]
% 65+ with osteoporosis		14.8%	[11.1	9% - 18.50%]	17.4%	[17.22% - 17.57%]
% 65+ with leukemias and lymphomas		1.7%	[1.1	1% - 2.31%]	2.0%	[1.91% - 2.05%]
% 65+ with lung cancer		1.4%	[0.8	7% - 1.94%]	1.6%	[1.54% - 1.66%]
% 65+ with colon cancer		2.6%	[1.8	9% - 3.27%]	2.4%	[2.37% - 2.51%]
% 65+ women with breast cancer		9.5%	[5.2	8% - 13.74%]	9.8%	[9.59% - 9.96%]
% 65+ women with endometrial cancer		2.0%	[1.1	1% - 2.83%]	1.7%	[1.60% - 1.76%]
% 65+ men with prostate cancer		10.8%	[6.1	8% - 15.41%]	11.5%	[11.24% - 11.68%]
% 65+ with benign prostatic hyperplasia		35.8%	[28.3	1% - 43.23%]	36.8%	[36.47% - 37.14%]
% 65+ with HIV/AIDS		0.05%	[0.0	2% - 0.08%]	0.05%	[0.04% - 0.06%]
% 65+ with hypothyroidism	В	13.2%	[9.5	9% - 16.77%]	20.8%	[20.62% - 20.99%]
% 65+ with anemia		32.3%	[27.2	7% - 37.27%]	37.3%	[37.12% - 37.57%]
% 65+ with chronic kidney disease	В	13.5%	[9.7	1% - 17.21%]	22.3%	[22.14% - 22.54%]
% 65+ with liver diseases		5.0%	-	0% - 7.48%]	6.9%	[6.75% - 6.99%]
% 65+ with fibromyalgia, chronic pain and fatigue		15.4%	[11.4	2% - 19.41%]	18.6%	[18.42% - 18.80%]
% 65+ with migraine and other chronic headache		3.8%	-	9% - 4.66%]	4.0%	[3.95% - 4.15%]
% 65+ with epilepsy		1.8%	•	8% - 2.38%]	2.1%	[2.00% - 2.14%]
% 65+ with traumatic brain injury		0.9%	[0.4	4% - 1.26%]	1.1%	[1.06% - 1.16%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders				N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		19.0%	[14.89% - 23.11%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract	В	55.0%	[49.47% - 60.47%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		6.6%	[3.77% - 9.35%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	В	46.2%	[40.80% - 51.63%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions	В	17.9%	[13.38% - 22.40%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment		12.7%	[9.15% - 16.16%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.7%	[0.32% - 1.01%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		2.7%	[1.96% - 3.43%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees		6.9%	[4.55% - 9.23%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid		6.7%	[4.36% - 8.97%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		96.4%	[94.19% - 98.69%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		9.7%	[4.77% - 14.66%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION						
Physician visits per year	*	3.9]	3.3 - 4.5]	6.3	[6.2 - 6.3
Emergency room visits/1000 persons 65+ years per year		549	[415 - 684]	572	[566 - 578
Part D monthly prescription fills per person per year	*	42.6	[37.9 - 47.3]	49.1	[48.9 - 49.3
Home health visits per year		2.0	[0.9 - 3.1]	2.5	[2.4 - 2.5
Durable medical equipment claims per year	*	1.3	[0.9 - 1.7]	2.0	[2.0 - 2.0
Inpatient hospital stays/1000 persons 65+ years per year	*	167	[110 - 225]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)	*	7.5%	[0.94% - 14.15%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year		69	[30 - 108]	76	[74 - 77
% 65+ getting Medicaid long term services and supports	*	1.8%	[0.56% - 3.00%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month		72.9%]	65.62% - 80.17%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt		73.0%	[66.15% - 79.79%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.
² All community and state margins of error are 95% confidence intervals.
³ The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu