## Nashua: Zip 03060 (Hillsborough)



|  |   |                       |   |                                 | FO                | U | NDAI        | ION        |
|--|---|-----------------------|---|---------------------------------|-------------------|---|-------------|------------|
| HEALTHY AGING INDICATORS                                       | BETTER /<br>WORSE<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE |   | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE |   | MARG<br>ERR |            |
| POPULATION CHARACTERISTICS                                     |   |                       |   |                                 |                   | П |             |            |
| Age-sex adjusted 1-year mortality rate                         |   | 3.9%                  | [ | 3.28% - 4.58%                   | ] 4.1%            | [ | 4.02%       | - 4.19% ]  |
| WELLNESS & PREVENTION  |   |                       |   |                                 |                   |   |             |            |
| % 60+ with any physical activity within last month             |   | 72.6%                 | [ | 67.41% - 77.77%                 | ] 74.5%           | [ | 73.36%      | - 75.59% ] |
| % 60+ met CDC guidelines for muscle-strengthening activity     |   | 23.1%                 | [ | 18.20% - 28.00%                 | ] 26.0%           | [ | 24.93%      | - 27.14% ] |
| % 60+ met CDC guidelines for aerobic physical activity         |   | 55.9%                 | [ | 50.23% - 61.50%                 | ] 58.6%           | [ | 57.33%      | - 59.82% ] |
| % 60+ met CDC guidelines for both types of physical activities |   | 17.4%                 | [ | 12.94% - 21.88%                 | ] 19.9%           | [ | 18.92%      | - 20.93% ] |
| % 60+ getting recommended hours of sleep                       |   | 62.1%                 | [ | 56.57% - 67.63%                 | ] 66.4%           | [ | 65.16%      | - 67.60% ] |
| % 60+ injured in a fall within last 12 months                  |   | 12.5%                 | [ | 8.53% - 16.47%                  | ] 10.4%           | [ | 9.65%       | - 11.16% ] |
| % 65+ had hip fracture   |   | 3.6%                  | [ | 2.95% - 4.25%                   | ] 3.3%            | [ | 3.19%       | - 3.34% ]  |
| % 60+ with self-reported fair or poor health status            |   | 20.9%                 | [ | 15.94% - 25.88%                 | ] 16.5%           | [ | 15.54%      | - 17.40% ] |
| % 60+ with 15+ physically unhealthy days last month            | W   | 19.7%                 | [ | 14.28% - 25.03%                 | ] 12.5%           | [ | 11.69%      | - 13.37% ] |
| % 60+ with physical exam/check-up in past year                 |   | 89.0%                 | [ | 85.69% - 92.34%                 | ] 86.5%           | [ | 85.66%      | - 87.39% ] |
| % 60+ met CDC preventive health screening goals                |   | 44.4%                 | [ | 37.25% - 51.50%                 | ] 40.3%           | [ | 38.77%      | - 41.81% ] |
| % 60+ flu shot past year                                       |   | 59.8%                 | [ | 53.56% - 66.07%                 | ] 59.3%           | [ | 58.06%      | - 60.63% ] |
| % 65+ with pneumonia vaccine                                   |   | 81.5%                 | [ | 74.93% - 87.99%                 | ] 77.8%           | [ | 76.56%      | - 79.12% ] |
| % 60+ with cholesterol screening                               |   | 93.8%                 | [ | 90.37% - 97.18%                 | ] 95.3%           | [ | 94.78%      | - 95.82% ] |
| % 60+ women with a mammogram within last 2 years               |   | 77.9%                 | [ | 69.66% - 86.13%                 | ] 79.3%           | [ | 77.96%      | - 80.69% ] |
| % 60+ with colorectal cancer screening                         |   | 75.8%                 | [ | 69.97% - 81.55%                 | ] 77.0%           | [ | 75.92%      | - 78.16% ] |
| % 60+ with HIV test  |   | 13.6%                 | [ | 8.63% - 18.48%                  | ] 13.5%           | [ | 12.50%      | - 14.41% ] |
| % 60+ current smokers  |   | 11.6%                 | [ | 6.80% - 16.49%                  | ] 8.1%            | [ | 7.38%       | - 8.79% ]  |
| Oral Health  |   |                       |   |                                 |                   |   |             |            |
| % 60+ with loss of 6 or more teeth                             |   | 28.4%                 | [ | 23.31% - 33.58%                 | ] 29.0%           | [ | 27.79%      | - 30.12% ] |
| % 60+ with annual dental exam                                  |   | 70.6%                 | [ | 64.95% - 76.23%                 | ] 75.7%           | [ | 74.65%      | - 76.84% ] |
| NUTRITION/DIET   |   |                       |   |                                 |                   |   |             |            |
| % 60+ with 5 or more servings of fruit or vegetables per day   |   | 18.9%                 | [ | 14.58% - 23.14%                 | ] 20.2%           | [ | 19.23%      | - 21.21% ] |
| % 60+ self-reported obese                                      |   | 26.0%                 | [ | 20.93% - 31.03%                 | ] 27.2%           | [ | 26.03%      | - 28.35% ] |
| % 65+ clinically diagnosed obese                               | W   | 19.4%                 | [ | 17.78% - 20.94%                 | ] 16.7%           | [ | 16.56%      | - 16.94% ] |
| % 65+ with high cholesterol                                    | W   | 77.7%                 | [ | 76.11% - 79.36%                 | ] 72.2%           | [ | 72.01%      | - 72.44% ] |
| % 60+ excessive drinking                                       |   | 7.4%                  | [ | 4.55% - 10.21%                  | ] 9.2%            | [ | 8.45%       | 9.94% ]    |
|  |   |                       |   |                                 |                   |   |             |            |

Nashua: Zip 03060 (Hillsborough)

| HEALTHY AGING INDICATORS   | BETTER /<br>WORSE<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE |   | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---|---------------------------------|-------------------|---------------------------------|
| BEHAVIORAL HEALTH  |   |                       | Т |                                 |                   |                                 |
| % 60+ with 15+ days poor mental health last month                |   | 9.2%                  | [ | 5.59% - 12.90% ]                | 6.9%              | [ 6.29% - 7.61% ]               |
| % 65+ with depression  | W   | 34.6%                 | [ | 32.78% - 36.35% ]               | 28.8%             | [ 28.59% - 29.01% ]             |
| % 65+ with anxiety disorders                                     | W   | 26.6%                 | [ | 24.86% - 28.33% ]               | 21.9%             | [ 21.74% - 22.15% ]             |
| % 65+ with bipolar disorders                                     | W   | 4.8%                  | [ | 3.92% - 5.70% ]                 | 3.1%              | [ 2.97% - 3.14% ]               |
| % 65+ with post-traumatic stress disorder                        |   | 1.5%                  | [ | 0.94% - 2.01% ]                 | 1.4%              | [ 1.35% - 1.48% ]               |
| % 65+ with schizophrenia & other psychotic disorders             | W   | 6.1%                  | [ | 5.20% - 7.00% ]                 | 4.9%              | [ 4.81% - 5.02% ]               |
| % 65+ with personality disorders                                 |   | 1.5%                  | [ | 1.01% - 2.02% ]                 | 1.1%              | [ 1.00% - 1.11% ]               |
| % 65+ with substance use disorders (drug use +/or alcohol abuse) | W   | 7.0%                  | [ | 5.97% - 8.10% ]                 | 5.5%              | [ 5.41% - 5.64% ]               |
| % 65+ with tobacco use disorders                                 | W   | 15.4%                 | [ | 13.91% - 16.90% ]               | 10.4%             | [ 10.23% - 10.54% ]             |
| CHRONIC DISEASE  |   |                       |   |                                 |                   |                                 |
| % 65+ with Alzheimer's disease or related dementias              |   | 12.8%                 | [ | 11.57% - 14.01% ]               | 12.0%             | [ 11.89% - 12.19% ]             |
| % 65+ with diabetes  | W   | 34.9%                 | [ | 33.01% - 36.77% ]               | 28.2%             | [ 27.97% - 28.41% ]             |
| % 65+ with stroke  | W   | 13.5%                 | [ | 12.29% - 14.76% ]               | 10.8%             | [ 10.64% - 10.92% ]             |
| % 65+ with chronic obstructive pulmonary disease                 | W   | 26.1%                 | [ | 24.46% - 27.72% ]               | 20.5%             | [ 20.36% - 20.73% ]             |
| % 65+ with asthma  | W   | 14.7%                 | [ | 13.42% - 16.06% ]               | 13.0%             | [ 12.89% - 13.21% ]             |
| % 65+ with hypertension  | W   | 77.4%                 | [ | 75.79% - 79.08% ]               | 70.2%             | [ 70.02% - 70.46% ]             |
| % 65+ ever had a heart attack                                    | W   | 6.0%                  | [ | 5.12% - 6.84% ]                 | 4.5%              | [ 4.37% - 4.56% ]               |
| % 65+ with ischemic heart disease                                | W   | 38.9%                 | [ | 37.03% - 40.83% ]               | 34.3%             | [ 34.12% - 34.58% ]             |
| % 65+ with congestive heart failure                              | W   | 21.7%                 | [ | 20.12% - 23.25% ]               | 17.7%             | [ 17.57% - 17.93% ]             |
| % 65+ with atrial fibrillation                                   | W   | 16.7%                 | [ | 15.38% - 18.10% ]               | 14.4%             | [ 14.28% - 14.60% ]             |
| % 65+ with peripheral vascular disease                           | W   | 16.8%                 | [ | 15.39% - 18.25% ]               | 14.7%             | [ 14.51% - 14.84% ]             |
| % 65+ with osteoarthritis/rheumatoid arthritis                   | W   | 51.7%                 | [ | 49.70% - 53.67% ]               | 49.1%             | [ 48.82% - 49.32% ]             |
| % 65+ with osteoporosis  | W   | 22.5%                 | [ | 21.02% - 24.03% ]               | 17.4%             | [ 17.22% - 17.57% ]             |
| % 65+ with leukemias and lymphomas                               |   | 2.3%                  | [ | 1.75% - 2.88% ]                 | 2.0%              | [ 1.91% - 2.05%]                |
| % 65+ with lung cancer   |   | 1.7%                  | [ | 1.24% - 2.20% ]                 | 1.6%              | [ 1.54% - 1.66% ]               |
| % 65+ with colon cancer  |   | 2.6%                  | [ | 2.02% - 3.15% ]                 | 2.4%              | [ 2.37% - 2.51%]                |
| % 65+ women with breast cancer                                   |   | 9.4%                  | [ | 8.07% - 10.80% ]                | 9.8%              | [ 9.59% - 9.96% ]               |
| % 65+ women with endometrial cancer                              |   | 1.8%                  | [ | 1.21% - 2.47% ]                 | 1.7%              | [ 1.60% - 1.76% ]               |
| % 65+ men with prostate cancer                                   |   | 11.5%                 | [ | 9.70% - 13.34% ]                | 11.5%             | [ 11.24% - 11.68% ]             |
| % 65+ with benign prostatic hyperplasia                          |   | 39.2%                 | [ | 36.28% - 42.09% ]               | 36.8%             | [ 36.47% - 37.14% ]             |
| % 65+ with HIV/AIDS  |   | 0.04%                 | [ | 0.02% - 0.07% ]                 | 0.05%             | [ 0.04% - 0.06% ]               |
| % 65+ with hypothyroidism  | W   | 23.8%                 | [ | 22.26% - 25.39% ]               | 20.8%             | [ 20.62% - 20.99% ]             |
| % 65+ with anemia  | W   | 47.6%                 | [ | 45.71% - 49.47% ]               | 37.3%             | [ 37.12% - 37.57% ]             |
| % 65+ with chronic kidney disease                                | W   | 31.3%                 | [ | 29.53% - 33.13% ]               | 22.3%             | [ 22.14% - 22.54% ]             |
| % 65+ with liver diseases  | W   | 8.4%                  | [ | 7.29% - 9.45% ]                 | 6.9%              | [ 6.75% - 6.99% ]               |
| % 65+ with fibromyalgia, chronic pain and fatigue                | W   | 21.8%                 | [ | 20.19% - 23.43% ]               | 18.6%             | [ 18.42% - 18.80% ]             |
| % 65+ with migraine and other chronic headache                   |   | 4.7%                  | [ | 3.82% - 5.53% ]                 | 4.0%              | [ 3.95% - 4.15% ]               |
| % 65+ with epilepsy  | W   | 2.9%                  | [ | 2.26% - 3.59% ]                 | 2.1%              | [ 2.00% - 2.14% ]               |
| % 65+ with traumatic brain injury                                |   | 1.3%                  | [ | 0.85% - 1.68% ]                 | 1.1%              | [ 1.06% - 1.16% ]               |

| HEALTHY AGING INDICATORS   | BETTER /<br>WORSE<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| % 65+ with autism spectrum disorders                             |   |                       | N/A                             | 0.03%             | [ 0.02% - 0.04%                 |
| % 65+ with glaucoma  |   | 21.3%                 | [ 19.80% - 22.82% ]             | 22.9%             | [ 22.74% - 23.13%               |
| % 65+ with cataract  | W   | 63.5%                 | [ 61.63% - 65.37% ]             | 61.2%             | [ 61.00% - 61.47%               |
| % 65+ with pressure ulcer or chronic ulcer                       |   | 7.8%                  | [ 6.76% - 8.80% ]               | 7.1%              | [ 6.99% - 7.24%                 |
| % 65+ with 4+ (out of 15) chronic conditions                     | W   | 64.1%                 | [ 62.29% - 66.00% ]             | 54.4%             | [ 54.19% - 54.66%               |
| % 65+ with 0 chronic conditions                                  | W   | 7.6%                  | [ 6.49% - 8.64% ]               | 10.3%             | [ 10.12% - 10.42%               |
| LIVING WITH DISABILITY   |   |                       |                                 |                   |                                 |
| % 65+ with clinical diagnosis of deafness or hearing impairment  |   | 13.3%                 | [ 12.06% - 14.60% ]             | 14.4%             | [ 14.23% - 14.56%               |
| % 65+ with clinical diagnosis of blindness or visual impairment  | W   | 1.4%                  | [ 0.92% - 1.79% ]               | 0.9%              | [ 0.81% - 0.90%                 |
| % 65+ with clinical diagnosis of mobility impairments            | W   | 4.6%                  | [ 3.81% - 5.44%]                | 3.2%              | [ 3.14% - 3.31%                 |
| ACCESS TO CARE   |   |                       |                                 |                   |                                 |
| Medicare (65+ population)  |   |                       |                                 |                   |                                 |
| % Medicare managed care enrollees                                | *   | 12.3%                 | [ 11.21% - 13.30% ]             | 7.9%              | [ 7.8% - 8.02%                  |
| % dually eligible for Medicare and Medicaid                      | *   | 12.7%                 | [ 11.67% - 13.79% ]             | 7.5%              | [ 7.40% - 7.62%                 |
| % 60+ with a regular doctor                                      |   | 95.9%                 | [ 93.45% - 98.42% ]             | 95.8%             | [ 95.34% - 96.33%               |
| % 60+ who did not see doctor when needed due to cost             |   | 7.8%                  | [ 4.07% - 11.53% ]              | 5.4%              | [ 4.75% - 5.97%                 |
| SERVICE UTILIZATION  |   |                       |                                 |                   |                                 |
| Physician visits per year  | *   | 7.5                   | [ 7.2 - 7.7 ]                   | 6.3               | [ 6.2 - 6.3                     |
| Emergency room visits/1000 persons 65+ years per year            | *   | 739                   | [ 675 - 803 ]                   | 572               | [ 566 - 578                     |
| Part D monthly prescription fills per person per year            | *   | 54.2                  | [ 52.6 - 55.9 ]                 | 49.1              | [ 48.9 - 49.3                   |
| Home health visits per year                                      | *   | 3.1                   | [ 2.6 - 3.6 ]                   | 2.5               | [ 2.4 - 2.5                     |
| Durable medical equipment claims per year                        | *   | 2.4                   | [ 2.2 - 2.6 ]                   | 2.0               | [ 2.0 - 2.0                     |
| Inpatient hospital stays/1000 persons 65+ years per year         | *   | 292                   | [ 261 - 322 ]                   | 237               | [ 234 - 241                     |
| Medicare inpatient hospital readmissions (as % of admissions)    |   | 15.0%                 | [ 11.85% - 18.09% ]             | 15.8%             | [ 15.41% - 16.22%               |
| # skilled nursing facility stays/1000 persons 65+ years per year |   | 87                    | [ 73 - 100 ]                    | 76                | [ 74 - 77                       |
| % 65+ getting Medicaid long term services and supports           |   | 4.0%                  | [ 3.35% - 4.59%]                | 3.7%              | [ 3.59% - 3.74%                 |
| COMMUNITY VARIABLES & CIVIC ENGAGEMENT                           |   |                       |                                 |                   |                                 |
| % 60+ who used Internet in last month                            |   | 71.0%                 | [ 65.00% - 76.97% ]             | 77.6%             | [ 76.53% - 78.64%               |
| SAFETY & TRANSPORTATION  |   |                       |                                 |                   |                                 |
| % 60+ who always drive wearing a seatbelt                        |   | 81.1%                 | [ 76.26% - 85.95% ]             | 77.1%             | [ 76.02% - 78.18%               |
| · •  |   |                       | · ·                             |                   | -                               |

Questions or suggestions? Beth.Dugan@umb.edu

Notes. 

Community-state differences where the margin of error do not overlap each other are noted B, W, or \*.

All community and state margins of error are 95% confidence intervals.

The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last montth, and did not see doctor when needed due to cost.