

Chester (Rockingham)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
POPULATION CHARACTERISTICS					
Age-sex adjusted 1-year mortality rate		3.5%	[1.73% - 5.36%]	4.1%	[4.02% - 4.19%]
WELLNESS & PREVENTION					
% 60+ with any physical activity within last month		70.1%	[63.05% - 77.10%]	74.5%	[73.36% - 75.59%]
% 60+ met CDC guidelines for muscle-strengthening activity		27.2%	[21.36% - 33.08%]	26.0%	[24.93% - 27.14%]
% 60+ met CDC guidelines for aerobic physical activity		52.0%	[45.41% - 58.64%]	58.6%	[57.33% - 59.82%]
% 60+ met CDC guidelines for both types of physical activities		20.4%	[14.95% - 25.79%]	19.9%	[18.92% - 20.93%]
% 60+ getting recommended hours of sleep		67.5%	[60.64% - 74.30%]	66.4%	[65.16% - 67.60%]
% 60+ injured in a fall within last 12 months		12.3%	[6.99% - 17.64%]	10.4%	[9.65% - 11.16%]
% 65+ had hip fracture	B	2.4%	[1.83% - 2.94%]	3.3%	[3.19% - 3.34%]
% 60+ with self-reported fair or poor health status		13.2%	[8.89% - 17.60%]	16.5%	[15.54% - 17.40%]
% 60+ with 15+ physically unhealthy days last month		12.1%	[7.85% - 16.44%]	12.5%	[11.69% - 13.37%]
% 60+ with physical exam/check-up in past year		87.9%	[82.46% - 93.34%]	86.5%	[85.66% - 87.39%]
% 60+ met CDC preventive health screening goals		47.8%	[38.48% - 57.15%]	40.3%	[38.77% - 41.81%]
% 60+ flu shot past year		67.1%	[60.12% - 74.15%]	59.3%	[58.06% - 60.63%]
% 65+ with pneumonia vaccine		79.7%	[72.76% - 86.58%]	77.8%	[76.56% - 79.12%]
% 60+ with cholesterol screening		96.4%	[94.20% - 98.64%]	95.3%	[94.78% - 95.82%]
% 60+ women with a mammogram within last 2 years	B	88.9%	[82.68% - 95.06%]	79.3%	[77.96% - 80.69%]
% 60+ with colorectal cancer screening		82.5%	[76.01% - 88.90%]	77.0%	[75.92% - 78.16%]
% 60+ with HIV test		16.1%	[10.28% - 21.95%]	13.5%	[12.50% - 14.41%]
% 60+ current smokers		6.1%	[2.99% - 9.21%]	8.1%	[7.38% - 8.79%]
Oral Health					
% 60+ with loss of 6 or more teeth		30.8%	[23.46% - 38.12%]	29.0%	[27.79% - 30.12%]
% 60+ with annual dental exam		80.9%	[74.41% - 87.38%]	75.7%	[74.65% - 76.84%]
NUTRITION/DIET					
% 60+ with 5 or more servings of fruit or vegetables per day		21.6%	[16.19% - 27.09%]	20.2%	[19.23% - 21.21%]
% 60+ self-reported obese		29.0%	[22.62% - 35.34%]	27.2%	[26.03% - 28.35%]
% 65+ clinically diagnosed obese	W	23.6%	[18.74% - 28.36%]	16.7%	[16.56% - 16.94%]
% 65+ with high cholesterol		73.9%	[69.47% - 78.32%]	72.2%	[72.01% - 72.44%]
% 60+ excessive drinking		7.8%	[4.08% - 11.48%]	9.2%	[8.45% - 9.94%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
BEHAVIORAL HEALTH					
% 60+ with 15+ days poor mental health last month		9.4%	[4.64% - 14.12%]	6.9%	[6.29% - 7.61%]
% 65+ with depression	B	24.1%	[19.97% - 28.13%]	28.8%	[28.59% - 29.01%]
% 65+ with anxiety disorders		19.4%	[15.05% - 23.68%]	21.9%	[21.74% - 22.15%]
% 65+ with bipolar disorders		2.8%	[2.11% - 3.46%]	3.1%	[2.97% - 3.14%]
% 65+ with post-traumatic stress disorder		1.4%	[0.86% - 1.86%]	1.4%	[1.35% - 1.48%]
% 65+ with schizophrenia & other psychotic disorders	B	3.8%	[3.01% - 4.49%]	4.9%	[4.81% - 5.02%]
% 65+ with personality disorders		0.7%	[0.31% - 1.01%]	1.1%	[1.00% - 1.11%]
% 65+ with substance use disorders (drug use +/- alcohol abuse)		3.8%	[2.20% - 5.48%]	5.5%	[5.41% - 5.64%]
% 65+ with tobacco use disorders		9.2%	[5.98% - 12.42%]	10.4%	[10.23% - 10.54%]
CHRONIC DISEASE					
% 65+ with Alzheimer's disease or related dementias	B	8.2%	[5.80% - 10.67%]	12.0%	[11.89% - 12.19%]
% 65+ with diabetes		28.1%	[23.19% - 33.04%]	28.2%	[27.97% - 28.41%]
% 65+ with stroke	B	7.5%	[5.17% - 9.93%]	10.8%	[10.64% - 10.92%]
% 65+ with chronic obstructive pulmonary disease		17.3%	[13.78% - 20.86%]	20.5%	[20.36% - 20.73%]
% 65+ with asthma		13.9%	[10.62% - 17.27%]	13.0%	[12.89% - 13.21%]
% 65+ with hypertension		71.5%	[66.93% - 76.11%]	70.2%	[70.02% - 70.46%]
% 65+ ever had a heart attack	B	2.7%	[1.21% - 4.10%]	4.5%	[4.37% - 4.56%]
% 65+ with ischemic heart disease		32.2%	[27.12% - 37.24%]	34.3%	[34.12% - 34.58%]
% 65+ with congestive heart failure	B	13.7%	[10.34% - 17.11%]	17.7%	[17.57% - 17.93%]
% 65+ with atrial fibrillation	B	10.6%	[7.72% - 13.49%]	14.4%	[14.28% - 14.60%]
% 65+ with peripheral vascular disease	B	9.7%	[6.91% - 12.57%]	14.7%	[14.51% - 14.84%]
% 65+ with osteoarthritis/rheumatoid arthritis		48.0%	[42.51% - 53.45%]	49.1%	[48.82% - 49.32%]
% 65+ with osteoporosis		14.4%	[11.28% - 17.62%]	17.4%	[17.22% - 17.57%]
% 65+ with leukemias and lymphomas		2.0%	[1.43% - 2.50%]	2.0%	[1.91% - 2.05%]
% 65+ with lung cancer		1.7%	[1.19% - 2.15%]	1.6%	[1.54% - 1.66%]
% 65+ with colon cancer		2.1%	[1.57% - 2.63%]	2.4%	[2.37% - 2.51%]
% 65+ women with breast cancer		8.9%	[5.06% - 12.70%]	9.8%	[9.59% - 9.96%]
% 65+ women with endometrial cancer	B	1.0%	[0.51% - 1.55%]	1.7%	[1.60% - 1.76%]
% 65+ men with prostate cancer		14.7%	[9.69% - 19.71%]	11.5%	[11.24% - 11.68%]
% 65+ with benign prostatic hyperplasia		36.2%	[29.60% - 42.83%]	36.8%	[36.47% - 37.14%]
% 65+ with HIV/AIDS		0.04%	[0.02% - 0.07%]	0.05%	[0.04% - 0.06%]
% 65+ with hypothyroidism		21.0%	[17.09% - 24.98%]	20.8%	[20.62% - 20.99%]
% 65+ with anemia	B	31.4%	[26.99% - 35.85%]	37.3%	[37.12% - 37.57%]
% 65+ with chronic kidney disease		21.0%	[16.78% - 25.26%]	22.3%	[22.14% - 22.54%]
% 65+ with liver diseases		7.9%	[4.99% - 10.78%]	6.9%	[6.75% - 6.99%]
% 65+ with fibromyalgia, chronic pain and fatigue		17.2%	[13.17% - 21.20%]	18.6%	[18.42% - 18.80%]
% 65+ with migraine and other chronic headache		3.4%	[2.65% - 4.08%]	4.0%	[3.95% - 4.15%]
% 65+ with epilepsy		2.4%	[1.77% - 3.02%]	2.1%	[2.00% - 2.14%]
% 65+ with traumatic brain injury		1.4%	[0.93% - 1.87%]	1.1%	[1.06% - 1.16%]

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders			N/A	0.03%	[0.02% - 0.04%]
% 65+ with glaucoma		23.2%	[19.16% - 27.21%]	22.9%	[22.74% - 23.13%]
% 65+ with cataract		61.3%	[56.35% - 66.16%]	61.2%	[61.00% - 61.47%]
% 65+ with pressure ulcer or chronic ulcer		5.0%	[2.85% - 7.24%]	7.1%	[6.99% - 7.24%]
% 65+ with 4+ (out of 15) chronic conditions	B	48.7%	[43.78% - 53.62%]	54.4%	[54.19% - 54.66%]
% 65+ with 0 chronic conditions		8.6%	[5.62% - 11.56%]	10.3%	[10.12% - 10.42%]
LIVING WITH DISABILITY					
% 65+ with clinical diagnosis of deafness or hearing impairment		14.1%	[10.49% - 17.65%]	14.4%	[14.23% - 14.56%]
% 65+ with clinical diagnosis of blindness or visual impairment		0.6%	[0.30% - 0.88%]	0.9%	[0.81% - 0.90%]
% 65+ with clinical diagnosis of mobility impairments		3.0%	[2.34% - 3.68%]	3.2%	[3.14% - 3.31%]
ACCESS TO CARE					
Medicare (65+ population)					
% Medicare managed care enrollees		9.5%	[7.19% - 11.81%]	7.9%	[7.8% - 8.02%]
% dually eligible for Medicare and Medicaid	*	3.1%	[1.70% - 4.41%]	7.5%	[7.40% - 7.62%]
% 60+ with a regular doctor		97.2%	[95.16% - 99.25%]	95.8%	[95.34% - 96.33%]
% 60+ who did not see doctor when needed due to cost		2.7%	[0.42% - 4.95%]	5.4%	[4.75% - 5.97%]
SERVICE UTILIZATION					
Physician visits per year	*	7.9	[7.1 - 8.6]	6.3	[6.2 - 6.3]
Emergency room visits/1000 persons 65+ years per year	*	447	[340 - 555]	572	[566 - 578]
Part D monthly prescription fills per person per year		46.6	[42.8 - 50.5]	49.1	[48.9 - 49.3]
Home health visits per year		1.7	[0.9 - 2.5]	2.5	[2.4 - 2.5]
Durable medical equipment claims per year		2.0	[1.5 - 2.5]	2.0	[2.0 - 2.0]
Inpatient hospital stays/1000 persons 65+ years per year		199	[133 - 265]	237	[234 - 241]
Medicare inpatient hospital readmissions (as % of admissions)		15.9%	[2.95% - 28.86%]	15.8%	[15.41% - 16.22%]
# skilled nursing facility stays/1000 persons 65+ years per year	*	45	[20 - 70]	76	[74 - 77]
% 65+ getting Medicaid long term services and supports	*	1.3%	[0.40% - 2.18%]	3.7%	[3.59% - 3.74%]
COMMUNITY VARIABLES & CIVIC ENGAGEMENT					
% 60+ who used Internet in last month		80.1%	[74.98% - 85.29%]	77.6%	[76.53% - 78.64%]
SAFETY & TRANSPORTATION					
% 60+ who always drive wearing a seatbelt		83.0%	[76.70% - 89.27%]	77.1%	[76.02% - 78.18%]

Notes. ¹ Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

² All community and state margins of error are 95% confidence intervals.

³ The margin of error of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last month, and did not see doctor when needed due to cost.

Questions or suggestions? Beth.Dugan@umb.edu