Conway (Carroll)



HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²		STATE ESTIMATE	MARGIN OF ERROR ²	
POPULATION CHARACTERISTICS							
Age-sex adjusted 1-year mortality rate		4.5%	[3.47% - 5.55%]	4.1%	[4.02% - 4.19%]	
WELLNESS & PREVENTION							
% 60+ with any physical activity within last month	В	80.6%	[75.68% - 85.59%]	74.5%	[73.36% - 75.59%]	
% 60+ met CDC guidelines for muscle-strengthening activity		28.5%	[21.37% - 35.54%]	26.0%	[24.93% - 27.14%]	
% 60+ met CDC guidelines for aerobic physical activity		65.4%	[57.93% - 72.77%]	58.6%	[57.33% - 59.82%]	
% 60+ met CDC guidelines for both types of physical activities		21.0%	[14.50% - 27.51%]	19.9%	[18.92% - 20.93%]	
% 60+ getting recommended hours of sleep		68.4%	[62.62% - 74.25%]	66.4%	[65.16% - 67.60%]	
% 60+ injured in a fall within last 12 months		11.1%	[6.95% - 15.22%]	10.4%	[9.65% - 11.16%]	
% 65+ had hip fracture	В	2.7%	[2.14% - 3.19%]	3.3%	[3.19% - 3.34%]	
% 60+ with self-reported fair or poor health status		14.9%	[10.66% - 19.18%]	16.5%	[15.54% - 17.40%]	
% 60+ with 15+ physically unhealthy days last month		12.6%	[8.79% - 16.50%]	12.5%	[11.69% - 13.37%]	
% 60+ with physical exam/check-up in past year		82.7%	[77.25% - 88.19%]	86.5%	[85.66% - 87.39%]	
% 60+ met CDC preventive health screening goals		39.7%	[32.41% - 47.03%]	40.3%	[38.77% - 41.81%]	
% 60+ flu shot past year		58.8%	[51.87% - 65.65%]	59.3%	[58.06% - 60.63%]	
% 65+ with pneumonia vaccine		72.5%	[65.14% - 79.91%]	77.8%	[76.56% - 79.12%]	
% 60+ with cholesterol screening		92.7%	[88.66% - 96.64%]	95.3%	[94.78% - 95.82%]	
% 60+ women with a mammogram within last 2 years		78.4%	[71.52% - 85.33%]	79.3%	[77.96% - 80.69%]	
% 60+ with colorectal cancer screening		73.0%	[66.37% - 79.71%]	77.0%	[75.92% - 78.16%]	
% 60+ with HIV test		11.4%	[6.96% - 15.84%]	13.5%	[12.50% - 14.41%]	
% 60+ current smokers		6.6%	[3.63% - 9.51%]	8.1%	[7.38% - 8.79%]	
Oral Health							
% 60+ with loss of 6 or more teeth		29.4%	[23.27% - 35.48%]	29.0%	[27.79% - 30.12%]	
% 60+ with annual dental exam		76.4%	[70.95% - 81.81%]	75.7%	[74.65% - 76.84%]	
NUTRITION/DIET							
% 60+ with 5 or more servings of fruit or vegetables per day		17.8%	[12.06% - 23.57%]	20.2%	[19.23% - 21.21%]	
% 60+ self-reported obese		25.9%	[19.27% - 32.58%]	27.2%	[26.03% - 28.35%]	
% 65+ clinically diagnosed obese		15.0%	[13.05% - 17.03%]	16.7%	[16.56% - 16.94%]	
% 65+ with high cholesterol		71.3%	[68.90% - 73.78%]	72.2%	[72.01% - 72.44%]	
% 60+ excessive drinking		12.3%	[8.03% - 16.50%]	9.2%	[8.45% - 9.94%]	

Conway (Carroll) Page 1

68 OH with 15+ days poor mental health last month B 3.8% 1.61% 5.94% 6.9% 6.29% 7.81% 66 SH with depression 29.5% [27.14% 31.94% 28.9% 28.9% 28.9% 28.9% 28.9% 28.9% 28.9% 29.0% 3.14% 29.0% 3.14% 29.0% 3.14% 4.28%	HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE	MARGIN OF ERROR ²		STATE ESTIMATE	MARGIN OF ERROR ²
86 Set with depression 29.5% 27.14% 31.94% 28.8% 28.99% 29.01% 66 Set with anwiety disorders 23.3% 20.92% -25.00% 21.9% 21.74% -22.15% 66 Set with pipolar disorders 1.4% 1.28% -2.25% 3.1% 12.97% 3.14% 66 Set with personality disorders 8 3.9% 1.32% -4.63% 4.9% 1.48% 1.05% 1.14% 1.05% 1.14% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.11% 1.00% 1.00% 1.11% 1.00% 1.00% 1.00% 1.00% 1.11% 1.00% 1.00% 1.11% 1.00% 1.00% 1.11% 1.00% 1.00%	BEHAVIORAL HEALTH						
8. 65+ with naviety disorders	% 60+ with 15+ days poor mental health last month	В	3.8%	[1.61% -	5.94%]	6.9%	[6.29% - 7.61%]
8. 65+ with bipolaid disorders B 1.8% [1.28% - 2.25%] 3.1% [2.97% - 3.14%	% 65+ with depression		29.5%	[27.14% - 3	1.94%]	28.8%	[28.59% - 29.01%]
66 + with post-traumatic stress disorder 1.4% [0.95% - 1.81%] 1.4% [1.36% - 1.48%] 6.4% 1.4% [1.36% - 1.48%] 6.4% 1.4% 6.95% - 1.81%] 1.4% 6.95% - 1.48% 6.65% 1.48% 5.02% 6.65% 1.05% - 1.13% 1.0% - 1.14% 5.02% 6.65% 1.05% - 1.29% 1.0% 1.0% - 1.14% 5.62% 6.65% 1.05% 1.05% - 1.29% 1.0% 1.05% - 1.05% 1.05% <td>% 65+ with anxiety disorders</td> <td></td> <td>23.3%</td> <td>[20.92% - 2</td> <td>5.60%]</td> <td>21.9%</td> <td>[21.74% - 22.15%]</td>	% 65+ with anxiety disorders		23.3%	[20.92% - 2	5.60%]	21.9%	[21.74% - 22.15%]
86 65+ with schizophrenia & other psychotic disorders B 3.9% [3.26% - 4.63%] 4.9% [4.81% - 5.02% 66 5+ with personality disorders B 0.7% [0.37% - 0.99%] 1.1% [1.00% - 1.11% 66 5+ with substance use disorders (drug use +/or alcohol abuse) 5.5% [4.25% - 6.78%] 5.5% [5.41% - 5.64% 66 5+ with lobacco use disorders 10.9% [9.16% - 12.69%] 10.4% [10.23% - 10.54% CHRONIC DISEASE 8 5.5% [2.351% - 28.40%] 12.0% [11.89% - 12.19% 66 5+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.89% - 12.19% 66 5+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.89% - 12.19% 66 5+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [12.19% - 12.19% 66 5+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [12.98% - 12.29% 66 5+ with Stroke 9.8% [8.27% - 11.33%] 10.0% [10.25% - 12.97%] 12.0% 12.19% <td>% 65+ with bipolar disorders</td> <td>В</td> <td>1.8%</td> <td>[1.28% -</td> <td>2.25%]</td> <td>3.1%</td> <td>[2.97% - 3.14%]</td>	% 65+ with bipolar disorders	В	1.8%	[1.28% -	2.25%]	3.1%	[2.97% - 3.14%]
B 0.7% 0.37% 0.99% 1.1% 1.00% 1.11% 1.00% 1.11% 65+ with substance use disorders (drug use +/or alcohol abuse) 5.5% 4.25% 6.78% 5.5% 5.41% 5.64% 685+ with substance use disorders 10.9% 9.16% 12.69% 10.4% 10.23% 10.54% 10.54% 65+ with hobacco use disorders 10.9% 9.16% 12.69% 10.4% 10.23% 10.54% 10.54% 65+ with hobacco use disorders 11.3% 9.65% 12.97% 12.07% 12.0% 11.89% 12.19% 65+ with diabetes 26.0% 23.51% 28.40% 28.2% 27.97% 28.41% 65+ with diabetes 26.0% 23.51% 28.2% 10.64% 10.92% 65+ with chronic obstructive pulmonary disease 22.6% 20.41% 24.77% 20.5% 20.36% 20.373% 65+ with chronic obstructive pulmonary disease 22.6% 20.41% 24.77% 20.5% 20.36% 20.373% 65+ with hypertension 71.0% 68.55% 73.46% 70.2% 70.02% 70.02% 70.046% 65+ with isothemic heart disease 32.0% 29.48% 34.61% 34.3% 34.12% 34.58% 65+ with isothemic heart disease 32.0% 29.48% 34.61% 34.3% 34.12% 34.58% 65+ with congestive heart failure 18.3% 16.23% 20.36% 17.7% 17.57% 17.93% 65+ with peripheral vascular disease 8 11.9% 10.17% 13.67% 14.4% 14.26% 46.60% 45+ with congestive heart failure 8 13.2% 11.14% 14.97% 17.4% 17.22% 17.57% 65+ with leukemias and lymphomas 1.5% 11.11% 1.98% 1.98% 2.98% 2.95	% 65+ with post-traumatic stress disorder		1.4%	[0.95% -	1.81%]	1.4%	[1.35% - 1.48%]
\$ 65+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.39% - 10.54% \$ 65+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.39% - 12.19% \$ 65+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.39% - 12.19% \$ 65+ with Alzheimer's disease or related dementias 11.3% [9.65% - 12.97%] 12.0% [11.39% - 12.19% \$ 65+ with diabetes 26.0% [23.51% - 28.40%] 28.2% [27.97% - 28.41% \$ 65+ with drivatic obstructive pulmonary disease 22.6% [20.41% - 24.77%] 10.0% [10.64% - 10.92% \$ 65+ with strivate 12.4% [10.65% - 14.11%] 13.0% [12.89% - 13.21% \$ 65+ with hypertension 71.0% [88.55% - 73.46%] 70.2% [70.02% - 70.46% \$ 65+ with hypertension 71.0% [88.55% - 73.46%] 70.2% [70.02% - 70.46% \$ 65+ with hypertension 71.0% [88.55% - 73.46%] 70.2% [70.02% - 70.46% \$ 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% \$ 65+ with peripheral vascular disease 8 11.9% [10.47% - 13.67%] 14.4% [14.28% - 14.60% \$ 65+ with peripheral vascular disease 8 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.89% \$ 65+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% \$ 65+ with hypertension 14.	% 65+ with schizophrenia & other psychotic disorders	В	3.9%	[3.26% -	4.63%]	4.9%	[4.81% - 5.02%]
## 65+ with tobacco use disorders 10.9%	% 65+ with personality disorders	В	0.7%	[0.37% -	0.99%]	1.1%	[1.00% - 1.11%]
## CHRONIC DISEASE ## 65+ with Alzheimer's disease or related dementias ## 11.3%	% 65+ with substance use disorders (drug use +/or alcohol abuse)		5.5%	[4.25% -	6.78%]	5.5%	[5.41% - 5.64%]
% 65+ with Alzheimer's disease or related dementias 11.3% [9.65% · 12.97%] 12.0% [11.89% · 12.19% % 65+ with diabetes 26.0% [2.351% · 28.40%] 28.2% [2.797% · 28.41% % 65+ with stroke 9.8% [8.27% · 11.33%] 10.8% [10.64% · 10.92% % 65+ with chronic obstructive pulmonary disease 22.6% [20.41% · 24.47%] 20.5% [20.36% · 20.73% % 65+ with hypertension 71.0% [68.55% · 14.11%] 13.0% [1.80% · 13.21% % 65+ with is pertension 71.0% [68.55% · 73.46%] 70.2% [70.02% · 70.46% % 65+ with congestive heart failure 18.3% [16.23% · 20.36%] 17.7% [4.57% · 17.93% % 65+ with strial fibililation 14.0% [12.16% · 15.77%] 14.4% [14.28% · 14.60% % 65+ with congestive heart failure 18.3% [10.15% · 13.67%] 14.4% [14.50% · 17.7%] 14.50% % 65+ with atrial fibililation 14.0% [12.16% · 15.7%] 14.4% [14.60% % 65+ with atrial fibililation 14.0% [10.17% · 13.67%] 14.4% [14.80% · 14.60% % 65+ with atrial fibililation 18.11.9% [10.17% · 1	% 65+ with tobacco use disorders		10.9%	[9.16% - 1	2.69%]	10.4%	[10.23% - 10.54%]
66 5+ with diabetes 26.0% [23.51% - 28.40%] 28.2% [27.97% - 28.41%] 68.41 (27.97% - 28.41%) 68.41 (27.97% - 28.41%) 68.41 (27.97% - 28.41%) 68.41 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 68.42 (27.97% - 28.41%) 69.42 (27.97%	CHRONIC DISEASE						
8 65+ with stroke 9.8% [8.27% - 11.33%] 10.8% [10.64% - 10.92% 65+ with chronic obstructive pulmonary disease 22.6% [2.0.41% - 24.77%] 20.5% [2.0.36% - 20.73% 65+ with chronic obstructive pulmonary disease 22.6% [2.0.41% - 24.77%] 20.5% [2.0.36% - 20.73% 65+ with chronic obstructive pulmonary disease 22.6% [2.0.41% - 24.77%] 20.5% [2.0.36% - 20.73% 65+ with sthma 12.4% [10.65% - 14.11%] 13.0% [12.69% - 13.21% 65+ with hypertension 71.0% [68.55% - 73.46%] 70.2% [7.0.2% 70.46% 65+ were had a heart attack 5.0% [68.55% - 73.46%] 70.2% [7.0.2% 70.46% 65+ with ischemic heart disease 32.0% [2.9.48% - 34.61%] 34.3% [34.12% - 34.56% 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% 65+ with atrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84% 65+ with osteoprorosis B 13.2% [11.44% - 14.91%] 17.4% [17.22% - 17.57% 65+ with luckemias and lymphomas 1.5% [11.14% - 1.91%] 17.4% [17.22% - 17.57% 65+ with luckemias and lymphomas 1.5% [11.44% - 14.91%] 17.4% [17.22% - 17.57% 65+ with luckemias and core B 11.5% [2.13% - 3.22%] 2.4% [2.33% - 2.51% 65+ with benign prostatic cancer 19.2% [7.14% - 11.26%] 9.8% [1.59% - 9.96% 65+ women with prostate cancer 19.2% [7.14% - 11.26%] 9.8% [1.69% - 3.14% 65+ with benign prostatic hyperplasia 37.7% [3.40%] 17.0% [3.64%] 17.0% [3.6	% 65+ with Alzheimer's disease or related dementias		11.3%	[9.65% - 1	2.97%]	12.0%	[11.89% - 12.19%]
66 6+ with chronic obstructive pulmonary disease 22.6% 20.41% - 24.7% 20.5% 20.36% 20.73% 66 6+ with asthma 12.4% 10.65% - 14.11% 13.0% 12.89% - 13.21% 6.20.36% 20.36% 20.73% 66 6+ with hypertension 71.0% 6.68.55% - 73.46% 70.2% 70.02% - 70.46% 6.69 with portan at attack 5.0% 3.393% - 6.17% 4.5% 4.57% 4.56% 66 6+ with ischemic heart disease 32.0% [29.48% - 34.61%] 34.3% [34.12% - 34.68%] 6.65 with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93%] 66 6+ with atrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.51% - 14.86%] 6.65 with peripheral vascular disease 8 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.88%] -14.60% 66 6+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.88%] -14.60% 66 5+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.52% - 49.32%] 66 5+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 1.0% [17.27% - 1.45%] 1.0% [15.66%]	% 65+ with diabetes		26.0%	[23.51% - 2	8.40%]	28.2%	[27.97% - 28.41%]
% 65+ with asthma 12.4% [10.65% - 14.11%] 13.0% [12.89% - 13.21% % 65+ with hypertension 71.0% [68.55% - 73.46%] 70.2% [70.02% - 70.46% % 65+ ever had a heart attack 5.0% [3.93% - 6.17%] 4.5% [4.37% - 4.56% % 65+ with ischemic heart disease 32.0% [29.48% - 34.61%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% % 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.4% [14.28% - 14.60% % 65+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32% % 65+ with leukemias and lymphomas 1.5% [11.14% - 1.98%] 2.0% [17.57% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.25% % 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.55% % 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.96% - 1.66% % 65+ with leukemias and lymphomas 1.5% [1	% 65+ with stroke		9.8%	[8.27% - 1	1.33%]	10.8%	[10.64% - 10.92%]
% 65+ with hypertension 71.0% [68.55% - 73.46% 70.2% [70.02% - 70.46% % 65+ ever had a heart attack 5.0% [3.93% - 6.17%] 4.5% [4.37% - 4.56% % 65+ with sichemic heart disease 32.0% [29.48% - 34.61%] 34.12% - 34.58% % 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% % 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% % 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.4% [14.28% - 14.84% % 65+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32% % 65+ with bosteoporosis B 13.2% [11.14% - 14.91%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.11% - 1.99%] 2.0% (54.94% - 14.91%] 1.7% [1.54% - 1.56%	% 65+ with chronic obstructive pulmonary disease		22.6%	[20.41% - 2	4.77%]	20.5%	[20.36% - 20.73%]
% 65+ ever had a heart attack 5.0% [3.93% - 6.17%] 4.5% [4.37% - 4.56% % 65+ with ischemic heart disease 32.0% [29.48% - 34.61%] 34.3% [34.12% - 34.58% % 65+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% % 65+ with attrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% % 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84% % 65+ with osteoporosis B 13.2% [11.44% - 14.91%] 17.4% [17.22% - 17.57% % 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05% % 65+ with long cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% % 65+ with long cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% % 65+ with long cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% % 65+ with long cancer B 1.1% [0.74% - 1.33%] 1.1% [1.54% - 1.66% % 65+ women with breast cancer <	% 65+ with asthma		12.4%	[10.65% - 1	4.11%]	13.0%	[12.89% - 13.21%]
66+ with ischemic heart disease 32.0% [29.48% - 34.61%] 34.3% [34.12% - 34.58% 66+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93% 65+ with atrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84% 65+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32% 65+ with osteoporosis B 13.2% [11.14% - 14.91%] 17.4% [17.22% - 17.57% 65+ with loug cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% 65+ with colon cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ women with breast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with prostate cancer 11.5% [9.10% - 1.39%] 11.24% - 11.26% 9.8% [9.59% - 9.96% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with hypothyroidism 19.	% 65+ with hypertension		71.0%	[68.55% - 7	3.46%]	70.2%	[70.02% - 70.46%]
66+ with congestive heart failure 18.3% [16.23% - 20.36%] 17.7% [17.57% - 17.93%] 66+ with atrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60%] 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84%] 65+ with osteoparthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32%] 65+ with osteoporosis B 13.2% [11.14% - 14.91%] 17.4% [17.22% - 17.57%] 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05%] 65+ with colon cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66%] 65+ women with breast cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51%] 65+ women with prostate cancer 11.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76%] 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 3.714%] 65+ with hypothyroidism 19.1% [1.707% - 21.16%] 20.8% [20.62% - 20.99%] 65+ with hippothyroidism 19.1% <	% 65+ ever had a heart attack		5.0%	[3.93% -	6.17%]	4.5%	[4.37% - 4.56%]
66+ with atrial fibrillation 14.0% [12.16% - 15.77%] 14.4% [14.28% - 14.60% 66+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84% 66+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32% 65+ with osteoporosis B 13.2% [11.44% - 14.91%] 17.4% [17.22% - 17.57% 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05% 65+ with colon cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% 65+ women with breast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 2.51% 65+ women with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with hirornoic kidney disease B <t< td=""><td>% 65+ with ischemic heart disease</td><td></td><td>32.0%</td><td>[29.48% - 3</td><td>4.61%]</td><td>34.3%</td><td>[34.12% - 34.58%]</td></t<>	% 65+ with ischemic heart disease		32.0%	[29.48% - 3	4.61%]	34.3%	[34.12% - 34.58%]
% 65+ with peripheral vascular disease B 11.9% [10.17% - 13.67%] 14.7% [14.51% - 14.84%	% 65+ with congestive heart failure		18.3%	[16.23% - 2	0.36%]	17.7%	[17.57% - 17.93%]
66+ with osteoarthritis/rheumatoid arthritis 50.6% [47.76% - 53.40%] 49.1% [48.82% - 49.32% 66+ with osteoporosis B 13.2% [11.44% - 14.91%] 17.4% [17.22% - 17.57% 66+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05% 66+ with lung cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% 65+ with colon cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ women with preast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with prostate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% - 1.76% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% 65+ with pypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 17.4% [15.88% - 20	% 65+ with atrial fibrillation		14.0%	[12.16% - 1	5.77%]	14.4%	[14.28% - 14.60%]
8 65+ with osteoporosis B 13.2% [11.44% - 14.91%] 17.4% [17.22% - 17.57% 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05% 65+ with lung cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% 65+ with colon cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ with osteoporosis and site of the colon cancer 3.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ women with breast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with endometrial cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with postate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ with hypothyroidism 1.9.1% [1.707% - 21.16%] 20.8% [2.062% - 20.99% 65+ with hypothyroidism 1.9.1% [1.707% - 21.16%] 20.8% [2.062% - 20.99% 65+ with chronic kidney disease 1.9% [1.54% - 1.43%] 1.6% [1.54% - 22.54% 65+ with fibromyalgia, chronic pain and fatigue 1.9% [1.54% - 1.43%] 1.6% [1.54% - 1.43%] 1.6% [1.84% - 1.88% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy	% 65+ with peripheral vascular disease	В	11.9%	[10.17% - 1	3.67%]	14.7%	[14.51% - 14.84%]
% 65+ with leukemias and lymphomas 1.5% [1.11% - 1.98%] 2.0% [1.91% - 2.05% % 65+ with lung cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% % 65+ with colon cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% % 65+ women with breast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% % 65+ women with prostate cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% % 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% % 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% % 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% % 65+ with chronic kidney disease B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% % 65+ with liver diseases B 4.7% [3.58% - 20.09%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% % 65+ with epilepsy B 1.3% [0.93% - 1.76%	% 65+ with osteoarthritis/rheumatoid arthritis		50.6%	[47.76% - 5	3.40%]	49.1%	[48.82% - 49.32%]
8 65+ with lung cancer B 1.1% [0.74% - 1.43%] 1.6% [1.54% - 1.66% 65+ with colon cancer 2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ with colon cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with breast cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with endometrial cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ men with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epil	% 65+ with osteoporosis	В	13.2%	[11.44% - 1	4.91%]	17.4%	[17.22% - 17.57%]
2.7% [2.13% - 3.22%] 2.4% [2.37% - 2.51% 65+ with colon cancer 9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with breast cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ men with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.9	% 65+ with leukemias and lymphomas		1.5%	[1.11% -	1.98%]	2.0%	[1.91% - 2.05%]
9.2% [7.14% - 11.26%] 9.8% [9.59% - 9.96% 65+ women with breast cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 65+ men with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy	% 65+ with lung cancer	В	1.1%	[0.74% -	1.43%]	1.6%	[1.54% - 1.66%]
26 65+ women with endometrial cancer 1.9% [1.25% - 2.52%] 1.7% [1.60% - 1.76% 26 65+ men with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 26 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 26 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% 26 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 26 65+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 26 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 26 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 26 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 26 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 26 65+ with epilepsy B 1.3% <td>% 65+ with colon cancer</td> <td></td> <td>2.7%</td> <td>[2.13% -</td> <td>3.22%]</td> <td>2.4%</td> <td>[2.37% - 2.51%]</td>	% 65+ with colon cancer		2.7%	[2.13% -	3.22%]	2.4%	[2.37% - 2.51%]
11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ men with prostate cancer 11.5% [9.10% - 13.94%] 11.5% [11.24% - 11.68% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93%	% 65+ women with breast cancer		9.2%	[7.14% - 1	1.26%]	9.8%	[9.59% - 9.96%]
37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with benign prostatic hyperplasia 37.7% [34.03% - 41.44%] 36.8% [36.47% - 37.14% 65+ with HIV/AIDS 0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 65+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [% 65+ women with endometrial cancer		1.9%	[1.25% -	2.52%]	1.7%	[1.60% - 1.76%]
0.04% [0.02% - 0.06%] 0.05% [0.04% - 0.06%] 0.05% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% 565+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy	% 65+ men with prostate cancer		11.5%	[9.10% - 1	3.94%]	11.5%	[11.24% - 11.68%]
% 65+ with hypothyroidism 19.1% [17.07% - 21.16%] 20.8% [20.62% - 20.99% % 65+ with anemia 34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% % 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% % 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% % 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% % 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14%	% 65+ with benign prostatic hyperplasia		37.7%	[34.03% - 4	1.44%]	36.8%	[36.47% - 37.14%]
34.7% [32.23% - 37.22%] 37.3% [37.12% - 37.57% 37.57% 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 65+ with fibromyalgia, chronic pain and fatigue B 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 65+ with migraine and other chronic headache B 1.3% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.76%] 2.1% [0.93% - 1.	% 65+ with HIV/AIDS		0.04%	[0.02% -	0.06%]	0.05%	[0.04% - 0.06%]
8 65+ with chronic kidney disease B 17.4% [15.36% - 19.47%] 22.3% [22.14% - 22.54% 8 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99% 8 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% 8 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 8 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14%	% 65+ with hypothyroidism		19.1%	[17.07% - 2	1.16%]	20.8%	[20.62% - 20.99%]
8 65+ with liver diseases B 4.7% [3.55% - 5.90%] 6.9% [6.75% - 6.99%	% 65+ with anemia		34.7%	[32.23% - 3	7.22%]	37.3%	[37.12% - 37.57%]
% 65+ with fibromyalgia, chronic pain and fatigue 18.0% [15.88% - 20.09%] 18.6% [18.42% - 18.80% % 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% % 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14%	% 65+ with chronic kidney disease	В	17.4%	[15.36% - 1	9.47%]	22.3%	[22.14% - 22.54%]
% 65+ with migraine and other chronic headache 3.6% [2.93% - 4.28%] 4.0% [3.95% - 4.15% 8 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14% 9 65+ with epilepsy	% 65+ with liver diseases	В	4.7%	[3.55% -	5.90%]	6.9%	[6.75% - 6.99%]
% 65+ with epilepsy B 1.3% [0.93% - 1.76%] 2.1% [2.00% - 2.14%	% 65+ with fibromyalgia, chronic pain and fatigue		18.0%	[15.88% - 2	0.09%]	18.6%	[18.42% - 18.80%]
	% 65+ with migraine and other chronic headache		3.6%	[2.93% -	4.28%]	4.0%	[3.95% - 4.15%]
% 65+ with traumatic brain injury 1.3% [0.90% - 1.67%] 1.1% [1.06% - 1.16%	% 65+ with epilepsy	В	1.3%	[0.93% -	1.76%]	2.1%	[2.00% - 2.14%]
	% 65+ with traumatic brain injury		1.3%	[0.90% -	1.67%]	1.1%	[1.06% - 1.16%]

Conway (Carroll) Page 2

HEALTHY AGING INDICATORS	BETTER / WORSE STATE RATE ¹	COMMUNITY ESTIMATE		MARGIN OF ERROR ²	STATE ESTIMATE	MARGIN OF ERROR ²
% 65+ with autism spectrum disorders				N/A	0.03%	[0.02% - 0.04
% 65+ with glaucoma	W	32.0%	[29.55% - 34.45%]	22.9%	[22.74% - 23.13
% 65+ with cataract		62.9%	[60.32% - 65.53%]	61.2%	[61.00% - 61.47
% 65+ with pressure ulcer or chronic ulcer		6.9%	[5.54% - 8.24%]	7.1%	[6.99% - 7.24
% 65+ with 4+ (out of 15) chronic conditions		52.8%	[50.14% - 55.46%]	54.4%	[54.19% - 54.66
% 65+ with 0 chronic conditions		10.0%	[8.39% - 11.70%]	10.3%	[10.12% - 10.42
LIVING WITH DISABILITY						
% 65+ with clinical diagnosis of deafness or hearing impairment	В	12.3%	[10.56% - 14.10%]	14.4%	[14.23% - 14.56
% 65+ with clinical diagnosis of blindness or visual impairment	В	0.5%	[0.24% - 0.71%]	0.9%	[0.81% - 0.90
% 65+ with clinical diagnosis of mobility impairments		2.8%	[2.19% - 3.31%]	3.2%	[3.14% - 3.31
ACCESS TO CARE						
Medicare (65+ population)						
% Medicare managed care enrollees		8.4%	[7.08% - 9.73%]	7.9%	[7.8% - 8.02
% dually eligible for Medicare and Medicaid		8.8%	[7.46% - 10.18%]	7.5%	[7.40% - 7.62
% 60+ with a regular doctor		94.2%	[90.66% - 97.67%]	95.8%	[95.34% - 96.33
% 60+ who did not see doctor when needed due to cost		6.0%	[3.00% - 8.93%]	5.4%	[4.75% - 5.97
SERVICE UTILIZATION						
Physician visits per year	*	2.9	[2.7 - 3.2]	6.3	[6.2 - 6.3
Emergency room visits/1000 persons 65+ years per year		621	[548 - 693]	572	[566 - 578
Part D monthly prescription fills per person per year		47.6	[45.2 - 50.0]	49.1	[48.9 - 49.3
Home health visits per year	*	1.6	[1.2 - 2.1]	2.5	[2.4 - 2.5
Durable medical equipment claims per year		2.2	[1.9 - 2.5]	2.0	[2.0 - 2.0
Inpatient hospital stays/1000 persons 65+ years per year		217	[186 - 248]	237	[234 - 241
Medicare inpatient hospital readmissions (as % of admissions)		13.6%	[9.45% - 17.79%]	15.8%	[15.41% - 16.22
# skilled nursing facility stays/1000 persons 65+ years per year		91	[71 - 112]	76	[74 - 77
% 65+ getting Medicaid long term services and supports		4.4%	[3.43% - 5.39%]	3.7%	[3.59% - 3.74
COMMUNITY VARIABLES & CIVIC ENGAGEMENT						
% 60+ who used Internet in last month		80.0%	[74.84% - 85.09%]	77.6%	[76.53% - 78.64
SAFETY & TRANSPORTATION						
% 60+ who always drive wearing a seatbelt		75.0%	[69.20% - 80.87%]	77.1%	[76.02% - 78.18

Questions or suggestions? Beth.Dugan@umb.edu

Conway (Carroll) Page 3

Notes.

Community-state differences where the margin of error do not overlap each other are noted B, W, or *.

All community and state margins of error are 95% confidence intervals.

The margin of erroe of the estimate is larger relative to the indicator value for current smokers, excessive drinking, 15+ days poor mental health last montth, and did not see doctor when needed due to cost.