

## Plainfield (Windham)



| HEALTHY AGING INDICATORS                                   | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>POPULATION CHARACTERISTICS</b>                          |   |                       |                                 |                   |                                 |
| Population 65 years or older                               |   |                       |                                 |                   |                                 |
| % 65-74 years  |   | 58.9%                 | [ 51.33% - 66.53% ]             | 55.7%             | [ 55.33% - 56.11% ]             |
| % 75-84 years  |   | 26.6%                 | [ 20.26% - 33.03% ]             | 29.1%             | [ 28.75% - 29.45% ]             |
| % 85 years or older  |   | 14.4%                 | [ 9.36% - 19.49% ]              | 15.2%             | [ 14.93% - 15.43% ]             |
| % 65+ population who are female                            |   | 62.0%                 | [ 55.01% - 68.98% ]             | 56.8%             | [ 56.36% - 57.18% ]             |
| % 85+ population who are female                            |   | 74.1%                 | [ 58.19% - 89.99% ]             | 66.6%             | [ 65.94% - 67.18% ]             |
| Race and ethnicity of the population 65+                   |   |                       |                                 |                   |                                 |
| % White  | *   | 97.5%                 | [ 91.45% - 103.55% ]            | 88.4%             | [ 87.85% - 88.93% ]             |
| % African American   | *   | 0.5%                  | [ -1.69% - 2.74% ]              | 6.8%              | [ 6.63% - 6.88% ]               |
| % Asian  |   | 1.1%                  | [ -1.17% - 3.34% ]              | 2.3%              | [ 2.20% - 2.35% ]               |
| % Hispanic/Latino  |   | 5.3%                  | [ -0.66% - 11.35% ]             | 5.8%              | [ 5.66% - 5.84% ]               |
| Marital status of the population 65+                       |   |                       |                                 |                   |                                 |
| % married  |   | 50.3%                 | [ 43.96% - 56.64% ]             | 53.8%             | [ 53.46% - 54.12% ]             |
| % divorced/separated                                       |   | 16.3%                 | [ 8.55% - 24.11% ]              | 14.9%             | [ 14.58% - 15.29% ]             |
| % widowed  |   | 26.4%                 | [ 19.83% - 33.01% ]             | 24.2%             | [ 23.77% - 24.57% ]             |
| % never married  |   | 7.0%                  | [ 2.41% - 11.49% ]              | 7.1%              | [ 6.85% - 7.36% ]               |
| Education of the population 65+                            |   |                       |                                 |                   |                                 |
| % with less than high school education                     | *   | 24.0%                 | [ 17.16% - 30.82% ]             | 14.7%             | [ 14.30% - 15.04% ]             |
| % with high school or some college                         | *   | 66.8%                 | [ 59.04% - 74.59% ]             | 53.3%             | [ 52.80% - 53.74% ]             |
| % with college degree                                      | *   | 9.2%                  | [ 4.71% - 13.68% ]              | 32.1%             | [ 31.72% - 32.40% ]             |
| % with graduate or professional degree                     | *   | 3.0%                  | [ -0.12% - 6.09% ]              | 16.8%             | [ 16.58% - 17.09% ]             |
| % of 65+ population who speak only English at home         | *   | 93.5%                 | [ 85.46% - 101.54% ]            | 84.3%             | [ 83.76% - 84.83% ]             |
| % of 65+ population who are veterans of military service   |   | 21.9%                 | [ 14.73% - 29.07% ]             | 17.5%             | [ 17.21% - 17.69% ]             |
| % of 60+ who are LGBT (County)                             |   | 3.9%                  | [ 2.02% - 5.73% ]               | 2.5%              | [ 2.17% - 2.74% ]               |
| Age-sex adjusted 1-year mortality rate                     | W   | 5.2%                  | [ 4.55% - 5.80% ]               | 4.1%              | [ 4.08% - 4.18% ]               |
| <b>WELLNESS</b>  |   |                       |                                 |                   |                                 |
| % 60+ getting the recommended hours of sleep               |   | 65.6%                 | [ 58.34% - 72.90% ]             | 65.2%             | [ 64.24% - 66.16% ]             |
| % 60+ doing any physical activity within last month        | W   | 63.0%                 | [ 56.72% - 69.36% ]             | 75.0%             | [ 74.20% - 75.79% ]             |
| % 60+ met CDC guidelines for muscle-strengthening activity |   | 22.2%                 | [ 16.50% - 27.87% ]             | 27.6%             | [ 26.62% - 28.52% ]             |
| % 60+ met CDC guidelines for aerobic physical activity     |   | 52.8%                 | [ 45.93% - 59.66% ]             | 58.1%             | [ 56.98% - 59.11% ]             |
| % 60+ with fair or poor health status                      |   | 17.4%                 | [ 12.53% - 22.27% ]             | 17.6%             | [ 16.90% - 18.31% ]             |
| % 60+ with 15+ physically unhealthy days in last month     |   | 13.4%                 | [ 8.90% - 17.90% ]              | 12.4%             | [ 11.74% - 12.96% ]             |

| HEALTHY AGING INDICATORS                                     | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>FALLS</b>   |   |                       |                                 |                   |                                 |
| % 60+ who fell within last year                              |   | 32.4%                 | [ 24.68% - 40.10% ]             | 26.3%             | [ 25.38% - 27.18% ]             |
| % 60+ who were injured in a fall within last year            |   | 10.9%                 | [ 6.40% - 15.46% ]              | 9.8%              | [ 9.22% - 10.46% ]              |
| % 65+ had hip fracture                                       |   | 3.1%                  | [ 2.58% - 3.65% ]               | 3.7%              | [ 3.63% - 3.75% ]               |
| <b>PREVENTION</b>  |   |                       |                                 |                   |                                 |
| % 60+ with physical exam/check-up in last year               |   | 91.3%                 | [ 87.54% - 94.98% ]             | 88.5%             | [ 87.93% - 89.09% ]             |
| % 60+ flu shot in last year                                  |   | 58.7%                 | [ 52.31% - 65.08% ]             | 57.5%             | [ 56.60% - 58.43% ]             |
| % 60+ with pneumonia vaccine                                 |   | 67.6%                 | [ 61.49% - 73.80% ]             | 63.9%             | [ 62.97% - 64.79% ]             |
| % 60+ with shingles vaccine                                  |   | 28.0%                 | [ 19.89% - 36.13% ]             | 33.5%             | [ 32.43% - 34.52% ]             |
| % 60+ women with a mammogram within last 2 years             |   | 81.2%                 | [ 73.16% - 89.15% ]             | 79.7%             | [ 78.56% - 80.79% ]             |
| % 60+ with colorectal cancer screening                       |   | 83.7%                 | [ 78.24% - 89.14% ]             | 78.2%             | [ 77.29% - 79.05% ]             |
| % 60+ with HIV test  |   | 17.4%                 | [ 12.17% - 22.61% ]             | 16.1%             | [ 15.40% - 16.83% ]             |
| % 60+ met CDC preventive health screening goals              |   | 45.7%                 | [ 35.67% - 55.68% ]             | 39.6%             | [ 38.38% - 40.80% ]             |
| <b>NUTRITION &amp; DIET</b>                                  |   |                       |                                 |                   |                                 |
| % 60+ with 5 or more servings of fruit or vegetables per day | W   | 13.1%                 | [ 8.43% - 17.74% ]              | 19.9%             | [ 19.00% - 20.72% ]             |
| % 60+ stressed about buying food in last month               |   | 9.7%                  | [ 5.44% - 13.89% ]              | 8.6%              | [ 7.95% - 9.17% ]               |
| % 60+ self-reported obese                                    | W   | 34.5%                 | [ 28.03% - 40.91% ]             | 26.1%             | [ 25.27% - 26.93% ]             |
| % 65+ clinically diagnosed obesity                           | W   | 28.1%                 | [ 26.41% - 29.70% ]             | 22.5%             | [ 22.35% - 22.66% ]             |
| % 65+ with high cholesterol                                  |   | 77.8%                 | [ 76.31% - 79.21% ]             | 77.0%             | [ 76.85% - 77.15% ]             |
| % 60+ with cholesterol screening                             |   | 96.7%                 | [ 94.69% - 98.66% ]             | 97.1%             | [ 96.74% - 97.42% ]             |
| <b>ORAL HEALTH</b>   |   |                       |                                 |                   |                                 |
| % 60+ with dental insurance                                  |   | 58.5%                 | [ 46.82% - 70.24% ]             | 59.2%             | [ 57.60% - 60.76% ]             |
| % 60+ with annual dental exam                                |   | 73.5%                 | [ 66.96% - 80.11% ]             | 80.0%             | [ 79.13% - 80.76% ]             |
| % 60+ with loss of 6 or more teeth                           | W   | 34.9%                 | [ 27.19% - 42.52% ]             | 23.8%             | [ 22.96% - 24.69% ]             |
| <b>CHRONIC DISEASE</b>                                       |   |                       |                                 |                   |                                 |
| % 65+ with Alzheimer's disease or related dementias          |   | 14.2%                 | [ 12.98% - 15.33% ]             | 14.4%             | [ 14.26% - 14.49% ]             |
| % 65+ with anemia  | B   | 45.8%                 | [ 44.16% - 47.53% ]             | 50.6%             | [ 50.41% - 50.76% ]             |
| % 65+ with asthma  | W   | 17.5%                 | [ 16.20% - 18.75% ]             | 14.1%             | [ 13.96% - 14.19% ]             |
| % 65+ with atrial fibrillation                               | W   | 17.5%                 | [ 16.26% - 18.76% ]             | 16.1%             | [ 15.94% - 16.18% ]             |
| % 65+ with autism spectrum disorder                          | *   | 0.07%                 | [ 0.04% - 0.10% ]               | 0.12%             | [ 0.11% - 0.14% ]               |
| % 65+ with benign prostatic hyperplasia (men)                |   | 43.0%                 | [ 40.42% - 45.51% ]             | 42.8%             | [ 42.49% - 43.01% ]             |
| % 65+ with breast cancer (women)                             | B   | 9.3%                  | [ 8.04% - 10.65% ]              | 11.4%             | [ 11.25% - 11.53% ]             |
| % 65+ with cataract  |   | 65.2%                 | [ 63.58% - 66.89% ]             | 64.8%             | [ 64.67% - 65.00% ]             |
| % 65+ with chronic kidney disease                            |   | 31.1%                 | [ 29.40% - 32.70% ]             | 29.9%             | [ 29.70% - 30.03% ]             |
| % 65+ with chronic obstructive pulmonary disease             | W   | 31.2%                 | [ 29.61% - 32.73% ]             | 21.0%             | [ 20.90% - 21.17% ]             |
| % 65+ with colon cancer                                      |   | 2.7%                  | [ 2.30% - 3.12% ]               | 2.7%              | [ 2.65% - 2.75% ]               |
| % 65+ with congestive heart failure                          |   | 24.1%                 | [ 22.59% - 25.58% ]             | 22.6%             | [ 22.43% - 22.72% ]             |
| % 65+ with diabetes  |   | 35.3%                 | [ 33.55% - 37.00% ]             | 33.8%             | [ 33.59% - 33.93% ]             |
| % 65+ with endometrial cancer (women)                        |   | 1.6%                  | [ 1.14% - 1.98% ]               | 2.0%              | [ 1.97% - 2.10% ]               |

| HEALTHY AGING INDICATORS                                | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| % 65+ with epilepsy                                     |   | 2.9%                  | [ 2.45% - 3.42% ]               | 2.9%              | [ 2.82% - 2.95% ]               |
| % 65+ with fibromyalgia, chronic pain, and fatigue      |   | 27.7%                 | [ 26.06% - 29.30% ]             | 26.6%             | [ 26.42% - 26.73% ]             |
| % 65+ with glaucoma                                     | B   | 23.7%                 | [ 22.31% - 25.15% ]             | 28.3%             | [ 28.11% - 28.41% ]             |
| % 65+ ever had a heart attack                           | W   | 6.4%                  | [ 5.61% - 7.20% ]               | 4.5%              | [ 4.40% - 4.53% ]               |
| % 65+ with HIV/AIDS                                     | *   | 0.11%                 | [ 0.08% - 0.15% ]               | 0.21%             | [ 0.19% - 0.23% ]               |
| % 65+ with hypertension                                 | W   | 80.8%                 | [ 79.39% - 82.17% ]             | 76.2%             | [ 76.07% - 76.37% ]             |
| % 65+ with hypothyroidism                               | B   | 22.3%                 | [ 20.90% - 23.68% ]             | 25.1%             | [ 24.99% - 25.28% ]             |
| % 65+ with ischemic heart disease                       | W   | 44.6%                 | [ 42.82% - 46.41% ]             | 40.7%             | [ 40.55% - 40.90% ]             |
| % 65+ with leukemias and lymphomas                      |   | 3.0%                  | [ 2.52% - 3.48% ]               | 2.7%              | [ 2.60% - 2.71% ]               |
| % 65+ with liver disease                                |   | 10.0%                 | [ 8.91% - 11.07% ]              | 9.5%              | [ 9.37% - 9.59% ]               |
| % 65+ with lung cancer                                  |   | 2.2%                  | [ 1.83% - 2.62% ]               | 1.9%              | [ 1.81% - 1.90% ]               |
| % 65+ with migraine and other chronic headache          |   | 5.2%                  | [ 4.41% - 6.05% ]               | 5.2%              | [ 5.13% - 5.29% ]               |
| % 65+ with osteoarthritis or rheumatoid arthritis       | W   | 57.3%                 | [ 55.46% - 59.10% ]             | 54.2%             | [ 53.97% - 54.34% ]             |
| % 65+ with osteoporosis                                 |   | 19.5%                 | [ 18.21% - 20.80% ]             | 20.2%             | [ 20.11% - 20.37% ]             |
| % 65+ with peripheral vascular disease                  |   | 21.1%                 | [ 19.66% - 22.51% ]             | 19.9%             | [ 19.72% - 19.99% ]             |
| % 65+ with pressure ulcer or chronic ulcer              |   | 9.3%                  | [ 8.31% - 10.30% ]              | 9.2%              | [ 9.11% - 9.31% ]               |
| % 65+ with prostate cancer (men)                        | B   | 9.2%                  | [ 7.81% - 10.68% ]              | 13.1%             | [ 12.93% - 13.28% ]             |
| % 65+ with stroke                                       | W   | 13.5%                 | [ 12.33% - 14.57% ]             | 11.9%             | [ 11.80% - 12.01% ]             |
| % 65+ with traumatic brain injury                       | B   | 0.9%                  | [ 0.68% - 1.22% ]               | 1.3%              | [ 1.25% - 1.33% ]               |
| % 65+ with 4+ (out of 15) chronic conditions            | W   | 66.3%                 | [ 64.63% - 67.92% ]             | 61.8%             | [ 61.61% - 61.95% ]             |
| % 65+ with 0 chronic conditions                         | W   | 6.1%                  | [ 5.24% - 6.98% ]               | 7.2%              | [ 7.10% - 7.29% ]               |
| <b>BEHAVIORAL HEALTH</b>                                |   |                       |                                 |                   |                                 |
| % 65+ with opioid use disorder                          |   | 1.7%                  | [ 1.35% - 2.12% ]               | 2.2%              | [ 2.10% - 2.21% ]               |
| % 65+ with substance use disorder                       | W   | 8.2%                  | [ 7.14% - 9.20% ]               | 6.8%              | [ 6.66% - 6.85% ]               |
| % 60+ who used marijuana in last month                  |   | N/A                   |                                 | 2.9%              | [ 2.30% - 3.53% ]               |
| % 60+ excessive drinking                                |   | 11.4%                 | [ 7.44% - 15.29% ]              | 9.2%              | [ 8.64% - 9.71% ]               |
| % 65+ with tobacco use disorder                         | W   | 15.1%                 | [ 13.76% - 16.46% ]             | 10.1%             | [ 9.94% - 10.17% ]              |
| % 60+ current smokers                                   | W   | 13.1%                 | [ 8.58% - 17.54% ]              | 7.7%              | [ 7.20% - 8.17% ]               |
| % 60+ ever used E-Cigarettes in last month              |   | 10.3%                 | [ 6.51% - 13.99% ]              | 6.4%              | [ 5.89% - 6.80% ]               |
| <b>MENTAL HEALTH</b>                                    |   |                       |                                 |                   |                                 |
| % 60+ who reported receiving adequate emotional support |   | 74.6%                 | [ 66.52% - 82.73% ]             | 78.3%             | [ 76.70% - 79.88% ]             |
| % 60+ with 15+ days poor mental health in last month    |   | 8.0%                  | [ 4.50% - 11.41% ]              | 6.9%              | [ 6.41% - 7.37% ]               |
| % 65+ with depression                                   | W   | 35.0%                 | [ 33.40% - 36.62% ]             | 30.3%             | [ 30.10% - 30.41% ]             |
| % 65+ with anxiety disorder                             | W   | 33.4%                 | [ 31.68% - 35.08% ]             | 25.5%             | [ 25.35% - 25.66% ]             |
| % 65+ with bipolar disorder                             | W   | 5.7%                  | [ 5.03% - 6.38% ]               | 3.9%              | [ 3.78% - 3.92% ]               |
| % 65+ with post-traumatic stress disorder               | W   | 2.0%                  | [ 1.56% - 2.41% ]               | 1.3%              | [ 1.20% - 1.29% ]               |
| % 65+ with schizophrenia & other psychotic disorder     |   | 4.8%                  | [ 4.16% - 5.34% ]               | 4.6%              | [ 4.53% - 4.68% ]               |
| % 65+ with personality disorder                         |   | 2.1%                  | [ 1.71% - 2.53% ]               | 2.6%              | [ 2.49% - 2.61% ]               |

| HEALTHY AGING INDICATORS   | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>LIVING WITH DISABILITY</b>                                    |   |                       |                                 |                   |                                 |
| % 65+ with self-reported hearing difficulty                      |   | 17.2%                 | [ 10.64% - 23.68% ]             | 12.4%             | [ 12.08% - 12.71% ]             |
| % 65+ with clinical diagnosis of deafness or hearing impairment  |   | 15.1%                 | [ 13.86% - 16.34% ]             | 15.7%             | [ 15.60% - 15.85% ]             |
| % 65+ with self-reported vision difficulty                       |   | 7.7%                  | [ 2.58% - 12.80% ]              | 4.8%              | [ 4.55% - 5.04% ]               |
| % 65+ with clinical diagnosis of blindness or visual impairment  |   | 1.0%                  | [ 0.75% - 1.28% ]               | 1.1%              | [ 1.05% - 1.12% ]               |
| % 65+ with self-reported cognition difficulty                    |   | 9.3%                  | [ 3.61% - 15.01% ]              | 7.6%              | [ 7.34% - 7.87% ]               |
| % 65+ with self-reported ambulatory difficulty                   |   | 26.1%                 | [ 18.30% - 33.86% ]             | 19.1%             | [ 18.71% - 19.52% ]             |
| % 65+ with clinical diagnosis of mobility impairment             |   | 4.5%                  | [ 3.77% - 5.21% ]               | 4.0%              | [ 3.95% - 4.09% ]               |
| % 65+ with self-reported self-care difficulty                    |   | 5.4%                  | [ 1.81% - 8.97% ]               | 7.5%              | [ 7.25% - 7.77% ]               |
| % 65+ with self-reported independent living difficulty           |   | 15.1%                 | [ 8.60% - 21.68% ]              | 13.6%             | [ 13.23% - 13.93% ]             |
| <b>CAREGIVING</b>  |   |                       |                                 |                   |                                 |
| % of 60+ who provide care to a family/friend in last month       |   | 27.4%                 | [ 17.40% - 37.45% ]             | 21.7%             | [ 19.87% - 23.57% ]             |
| % of grandparents raising grandchildren                          |   | 1.4%                  | [ -0.31% - 3.12% ]              | 0.8%              | [ 0.73% - 0.86% ]               |
| % of grandparents who live with grandchildren                    |   | 3.2%                  | [ 1.20% - 5.25% ]               | 2.7%              | [ 2.62% - 2.83% ]               |
| <b>ACCESS TO CARE</b>  |   |                       |                                 |                   |                                 |
| % 65+ dually eligible for Medicare and Medicaid                  | *   | 32.1%                 | [ 30.87% - 33.27% ]             | 22.1%             | [ 21.96% - 22.17% ]             |
| % 65+ Medicare managed care enrollees                            |   | 28.7%                 | [ 27.52% - 29.85% ]             | 29.9%             | [ 29.76% - 29.99% ]             |
| % 60+ with a regular doctor                                      |   | 96.1%                 | [ 94.06% - 98.24% ]             | 96.0%             | [ 95.68% - 96.39% ]             |
| % 60+ who did not see a doctor when needed due to cost           |   | 6.1%                  | [ 3.06% - 9.21% ]               | 4.5%              | [ 4.14% - 4.92% ]               |
| <b>SERVICE UTILIZATION</b>                                       |   |                       |                                 |                   |                                 |
| # physician visits per year                                      | *   | 7.9                   | [ 7.64 - 8.15 ]                 | 8.6               | [ 8.60 - 8.66 ]                 |
| # emergency room visits/1000 persons 65+ years annually          | *   | 876.0                 | [ 802.36 - 949.59 ]             | 636.7             | [ 631.77 - 641.56 ]             |
| # Part D monthly prescription fills per person annually          | *   | 57.6                  | [ 56.25 - 58.94 ]               | 50.3              | [ 50.15 - 50.39 ]               |
| # home health visits annually                                    | *   | 3.5                   | [ 3.00 - 3.98 ]                 | 4.1               | [ 4.02 - 4.14 ]                 |
| # durable medical equipment claims annually                      | *   | 2.4                   | [ 2.15 - 2.59 ]                 | 1.9               | [ 1.88 - 1.91 ]                 |
| # inpatient hospital stays/1000 persons 65+ years annually       |   | 299.4                 | [ 273.23 - 325.57 ]             | 273.3             | [ 270.72 - 275.80 ]             |
| % Medicare inpatient hospital readmissions (as % of admissions)  |   | 16.1%                 | [ 13.67% - 18.51% ]             | 17.2%             | [ 16.89% - 17.47% ]             |
| # skilled nursing facility stays/1000 persons 65+ years annually |   | 119.9                 | [ 105.19 - 134.69 ]             | 104.7             | [ 103.23 - 106.08 ]             |
| % 65+ getting Medicaid long term services and supports           | *   | 7.7%                  | [ 6.99% - 8.36% ]               | 5.5%              | [ 5.48% - 5.60% ]               |
| % 65+ hospice users  | *   | 1.8%                  | [ 1.48% - 2.20% ]               | 2.8%              | [ 2.70% - 2.79% ]               |
| % 65+ hospice users as % of decedents                            | *   | 28.0%                 | [ 22.56% - 33.39% ]             | 45.5%             | [ 44.85% - 46.14% ]             |
| Median hospice days per hospice user (65+, deceased)             |   | 12.0                  | [ 8.0 - 17.0 ]                  | 10.0              | [ 10.00 - 10.00 ]               |
| Median hospice payment (Medicare + other) per hospice user       |   | \$3,401               | [ \$2,640 - \$4,216 ]           | \$3,741           | [ \$3,650 - \$3,816 ]           |
| <b>COMMUNITY</b>   |   |                       |                                 |                   |                                 |
| % 60+ who used Internet in last month                            |   | 68.1%                 | [ 61.77% - 74.43% ]             | 75.1%             | [ 74.32% - 75.89% ]             |
| <b>TRANSPORTATION</b>  |   |                       |                                 |                   |                                 |
| % 65+ who own a motor vehicle                                    |   | 85.2%                 | [ 79.87% - 90.43% ]             | 87.3%             | [ 87.15% - 87.49% ]             |
| % 60+ who always drive or ride wearing a seatbelt                |   | 94.3%                 | [ 91.78% - 96.80% ]             | 93.0%             | [ 92.53% - 93.49% ]             |
| % 60+ who drove under the influence of drinking in last month    |   | N/A                   |                                 | 2.2%              | [ 1.76% - 2.56% ]               |

| HEALTHY AGING INDICATORS                                  | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>HOUSING</b>  |   |                       |                                 |                   |                                 |
| % 65+ population who live alone                           |   | 29.3%                 | [ 23.17% - 35.34% ]             | 28.4%             | [ 27.99% - 28.89% ]             |
| % 60+ own home  |   | 73.8%                 | [ 69.70% - 77.92% ]             | 76.9%             | [ 76.70% - 77.17% ]             |
| % 60+ homeowners who have mortgage                        |   | 46.8%                 | [ 37.48% - 56.11% ]             | 47.1%             | [ 46.58% - 47.64% ]             |
| % 60+ stressed about paying rent/mortgage in last month   |   | 15.9%                 | [ 10.65% - 21.22% ]             | 17.4%             | [ 16.52% - 18.19% ]             |
| % 65+ households (renter) spend >35% of income on housing |   | 37.0%                 | [ 13.32% - 60.77% ]             | 44.0%             | [ 42.55% - 45.42% ]             |
| % 65+ households (owner) spend >35% of income on housing  |   | 23.4%                 | [ 11.81% - 35.07% ]             | 29.5%             | [ 28.89% - 30.15% ]             |
| % 65+ moved within same county in last year               |   | 7.3%                  | [ 1.92% - 12.73% ]              | 3.7%              | [ 3.48% - 3.89% ]               |
| % 65+ moved from different county in last year            |   | 0.2%                  | [ -1.12% - 1.50% ]              | 0.8%              | [ 0.71% - 0.90% ]               |
| % 65+ moved from different state in last year             |   | 1.6%                  | [ -1.60% - 4.81% ]              | 0.9%              | [ 0.82% - 1.05% ]               |
| % of vacant homes in community                            |   | 10.3%                 | [ 6.35% - 14.29% ]              | 9.6%              | [ 9.29% - 9.87% ]               |
| <b>ECONOMIC</b>   |   |                       |                                 |                   |                                 |
| % 60+ receiving food benefits in last year                |   | 12.0%                 | [ 6.88% - 17.19% ]              | 10.9%             | [ 10.53% - 11.21% ]             |
| % 65+ employed in last year                               | *   | 14.7%                 | [ 8.70% - 20.74% ]              | 24.8%             | [ 24.48% - 25.15% ]             |
| % 65+ with income below the poverty line in last year     |   | 10.4%                 | [ 4.07% - 16.79% ]              | 7.0%              | [ 6.79% - 7.30% ]               |
| % 65+ households with annual income < \$20,000            |   | 22.4%                 | [ 13.78% - 30.96% ]             | 17.1%             | [ 16.57% - 17.55% ]             |
| % 65+ households with annual income \$20,000-\$49,999     | *   | 45.7%                 | [ 32.15% - 59.33% ]             | 31.0%             | [ 30.35% - 31.62% ]             |
| % 65+ households with annual income \$50,000-\$99,999     |   | 24.6%                 | [ 16.55% - 32.58% ]             | 27.6%             | [ 27.04% - 28.17% ]             |
| % 65+ households with annual income \$100,000+            | *   | 7.3%                  | [ 2.90% - 11.76% ]              | 24.4%             | [ 23.82% - 24.89% ]             |

Notes. 1 We use B, W, or \* to denote significant differences in rates where the margins of error do not overlap (B=rate is better than the comparison, W=rate is worse than comparison, \*=rate is different than comparison).

2 All community and state margins of error are 95% confidence intervals.

Questions or suggestions? [Beth.Dugan@umb.edu](mailto:Beth.Dugan@umb.edu)