

## Shelton (Fairfield)



| HEALTHY AGING INDICATORS                                   | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>POPULATION CHARACTERISTICS</b>                          |   |                       |                                 |                   |                                 |
| Population 65 years or older                               |   |                       |                                 |                   |                                 |
| % 65-74 years  |   | 55.8%                 | [ 52.00% - 59.70% ]             | 55.7%             | [ 55.33% - 56.11% ]             |
| % 75-84 years  |   | 27.5%                 | [ 23.81% - 31.12% ]             | 29.1%             | [ 28.75% - 29.45% ]             |
| % 85 years or older  |   | 16.7%                 | [ 12.99% - 20.38% ]             | 15.2%             | [ 14.93% - 15.43% ]             |
| % 65+ population who are female                            |   | 57.2%                 | [ 52.80% - 61.52% ]             | 56.8%             | [ 56.36% - 57.18% ]             |
| % 85+ population who are female                            |   | 73.4%                 | [ 61.23% - 85.66% ]             | 66.6%             | [ 65.94% - 67.18% ]             |
| Race and ethnicity of the population 65+                   |   |                       |                                 |                   |                                 |
| % White  | *   | 97.4%                 | [ 93.74% - 101.11% ]            | 88.4%             | [ 87.85% - 88.93% ]             |
| % African American   | *   | 0.6%                  | [ -0.14% - 1.30% ]              | 6.8%              | [ 6.63% - 6.88% ]               |
| % Asian  | *   | 1.0%                  | [ 0.09% - 1.83% ]               | 2.3%              | [ 2.20% - 2.35% ]               |
| % Hispanic/Latino  | *   | 3.0%                  | [ 1.45% - 4.48% ]               | 5.8%              | [ 5.66% - 5.84% ]               |
| Marital status of the population 65+                       |   |                       |                                 |                   |                                 |
| % married  | *   | 58.1%                 | [ 55.27% - 60.94% ]             | 53.8%             | [ 53.46% - 54.12% ]             |
| % divorced/separated                                       | *   | 11.8%                 | [ 9.08% - 14.43% ]              | 14.9%             | [ 14.58% - 15.29% ]             |
| % widowed  |   | 27.8%                 | [ 23.62% - 31.96% ]             | 24.2%             | [ 23.77% - 24.57% ]             |
| % never married  | *   | 2.4%                  | [ 1.29% - 3.42% ]               | 7.1%              | [ 6.85% - 7.36% ]               |
| Education of the population 65+                            |   |                       |                                 |                   |                                 |
| % with less than high school education                     | *   | 10.1%                 | [ 7.36% - 12.77% ]              | 14.7%             | [ 14.30% - 15.04% ]             |
| % with high school or some college                         | *   | 61.2%                 | [ 57.86% - 64.52% ]             | 53.3%             | [ 52.80% - 53.74% ]             |
| % with college degree                                      |   | 28.7%                 | [ 25.40% - 32.09% ]             | 32.1%             | [ 31.72% - 32.40% ]             |
| % with graduate or professional degree                     |   | 14.4%                 | [ 11.86% - 16.90% ]             | 16.8%             | [ 16.58% - 17.09% ]             |
| % of 65+ population who speak only English at home         |   | 87.3%                 | [ 82.12% - 92.48% ]             | 84.3%             | [ 83.76% - 84.83% ]             |
| % of 65+ population who are veterans of military service   |   | 16.3%                 | [ 13.33% - 19.31% ]             | 17.5%             | [ 17.21% - 17.69% ]             |
| % of 60+ who are LGBT (County)                             |   | 2.1%                  | [ 1.50% - 2.60% ]               | 2.5%              | [ 2.17% - 2.74% ]               |
| Age-sex adjusted 1-year mortality rate                     |   | 4.3%                  | [ 3.82% - 4.70% ]               | 4.1%              | [ 4.08% - 4.18% ]               |
| <b>WELLNESS</b>  |   |                       |                                 |                   |                                 |
| % 60+ getting the recommended hours of sleep               | W   | 57.1%                 | [ 50.70% - 63.57% ]             | 65.2%             | [ 64.24% - 66.16% ]             |
| % 60+ doing any physical activity within last month        |   | 71.8%                 | [ 66.68% - 76.83% ]             | 75.0%             | [ 74.20% - 75.79% ]             |
| % 60+ met CDC guidelines for muscle-strengthening activity |   | 27.4%                 | [ 21.02% - 33.75% ]             | 27.6%             | [ 26.62% - 28.52% ]             |
| % 60+ met CDC guidelines for aerobic physical activity     |   | 53.8%                 | [ 46.44% - 61.17% ]             | 58.1%             | [ 56.98% - 59.11% ]             |
| % 60+ with fair or poor health status                      |   | 19.6%                 | [ 15.16% - 24.11% ]             | 17.6%             | [ 16.90% - 18.31% ]             |
| % 60+ with 15+ physically unhealthy days in last month     |   | 10.3%                 | [ 6.96% - 13.62% ]              | 12.4%             | [ 11.74% - 12.96% ]             |

| HEALTHY AGING INDICATORS                                     | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>FALLS</b>   |   |                       |                                 |                   |                                 |
| % 60+ who fell within last year                              |   | 24.0%                 | [ 18.20% - 29.79% ]             | 26.3%             | [ 25.38% - 27.18% ]             |
| % 60+ who were injured in a fall within last year            |   | 8.1%                  | [ 4.23% - 12.05% ]              | 9.8%              | [ 9.22% - 10.46% ]              |
| % 65+ had hip fracture                                       |   | 3.9%                  | [ 3.34% - 4.42% ]               | 3.7%              | [ 3.63% - 3.75% ]               |
| <b>PREVENTION</b>  |   |                       |                                 |                   |                                 |
| % 60+ with physical exam/check-up in last year               |   | 89.5%                 | [ 86.21% - 92.76% ]             | 88.5%             | [ 87.93% - 89.09% ]             |
| % 60+ flu shot in last year                                  |   | 57.1%                 | [ 51.47% - 62.81% ]             | 57.5%             | [ 56.60% - 58.43% ]             |
| % 60+ with pneumonia vaccine                                 |   | 59.1%                 | [ 53.22% - 64.91% ]             | 63.9%             | [ 62.97% - 64.79% ]             |
| % 60+ with shingles vaccine                                  | W   | 26.1%                 | [ 20.01% - 32.24% ]             | 33.5%             | [ 32.43% - 34.52% ]             |
| % 60+ women with a mammogram within last 2 years             |   | 77.2%                 | [ 70.03% - 84.41% ]             | 79.7%             | [ 78.56% - 80.79% ]             |
| % 60+ with colorectal cancer screening                       |   | 74.5%                 | [ 68.73% - 80.21% ]             | 78.2%             | [ 77.29% - 79.05% ]             |
| % 60+ with HIV test  |   | 13.6%                 | [ 9.45% - 17.70% ]              | 16.1%             | [ 15.40% - 16.83% ]             |
| % 60+ met CDC preventive health screening goals              |   | 31.5%                 | [ 24.31% - 38.76% ]             | 39.6%             | [ 38.38% - 40.80% ]             |
| <b>NUTRITION &amp; DIET</b>                                  |   |                       |                                 |                   |                                 |
| % 60+ with 5 or more servings of fruit or vegetables per day |   | 18.5%                 | [ 12.69% - 24.27% ]             | 19.9%             | [ 19.00% - 20.72% ]             |
| % 60+ stressed about buying food in last month               |   | 8.4%                  | [ 4.78% - 11.97% ]              | 8.6%              | [ 7.95% - 9.17% ]               |
| % 60+ self-reported obese                                    |   | 29.4%                 | [ 24.18% - 34.54% ]             | 26.1%             | [ 25.27% - 26.93% ]             |
| % 65+ clinically diagnosed obesity                           | B   | 19.6%                 | [ 18.29% - 20.85% ]             | 22.5%             | [ 22.35% - 22.66% ]             |
| % 65+ with high cholesterol                                  | W   | 79.3%                 | [ 78.04% - 80.58% ]             | 77.0%             | [ 76.85% - 77.15% ]             |
| % 60+ with cholesterol screening                             |   | 98.0%                 | [ 96.26% - 99.82% ]             | 97.1%             | [ 96.74% - 97.42% ]             |
| <b>ORAL HEALTH</b>   |   |                       |                                 |                   |                                 |
| % 60+ with dental insurance                                  |   | 56.8%                 | [ 46.80% - 66.75% ]             | 59.2%             | [ 57.60% - 60.76% ]             |
| % 60+ with annual dental exam                                |   | 77.0%                 | [ 71.44% - 82.46% ]             | 80.0%             | [ 79.13% - 80.76% ]             |
| % 60+ with loss of 6 or more teeth                           |   | 26.0%                 | [ 20.04% - 31.89% ]             | 23.8%             | [ 22.96% - 24.69% ]             |
| <b>CHRONIC DISEASE</b>                                       |   |                       |                                 |                   |                                 |
| % 65+ with Alzheimer's disease or related dementias          |   | 14.6%                 | [ 13.56% - 15.66% ]             | 14.4%             | [ 14.26% - 14.49% ]             |
| % 65+ with anemia  | W   | 56.0%                 | [ 54.51% - 57.54% ]             | 50.6%             | [ 50.41% - 50.76% ]             |
| % 65+ with asthma  |   | 13.1%                 | [ 12.13% - 14.14% ]             | 14.1%             | [ 13.96% - 14.19% ]             |
| % 65+ with atrial fibrillation                               |   | 16.7%                 | [ 15.65% - 17.83% ]             | 16.1%             | [ 15.94% - 16.18% ]             |
| % 65+ with autism spectrum disorder                          |   | 0.09%                 | [ 0.06% - 0.13% ]               | 0.12%             | [ 0.11% - 0.14% ]               |
| % 65+ with benign prostatic hyperplasia (men)                |   | 45.3%                 | [ 42.97% - 47.56% ]             | 42.8%             | [ 42.49% - 43.01% ]             |
| % 65+ with breast cancer (women)                             |   | 10.9%                 | [ 9.70% - 12.16% ]              | 11.4%             | [ 11.25% - 11.53% ]             |
| % 65+ with cataract  | B   | 63.0%                 | [ 61.55% - 64.53% ]             | 64.8%             | [ 64.67% - 65.00% ]             |
| % 65+ with chronic kidney disease                            |   | 28.3%                 | [ 26.90% - 29.73% ]             | 29.9%             | [ 29.70% - 30.03% ]             |
| % 65+ with chronic obstructive pulmonary disease             |   | 20.8%                 | [ 19.63% - 22.03% ]             | 21.0%             | [ 20.90% - 21.17% ]             |
| % 65+ with colon cancer                                      |   | 2.9%                  | [ 2.46% - 3.44% ]               | 2.7%              | [ 2.65% - 2.75% ]               |
| % 65+ with congestive heart failure                          |   | 22.6%                 | [ 21.27% - 23.83% ]             | 22.6%             | [ 22.43% - 22.72% ]             |
| % 65+ with diabetes  | W   | 37.2%                 | [ 35.66% - 38.75% ]             | 33.8%             | [ 33.59% - 33.93% ]             |
| % 65+ with endometrial cancer (women)                        |   | 2.7%                  | [ 2.04% - 3.32% ]               | 2.0%              | [ 1.97% - 2.10% ]               |

| HEALTHY AGING INDICATORS                                | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| % 65+ with epilepsy                                     |   | 2.4%                  | [ 1.89% - 2.82% ]               | 2.9%              | [ 2.82% - 2.95% ]               |
| % 65+ with fibromyalgia, chronic pain, and fatigue      |   | 26.3%                 | [ 24.88% - 27.68% ]             | 26.6%             | [ 26.42% - 26.73% ]             |
| % 65+ with glaucoma                                     |   | 27.0%                 | [ 25.71% - 28.36% ]             | 28.3%             | [ 28.11% - 28.41% ]             |
| % 65+ ever had a heart attack                           |   | 4.8%                  | [ 4.22% - 5.46% ]               | 4.5%              | [ 4.40% - 4.53% ]               |
| % 65+ with HIV/AIDS                                     |   | 0.21%                 | [ 0.15% - 0.27% ]               | 0.21%             | [ 0.19% - 0.23% ]               |
| % 65+ with hypertension                                 | W   | 78.5%                 | [ 77.17% - 79.75% ]             | 76.2%             | [ 76.07% - 76.37% ]             |
| % 65+ with hypothyroidism                               | W   | 29.8%                 | [ 28.38% - 31.12% ]             | 25.1%             | [ 24.99% - 25.28% ]             |
| % 65+ with ischemic heart disease                       | W   | 43.3%                 | [ 41.73% - 44.89% ]             | 40.7%             | [ 40.55% - 40.90% ]             |
| % 65+ with leukemias and lymphomas                      |   | 2.9%                  | [ 2.33% - 3.39% ]               | 2.7%              | [ 2.60% - 2.71% ]               |
| % 65+ with liver disease                                |   | 10.5%                 | [ 9.52% - 11.45% ]              | 9.5%              | [ 9.37% - 9.59% ]               |
| % 65+ with lung cancer                                  |   | 2.1%                  | [ 1.68% - 2.53% ]               | 1.9%              | [ 1.81% - 1.90% ]               |
| % 65+ with migraine and other chronic headache          |   | 5.4%                  | [ 4.69% - 6.10% ]               | 5.2%              | [ 5.13% - 5.29% ]               |
| % 65+ with osteoarthritis or rheumatoid arthritis       |   | 53.8%                 | [ 52.17% - 55.41% ]             | 54.2%             | [ 53.97% - 54.34% ]             |
| % 65+ with osteoporosis                                 | W   | 22.2%                 | [ 20.99% - 23.44% ]             | 20.2%             | [ 20.11% - 20.37% ]             |
| % 65+ with peripheral vascular disease                  | W   | 25.6%                 | [ 24.30% - 26.98% ]             | 19.9%             | [ 19.72% - 19.99% ]             |
| % 65+ with pressure ulcer or chronic ulcer              | W   | 11.2%                 | [ 10.20% - 12.10% ]             | 9.2%              | [ 9.11% - 9.31% ]               |
| % 65+ with prostate cancer (men)                        |   | 13.1%                 | [ 11.59% - 14.61% ]             | 13.1%             | [ 12.93% - 13.28% ]             |
| % 65+ with stroke                                       |   | 11.6%                 | [ 10.64% - 12.50% ]             | 11.9%             | [ 11.80% - 12.01% ]             |
| % 65+ with traumatic brain injury                       |   | 1.1%                  | [ 0.83% - 1.47% ]               | 1.3%              | [ 1.25% - 1.33% ]               |
| % 65+ with 4+ (out of 15) chronic conditions            | W   | 64.1%                 | [ 62.61% - 65.58% ]             | 61.8%             | [ 61.61% - 61.95% ]             |
| % 65+ with 0 chronic conditions                         |   | 6.6%                  | [ 5.84% - 7.45% ]               | 7.2%              | [ 7.10% - 7.29% ]               |
| <b>BEHAVIORAL HEALTH</b>                                |   |                       |                                 |                   |                                 |
| % 65+ with opioid use disorder                          |   | 2.1%                  | [ 1.65% - 2.60% ]               | 2.2%              | [ 2.10% - 2.21% ]               |
| % 65+ with substance use disorder                       | B   | 5.3%                  | [ 4.55% - 5.98% ]               | 6.8%              | [ 6.66% - 6.85% ]               |
| % 60+ who used marijuana in last month                  |   | N/A                   |                                 | 2.9%              | [ 2.30% - 3.53% ]               |
| % 60+ excessive drinking                                |   | 7.5%                  | [ 4.27% - 10.78% ]              | 9.2%              | [ 8.64% - 9.71% ]               |
| % 65+ with tobacco use disorder                         | B   | 8.4%                  | [ 7.51% - 9.35% ]               | 10.1%             | [ 9.94% - 10.17% ]              |
| % 60+ current smokers                                   |   | 8.5%                  | [ 5.40% - 11.64% ]              | 7.7%              | [ 7.20% - 8.17% ]               |
| % 60+ ever used E-Cigarettes in last month              |   | 6.5%                  | [ 3.87% - 9.19% ]               | 6.4%              | [ 5.89% - 6.80% ]               |
| <b>MENTAL HEALTH</b>                                    |   |                       |                                 |                   |                                 |
| % 60+ who reported receiving adequate emotional support |   | 78.2%                 | [ 75.03% - 81.46% ]             | 78.3%             | [ 76.70% - 79.88% ]             |
| % 60+ with 15+ days poor mental health in last month    |   | 6.6%                  | [ 4.04% - 9.11% ]               | 6.9%              | [ 6.41% - 7.37% ]               |
| % 65+ with depression                                   |   | 28.8%                 | [ 27.42% - 30.13% ]             | 30.3%             | [ 30.10% - 30.41% ]             |
| % 65+ with anxiety disorder                             | B   | 23.6%                 | [ 22.23% - 24.90% ]             | 25.5%             | [ 25.35% - 25.66% ]             |
| % 65+ with bipolar disorder                             |   | 3.7%                  | [ 3.09% - 4.30% ]               | 3.9%              | [ 3.78% - 3.92% ]               |
| % 65+ with post-traumatic stress disorder               |   | 0.9%                  | [ 0.60% - 1.26% ]               | 1.3%              | [ 1.20% - 1.29% ]               |
| % 65+ with schizophrenia & other psychotic disorder     |   | 4.6%                  | [ 3.99% - 5.31% ]               | 4.6%              | [ 4.53% - 4.68% ]               |
| % 65+ with personality disorder                         |   | 2.4%                  | [ 1.91% - 2.90% ]               | 2.6%              | [ 2.49% - 2.61% ]               |

| HEALTHY AGING INDICATORS   | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>LIVING WITH DISABILITY</b>                                    |   |                       |                                 |                   |                                 |
| % 65+ with self-reported hearing difficulty                      |   | 9.9%                  | [ 7.37% - 12.35% ]              | 12.4%             | [ 12.08% - 12.71% ]             |
| % 65+ with clinical diagnosis of deafness or hearing impairment  |   | 16.2%                 | [ 15.11% - 17.37% ]             | 15.7%             | [ 15.60% - 15.85% ]             |
| % 65+ with self-reported vision difficulty                       |   | 3.3%                  | [ 1.77% - 4.75% ]               | 4.8%              | [ 4.55% - 5.04% ]               |
| % 65+ with clinical diagnosis of blindness or visual impairment  |   | 1.1%                  | [ 0.78% - 1.40% ]               | 1.1%              | [ 1.05% - 1.12% ]               |
| % 65+ with self-reported cognition difficulty                    | *   | 4.4%                  | [ 2.48% - 6.28% ]               | 7.6%              | [ 7.34% - 7.87% ]               |
| % 65+ with self-reported ambulatory difficulty                   | *   | 14.5%                 | [ 11.47% - 17.62% ]             | 19.1%             | [ 18.71% - 19.52% ]             |
| % 65+ with clinical diagnosis of mobility impairment             |   | 3.5%                  | [ 2.90% - 4.03% ]               | 4.0%              | [ 3.95% - 4.09% ]               |
| % 65+ with self-reported self-care difficulty                    | *   | 4.6%                  | [ 2.79% - 6.38% ]               | 7.5%              | [ 7.25% - 7.77% ]               |
| % 65+ with self-reported independent living difficulty           | *   | 9.6%                  | [ 7.07% - 12.13% ]              | 13.6%             | [ 13.23% - 13.93% ]             |
| <b>CAREGIVING</b>  |   |                       |                                 |                   |                                 |
| % of 60+ who provide care to a family/friend in last month       |   | 18.5%                 | [ 15.05% - 22.05% ]             | 21.7%             | [ 19.87% - 23.57% ]             |
| % of grandparents raising grandchildren                          |   | 0.7%                  | [ 0.20% - 1.11% ]               | 0.8%              | [ 0.73% - 0.86% ]               |
| % of grandparents who live with grandchildren                    | *   | 1.7%                  | [ 1.00% - 2.39% ]               | 2.7%              | [ 2.62% - 2.83% ]               |
| <b>ACCESS TO CARE</b>  |   |                       |                                 |                   |                                 |
| % 65+ dually eligible for Medicare and Medicaid                  | *   | 18.3%                 | [ 17.50% - 19.14% ]             | 22.1%             | [ 21.96% - 22.17% ]             |
| % 65+ Medicare managed care enrollees                            | *   | 37.5%                 | [ 36.51% - 38.56% ]             | 29.9%             | [ 29.76% - 29.99% ]             |
| % 60+ with a regular doctor                                      |   | 96.0%                 | [ 93.69% - 98.24% ]             | 96.0%             | [ 95.68% - 96.39% ]             |
| % 60+ who did not see a doctor when needed due to cost           |   | 4.8%                  | [ 2.47% - 7.15% ]               | 4.5%              | [ 4.14% - 4.92% ]               |
| <b>SERVICE UTILIZATION</b>                                       |   |                       |                                 |                   |                                 |
| # physician visits per year                                      | *   | 9.2                   | [ 8.90 - 9.43 ]                 | 8.6               | [ 8.60 - 8.66 ]                 |
| # emergency room visits/1000 persons 65+ years annually          | *   | 558.0                 | [ 516.30 - 599.75 ]             | 636.7             | [ 631.77 - 641.56 ]             |
| # Part D monthly prescription fills per person annually          | *   | 48.5                  | [ 47.53 - 49.39 ]               | 50.3              | [ 50.15 - 50.39 ]               |
| # home health visits annually                                    |   | 4.6                   | [ 4.00 - 5.25 ]                 | 4.1               | [ 4.02 - 4.14 ]                 |
| # durable medical equipment claims annually                      |   | 1.8                   | [ 1.61 - 1.94 ]                 | 1.9               | [ 1.88 - 1.91 ]                 |
| # inpatient hospital stays/1000 persons 65+ years annually       |   | 282.1                 | [ 257.62 - 306.54 ]             | 273.3             | [ 270.72 - 275.80 ]             |
| % Medicare inpatient hospital readmissions (as % of admissions)  |   | 18.2%                 | [ 15.08% - 21.37% ]             | 17.2%             | [ 16.89% - 17.47% ]             |
| # skilled nursing facility stays/1000 persons 65+ years annually |   | 113.2                 | [ 99.41 - 127.07 ]              | 104.7             | [ 103.23 - 106.08 ]             |
| % 65+ getting Medicaid long term services and supports           |   | 5.4%                  | [ 4.87% - 5.83% ]               | 5.5%              | [ 5.48% - 5.60% ]               |
| % 65+ hospice users  |   | 3.0%                  | [ 2.58% - 3.34% ]               | 2.8%              | [ 2.70% - 2.79% ]               |
| % 65+ hospice users as % of decedents                            |   | 51.2%                 | [ 46.13% - 56.32% ]             | 45.5%             | [ 44.85% - 46.14% ]             |
| Median hospice days per hospice user (65+, deceased)             | *   | 7.0                   | [ 6.0 - 9.0 ]                   | 10.0              | [ 10.00 - 10.00 ]               |
| Median hospice payment (Medicare + other) per hospice user       |   | \$3,498               | [ \$2,943 - \$4,039 ]           | \$3,741           | [ \$3,650 - \$3,816 ]           |
| <b>COMMUNITY</b>   |   |                       |                                 |                   |                                 |
| % 60+ who used Internet in last month                            |   | 74.0%                 | [ 68.89% - 79.19% ]             | 75.1%             | [ 74.32% - 75.89% ]             |
| <b>TRANSPORTATION</b>  |   |                       |                                 |                   |                                 |
| % 65+ who own a motor vehicle                                    | *   | 90.5%                 | [ 88.10% - 92.81% ]             | 87.3%             | [ 87.15% - 87.49% ]             |
| % 60+ who always drive or ride wearing a seatbelt                |   | 91.2%                 | [ 87.72% - 94.62% ]             | 93.0%             | [ 92.53% - 93.49% ]             |
| % 60+ who drove under the influence of drinking in last month    |   | N/A                   |                                 | 2.2%              | [ 1.76% - 2.56% ]               |

| HEALTHY AGING INDICATORS                                  | HIGHER /<br>LOWER<br>STATE<br>RATE <sup>1</sup> | COMMUNITY<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> | STATE<br>ESTIMATE | MARGIN OF<br>ERROR <sup>2</sup> |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| <b>HOUSING</b>  |   |                       |                                 |                   |                                 |
| % 65+ population who live alone                           |   | 26.4%                 | [ 22.33% - 30.42% ]             | 28.4%             | [ 27.99% - 28.89% ]             |
| % 60+ own home  | *   | 87.7%                 | [ 85.24% - 90.07% ]             | 76.9%             | [ 76.70% - 77.17% ]             |
| % 60+ homeowners who have mortgage                        |   | 47.8%                 | [ 42.83% - 52.76% ]             | 47.1%             | [ 46.58% - 47.64% ]             |
| % 60+ stressed about paying rent/mortgage in last month   |   | 20.0%                 | [ 14.45% - 25.63% ]             | 17.4%             | [ 16.52% - 18.19% ]             |
| % 65+ households (renter) spend >35% of income on housing |   | 41.2%                 | [ 27.78% - 54.72% ]             | 44.0%             | [ 42.55% - 45.42% ]             |
| % 65+ households (owner) spend >35% of income on housing  |   | 33.3%                 | [ 27.17% - 39.37% ]             | 29.5%             | [ 28.89% - 30.15% ]             |
| % 65+ moved within same county in last year               |   | 2.7%                  | [ 1.36% - 4.05% ]               | 3.7%              | [ 3.48% - 3.89% ]               |
| % 65+ moved from different county in last year            |   | 1.4%                  | [ -0.02% - 2.76% ]              | 0.8%              | [ 0.71% - 0.90% ]               |
| % 65+ moved from different state in last year             |   | 0.3%                  | [ -0.37% - 1.00% ]              | 0.9%              | [ 0.82% - 1.05% ]               |
| % of vacant homes in community                            | *   | 6.1%                  | [ 4.35% - 7.87% ]               | 9.6%              | [ 9.29% - 9.87% ]               |
| <b>ECONOMIC</b>   |   |                       |                                 |                   |                                 |
| % 60+ receiving food benefits in last year                | *   | 5.9%                  | [ 3.46% - 8.37% ]               | 10.9%             | [ 10.53% - 11.21% ]             |
| % 65+ employed in last year                               |   | 27.4%                 | [ 23.99% - 30.78% ]             | 24.8%             | [ 24.48% - 25.15% ]             |
| % 65+ with income below the poverty line in last year     | *   | 4.2%                  | [ 2.31% - 6.15% ]               | 7.0%              | [ 6.79% - 7.30% ]               |
| % 65+ households with annual income < \$20,000            |   | 14.2%                 | [ 9.99% - 18.51% ]              | 17.1%             | [ 16.57% - 17.55% ]             |
| % 65+ households with annual income \$20,000-\$49,999     |   | 34.0%                 | [ 27.66% - 40.37% ]             | 31.0%             | [ 30.35% - 31.62% ]             |
| % 65+ households with annual income \$50,000-\$99,999     |   | 31.3%                 | [ 26.65% - 35.88% ]             | 27.6%             | [ 27.04% - 28.17% ]             |
| % 65+ households with annual income \$100,000+            |   | 20.5%                 | [ 16.17% - 24.77% ]             | 24.4%             | [ 23.82% - 24.89% ]             |

Notes. 1 We use B, W, or \* to denote significant differences in rates where the margins of error do not overlap (B=rate is better than the comparison, W=rate is worse than comparison, \*=rate is different than comparison).

2 All community and state margins of error are 95% confidence intervals.

Questions or suggestions? [Beth.Dugan@umb.edu](mailto:Beth.Dugan@umb.edu)