

Newtown (Fairfield)



| HEALTHY AGING INDICATORS | HIGHER / LOWER STATE RATE ¹ | COMMUNITY ESTIMATE | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| POPULATION CHARACTERISTICS | | | | | |
| Population 65 years or older | | | | | |
| % 65-74 years | * | 50.8% | [46.67% - 54.85%] | 55.7% | [55.33% - 56.11%] |
| % 75-84 years | | 31.2% | [27.14% - 35.32%] | 29.1% | [28.75% - 29.45%] |
| % 85 years or older | | 18.0% | [12.84% - 23.18%] | 15.2% | [14.93% - 15.43%] |
| % 65+ population who are female | | 55.6% | [50.21% - 60.91%] | 56.8% | [56.36% - 57.18%] |
| % 85+ population who are female | | 68.9% | [50.75% - 87.07%] | 66.6% | [65.94% - 67.18%] |
| Race and ethnicity of the population 65+ | | | | | |
| % White | * | 98.8% | [96.41% - 101.22%] | 88.4% | [87.85% - 88.93%] |
| % African American | * | 0.2% | [-0.92% - 1.36%] | 6.8% | [6.63% - 6.88%] |
| % Asian | | 1.0% | [-0.52% - 2.44%] | 2.3% | [2.20% - 2.35%] |
| % Hispanic/Latino | * | 0.8% | [-0.59% - 2.25%] | 5.8% | [5.66% - 5.84%] |
| Marital status of the population 65+ | | | | | |
| % married | * | 64.6% | [61.35% - 67.77%] | 53.8% | [53.46% - 54.12%] |
| % divorced/separated | * | 8.1% | [5.28% - 10.94%] | 14.9% | [14.58% - 15.29%] |
| % widowed | | 23.0% | [18.59% - 27.42%] | 24.2% | [23.77% - 24.57%] |
| % never married | * | 4.3% | [2.67% - 5.98%] | 7.1% | [6.85% - 7.36%] |
| Education of the population 65+ | | | | | |
| % with less than high school education | * | 6.4% | [3.69% - 9.17%] | 14.7% | [14.30% - 15.04%] |
| % with high school or some college | | 49.9% | [44.97% - 54.90%] | 53.3% | [52.80% - 53.74%] |
| % with college degree | * | 43.6% | [40.66% - 46.62%] | 32.1% | [31.72% - 32.40%] |
| % with graduate or professional degree | * | 22.6% | [19.58% - 25.67%] | 16.8% | [16.58% - 17.09%] |
| % of 65+ population who speak only English at home | * | 93.8% | [86.79% - 100.80%] | 84.3% | [83.76% - 84.83%] |
| % of 65+ population who are veterans of military service | | 19.4% | [15.76% - 22.95%] | 17.5% | [17.21% - 17.69%] |
| % of 60+ who are LGBT (County) | | 2.1% | [1.50% - 2.60%] | 2.5% | [2.17% - 2.74%] |
| Age-sex adjusted 1-year mortality rate | | 3.7% | [3.16% - 4.32%] | 4.1% | [4.08% - 4.18%] |
| WELLNESS | | | | | |
| % 60+ getting the recommended hours of sleep | | 68.3% | [62.70% - 73.85%] | 65.2% | [64.24% - 66.16%] |
| % 60+ doing any physical activity within last month | B | 81.0% | [77.07% - 84.92%] | 75.0% | [74.20% - 75.79%] |
| % 60+ met CDC guidelines for muscle-strengthening activity | | 30.1% | [24.66% - 35.48%] | 27.6% | [26.62% - 28.52%] |
| % 60+ met CDC guidelines for aerobic physical activity | | 62.4% | [56.52% - 68.19%] | 58.1% | [56.98% - 59.11%] |
| % 60+ with fair or poor health status | B | 12.9% | [9.53% - 16.30%] | 17.6% | [16.90% - 18.31%] |
| % 60+ with 15+ physically unhealthy days in last month | | 10.7% | [7.35% - 14.05%] | 12.4% | [11.74% - 12.96%] |

| HEALTHY AGING INDICATORS | HIGHER / LOWER STATE RATE ¹ | COMMUNITY ESTIMATE | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| FALLS | | | | | |
| % 60+ who fell within last year | | 25.4% | [20.26% - 30.59%] | 26.3% | [25.38% - 27.18%] |
| % 60+ who were injured in a fall within last year | | 8.5% | [5.34% - 11.58%] | 9.8% | [9.22% - 10.46%] |
| % 65+ had hip fracture | | 4.0% | [3.31% - 4.62%] | 3.7% | [3.63% - 3.75%] |
| PREVENTION | | | | | |
| % 60+ with physical exam/check-up in last year | | 84.9% | [81.17% - 88.65%] | 88.5% | [87.93% - 89.09%] |
| % 60+ flu shot in last year | | 63.3% | [58.34% - 68.35%] | 57.5% | [56.60% - 58.43%] |
| % 60+ with pneumonia vaccine | | 60.5% | [55.13% - 65.88%] | 63.9% | [62.97% - 64.79%] |
| % 60+ with shingles vaccine | | 35.4% | [28.83% - 41.87%] | 33.5% | [32.43% - 34.52%] |
| % 60+ women with a mammogram within last 2 years | | 78.3% | [70.99% - 85.61%] | 79.7% | [78.56% - 80.79%] |
| % 60+ with colorectal cancer screening | | 79.6% | [74.40% - 84.80%] | 78.2% | [77.29% - 79.05%] |
| % 60+ with HIV test | | 13.2% | [9.54% - 16.88%] | 16.1% | [15.40% - 16.83%] |
| % 60+ met CDC preventive health screening goals | | 36.7% | [29.78% - 43.57%] | 39.6% | [38.38% - 40.80%] |
| NUTRITION & DIET | | | | | |
| % 60+ with 5 or more servings of fruit or vegetables per day | | 23.8% | [18.80% - 28.86%] | 19.9% | [19.00% - 20.72%] |
| % 60+ stressed about buying food in last month | | 8.1% | [4.89% - 11.34%] | 8.6% | [7.95% - 9.17%] |
| % 60+ self-reported obese | | 22.0% | [17.55% - 26.41%] | 26.1% | [25.27% - 26.93%] |
| % 65+ clinically diagnosed obesity | W | 24.6% | [22.89% - 26.33%] | 22.5% | [22.35% - 22.66%] |
| % 65+ with high cholesterol | | 78.1% | [76.54% - 79.72%] | 77.0% | [76.85% - 77.15%] |
| % 60+ with cholesterol screening | | 97.9% | [96.45% - 99.33%] | 97.1% | [96.74% - 97.42%] |
| ORAL HEALTH | | | | | |
| % 60+ with dental insurance | | 58.3% | [48.55% - 67.99%] | 59.2% | [57.60% - 60.76%] |
| % 60+ with annual dental exam | | 83.6% | [79.18% - 87.93%] | 80.0% | [79.13% - 80.76%] |
| % 60+ with loss of 6 or more teeth | | 21.3% | [16.28% - 26.29%] | 23.8% | [22.96% - 24.69%] |
| CHRONIC DISEASE | | | | | |
| % 65+ with Alzheimer's disease or related dementias | B | 13.0% | [11.75% - 14.17%] | 14.4% | [14.26% - 14.49%] |
| % 65+ with anemia | | 49.5% | [47.63% - 51.33%] | 50.6% | [50.41% - 50.76%] |
| % 65+ with asthma | W | 16.8% | [15.45% - 18.12%] | 14.1% | [13.96% - 14.19%] |
| % 65+ with atrial fibrillation | B | 14.6% | [13.40% - 15.89%] | 16.1% | [15.94% - 16.18%] |
| % 65+ with autism spectrum disorder | | 0.07% | [0.03% - 0.11%] | 0.12% | [0.11% - 0.14%] |
| % 65+ with benign prostatic hyperplasia (men) | | 45.4% | [42.65% - 48.25%] | 42.8% | [42.49% - 43.01%] |
| % 65+ with breast cancer (women) | | 11.2% | [9.70% - 12.72%] | 11.4% | [11.25% - 11.53%] |
| % 65+ with cataract | | 64.0% | [62.17% - 65.83%] | 64.8% | [64.67% - 65.00%] |
| % 65+ with chronic kidney disease | | 28.5% | [26.79% - 30.24%] | 29.9% | [29.70% - 30.03%] |
| % 65+ with chronic obstructive pulmonary disease | | 19.6% | [18.15% - 20.97%] | 21.0% | [20.90% - 21.17%] |
| % 65+ with colon cancer | | 2.2% | [1.72% - 2.72%] | 2.7% | [2.65% - 2.75%] |
| % 65+ with congestive heart failure | B | 16.7% | [15.32% - 18.00%] | 22.6% | [22.43% - 22.72%] |
| % 65+ with diabetes | B | 31.7% | [29.89% - 33.53%] | 33.8% | [33.59% - 33.93%] |
| % 65+ with endometrial cancer (women) | | 2.0% | [1.37% - 2.72%] | 2.0% | [1.97% - 2.10%] |

| HEALTHY AGING INDICATORS | HIGHER / LOWER STATE RATE ¹ | COMMUNITY ESTIMATE | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| % 65+ with epilepsy | | 2.7% | [2.07% - 3.40%] | 2.9% | [2.82% - 2.95%] |
| % 65+ with fibromyalgia, chronic pain, and fatigue | W | 31.3% | [29.51% - 33.12%] | 26.6% | [26.42% - 26.73%] |
| % 65+ with glaucoma | | 26.6% | [24.98% - 28.14%] | 28.3% | [28.11% - 28.41%] |
| % 65+ ever had a heart attack | B | 3.6% | [2.95% - 4.22%] | 4.5% | [4.40% - 4.53%] |
| % 65+ with HIV/AIDS | * | 0.14% | [0.08% - 0.19%] | 0.21% | [0.19% - 0.23%] |
| % 65+ with hypertension | | 74.4% | [72.73% - 76.09%] | 76.2% | [76.07% - 76.37%] |
| % 65+ with hypothyroidism | | 25.8% | [24.18% - 27.33%] | 25.1% | [24.99% - 25.28%] |
| % 65+ with ischemic heart disease | | 39.1% | [37.21% - 40.99%] | 40.7% | [40.55% - 40.90%] |
| % 65+ with leukemias and lymphomas | | 2.3% | [1.77% - 2.91%] | 2.7% | [2.60% - 2.71%] |
| % 65+ with liver disease | | 10.4% | [9.24% - 11.64%] | 9.5% | [9.37% - 9.59%] |
| % 65+ with lung cancer | | 1.6% | [1.10% - 2.03%] | 1.9% | [1.81% - 1.90%] |
| % 65+ with migraine and other chronic headache | | 6.0% | [5.04% - 6.90%] | 5.2% | [5.13% - 5.29%] |
| % 65+ with osteoarthritis or rheumatoid arthritis | | 55.8% | [53.81% - 57.82%] | 54.2% | [53.97% - 54.34%] |
| % 65+ with osteoporosis | | 20.7% | [19.27% - 22.12%] | 20.2% | [20.11% - 20.37%] |
| % 65+ with peripheral vascular disease | W | 23.5% | [21.90% - 25.06%] | 19.9% | [19.72% - 19.99%] |
| % 65+ with pressure ulcer or chronic ulcer | | 9.7% | [8.63% - 10.78%] | 9.2% | [9.11% - 9.31%] |
| % 65+ with prostate cancer (men) | W | 15.2% | [13.32% - 17.18%] | 13.1% | [12.93% - 13.28%] |
| % 65+ with stroke | | 11.3% | [10.17% - 12.37%] | 11.9% | [11.80% - 12.01%] |
| % 65+ with traumatic brain injury | | 1.5% | [1.01% - 1.98%] | 1.3% | [1.25% - 1.33%] |
| % 65+ with 4+ (out of 15) chronic conditions | | 59.8% | [57.95% - 61.65%] | 61.8% | [61.61% - 61.95%] |
| % 65+ with 0 chronic conditions | | 6.9% | [5.84% - 7.87%] | 7.2% | [7.10% - 7.29%] |
| BEHAVIORAL HEALTH | | | | | |
| % 65+ with opioid use disorder | | 2.8% | [2.12% - 3.42%] | 2.2% | [2.10% - 2.21%] |
| % 65+ with substance use disorder | B | 5.6% | [4.68% - 6.48%] | 6.8% | [6.66% - 6.85%] |
| % 60+ who used marijuana in last month | | N/A | | 2.9% | [2.30% - 3.53%] |
| % 60+ excessive drinking | | 8.9% | [5.88% - 11.85%] | 9.2% | [8.64% - 9.71%] |
| % 65+ with tobacco use disorder | B | 8.1% | [6.98% - 9.17%] | 10.1% | [9.94% - 10.17%] |
| % 60+ current smokers | B | 4.3% | [2.35% - 6.29%] | 7.7% | [7.20% - 8.17%] |
| % 60+ ever used E-Cigarettes in last month | | 5.5% | [3.11% - 7.93%] | 6.4% | [5.89% - 6.80%] |
| MENTAL HEALTH | | | | | |
| % 60+ who reported receiving adequate emotional support | | 78.2% | [75.03% - 81.46%] | 78.3% | [76.70% - 79.88%] |
| % 60+ with 15+ days poor mental health in last month | | 8.3% | [5.20% - 11.30%] | 6.9% | [6.41% - 7.37%] |
| % 65+ with depression | | 29.7% | [28.05% - 31.38%] | 30.3% | [30.10% - 30.41%] |
| % 65+ with anxiety disorder | | 25.2% | [23.52% - 26.89%] | 25.5% | [25.35% - 25.66%] |
| % 65+ with bipolar disorder | | 3.2% | [2.49% - 3.85%] | 3.9% | [3.78% - 3.92%] |
| % 65+ with post-traumatic stress disorder | | 0.8% | [0.46% - 1.21%] | 1.3% | [1.20% - 1.29%] |
| % 65+ with schizophrenia & other psychotic disorder | B | 2.9% | [2.29% - 3.53%] | 4.6% | [4.53% - 4.68%] |
| % 65+ with personality disorder | | 2.3% | [1.76% - 2.91%] | 2.6% | [2.49% - 2.61%] |

| HEALTHY AGING INDICATORS | HIGHER / LOWER STATE RATE ¹ | COMMUNITY ESTIMATE | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|--|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| LIVING WITH DISABILITY | | | | | |
| % 65+ with self-reported hearing difficulty | | 13.0% | [8.85% - 17.13%] | 12.4% | [12.08% - 12.71%] |
| % 65+ with clinical diagnosis of deafness or hearing impairment | | 16.8% | [15.37% - 18.17%] | 15.7% | [15.60% - 15.85%] |
| % 65+ with self-reported vision difficulty | | 3.8% | [2.23% - 5.34%] | 4.8% | [4.55% - 5.04%] |
| % 65+ with clinical diagnosis of blindness or visual impairment | W | 1.7% | [1.21% - 2.11%] | 1.1% | [1.05% - 1.12%] |
| % 65+ with self-reported cognition difficulty | | 9.1% | [5.13% - 13.14%] | 7.6% | [7.34% - 7.87%] |
| % 65+ with self-reported ambulatory difficulty | | 16.3% | [12.12% - 20.47%] | 19.1% | [18.71% - 19.52%] |
| % 65+ with clinical diagnosis of mobility impairment | | 4.8% | [3.98% - 5.64%] | 4.0% | [3.95% - 4.09%] |
| % 65+ with self-reported self-care difficulty | | 9.2% | [5.39% - 13.07%] | 7.5% | [7.25% - 7.77%] |
| % 65+ with self-reported independent living difficulty | | 16.6% | [11.65% - 21.54%] | 13.6% | [13.23% - 13.93%] |
| CAREGIVING | | | | | |
| % of 60+ who provide care to a family/friend in last month | | 18.5% | [15.05% - 22.05%] | 21.7% | [19.87% - 23.57%] |
| % of grandparents raising grandchildren | | 0.9% | [0.21% - 1.50%] | 0.8% | [0.73% - 0.86%] |
| % of grandparents who live with grandchildren | | 2.6% | [1.52% - 3.74%] | 2.7% | [2.62% - 2.83%] |
| ACCESS TO CARE | | | | | |
| % 65+ dually eligible for Medicare and Medicaid | * | 11.9% | [10.98% - 12.88%] | 22.1% | [21.96% - 22.17%] |
| % 65+ Medicare managed care enrollees | * | 20.4% | [19.19% - 21.55%] | 29.9% | [29.76% - 29.99%] |
| % 60+ with a regular doctor | | 95.7% | [93.46% - 97.86%] | 96.0% | [95.68% - 96.39%] |
| % 60+ who did not see a doctor when needed due to cost | | 6.4% | [3.65% - 9.10%] | 4.5% | [4.14% - 4.92%] |
| SERVICE UTILIZATION | | | | | |
| # physician visits per year | * | 10.1 | [9.74 - 10.43] | 8.6 | [8.60 - 8.66] |
| # emergency room visits/1000 persons 65+ years annually | * | 445.0 | [406.70 - 483.29] | 636.7 | [631.77 - 641.56] |
| # Part D monthly prescription fills per person annually | * | 47.2 | [45.86 - 48.45] | 50.3 | [50.15 - 50.39] |
| # home health visits annually | | 4.1 | [3.27 - 4.96] | 4.1 | [4.02 - 4.14] |
| # durable medical equipment claims annually | * | 1.6 | [1.43 - 1.84] | 1.9 | [1.88 - 1.91] |
| # inpatient hospital stays/1000 persons 65+ years annually | * | 223.4 | [200.70 - 246.05] | 273.3 | [270.72 - 275.80] |
| % Medicare inpatient hospital readmissions (as % of admissions) | * | 13.7% | [10.72% - 16.73%] | 17.2% | [16.89% - 17.47%] |
| # skilled nursing facility stays/1000 persons 65+ years annually | | 94.2 | [80.48 - 107.83] | 104.7 | [103.23 - 106.08] |
| % 65+ getting Medicaid long term services and supports | * | 3.1% | [2.56% - 3.56%] | 5.5% | [5.48% - 5.60%] |
| % 65+ hospice users | | 2.6% | [2.12% - 3.09%] | 2.8% | [2.70% - 2.79%] |
| % 65+ hospice users as % of decedents | | 47.9% | [35.85% - 59.92%] | 45.5% | [44.85% - 46.14%] |
| Median hospice days per hospice user (65+, deceased) | | 11.0 | [9.0 - 16.0] | 10.0 | [10.00 - 10.00] |
| Median hospice payment (Medicare + other) per hospice user | | \$3,410 | [\$2,362 - \$4,439] | \$3,741 | [\$3,650 - \$3,816] |
| COMMUNITY | | | | | |
| % 60+ who used Internet in last month | * | 85.5% | [82.15% - 88.81%] | 75.1% | [74.32% - 75.89%] |
| TRANSPORTATION | | | | | |
| % 65+ who own a motor vehicle | * | 92.1% | [90.43% - 93.79%] | 87.3% | [87.15% - 87.49%] |
| % 60+ who always drive or ride wearing a seatbelt | | 94.7% | [92.56% - 96.94%] | 93.0% | [92.53% - 93.49%] |
| % 60+ who drove under the influence of drinking in last month | | N/A | | 2.2% | [1.76% - 2.56%] |

| HEALTHY AGING INDICATORS | HIGHER / LOWER STATE RATE ¹ | COMMUNITY ESTIMATE | MARGIN OF ERROR ² | STATE ESTIMATE | MARGIN OF ERROR ² |
|---|---|-----------------------|---------------------------------|-------------------|---------------------------------|
| HOUSING | | | | | |
| % 65+ population who live alone | | 25.8% | [20.56% - 31.01%] | 28.4% | [27.99% - 28.89%] |
| % 60+ own home | * | 83.6% | [80.71% - 86.59%] | 76.9% | [76.70% - 77.17%] |
| % 60+ homeowners who have mortgage | | 44.7% | [39.10% - 50.28%] | 47.1% | [46.58% - 47.64%] |
| % 60+ stressed about paying rent/mortgage in last month | | 17.3% | [12.60% - 22.02%] | 17.4% | [16.52% - 18.19%] |
| % 65+ households (renter) spend >35% of income on housing | | 60.9% | [34.30% - 87.41%] | 44.0% | [42.55% - 45.42%] |
| % 65+ households (owner) spend >35% of income on housing | | 29.3% | [21.98% - 36.62%] | 29.5% | [28.89% - 30.15%] |
| % 65+ moved within same county in last year | | 2.3% | [-0.11% - 4.77%] | 3.7% | [3.48% - 3.89%] |
| % 65+ moved from different county in last year | | 0.4% | [-0.66% - 1.56%] | 0.8% | [0.71% - 0.90%] |
| % 65+ moved from different state in last year | | 1.2% | [0.26% - 2.15%] | 0.9% | [0.82% - 1.05%] |
| % of vacant homes in community | * | 6.4% | [4.18% - 8.64%] | 9.6% | [9.29% - 9.87%] |
| ECONOMIC | | | | | |
| % 60+ receiving food benefits in last year | * | 2.8% | [1.06% - 4.48%] | 10.9% | [10.53% - 11.21%] |
| % 65+ employed in last year | | 21.7% | [18.12% - 25.34%] | 24.8% | [24.48% - 25.15%] |
| % 65+ with income below the poverty line in last year | | 4.5% | [1.97% - 7.08%] | 7.0% | [6.79% - 7.30%] |
| % 65+ households with annual income < \$20,000 | * | 10.3% | [5.49% - 15.16%] | 17.1% | [16.57% - 17.55%] |
| % 65+ households with annual income \$20,000-\$49,999 | | 34.3% | [26.85% - 41.80%] | 31.0% | [30.35% - 31.62%] |
| % 65+ households with annual income \$50,000-\$99,999 | | 22.6% | [17.52% - 27.64%] | 27.6% | [27.04% - 28.17%] |
| % 65+ households with annual income \$100,000+ | * | 32.8% | [26.44% - 39.08%] | 24.4% | [23.82% - 24.89%] |

Notes. 1 We use B, W, or * to denote significant differences in rates where the margins of error do not overlap (B=rate is better than the comparison, W=rate is worse than comparison, *=rate is different than comparison).

2 All community and state margins of error are 95% confidence intervals.

Questions or suggestions? Beth.Dugan@umb.edu